

INEQUALITIES FOR TRIGONOMETRIC SUMS IN TWO VARIABLES

V. E. S. SZABÓ

Abstract. We prove sharp inequalities for the two-variable versions of the sum $\sum_{k=1}^n \frac{\sin(kx)}{k}$.

Mathematics subject classification (2020): 26D05, 42A05.

Keywords and phrases: Trigonometric sum, two variables, inequalities.

REFERENCES

- [1] H. ALZER, S. KOUMANDOS, *Sharp inequalities for trigonometric sums*, Math. Proc. Cambridge Philos. Soc. **134**, 1 (2003), 139–52.
- [2] H. ALZER, S. KOUMANDOS, *Inequalities of Fejér-Jackson Type*, Monatsh. Math. **139**, 2 (2003), 89–103.
- [3] H. ALZER, X. SHI, *Sharp bounds for trigonometric polynomials in two variables*, Anal. and Appl. **7**, 4 (2009), 341–350.
- [4] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER, F. G. TRICOMI, *Higher Transcendental Functions, Vol. I*, McGraw-Hill, New York, 1953.
- [5] D. JACKSON, *Über eine trigonometrische Summe*, Rend. Circ. Mat. Palermo **32**, 1 (1911), 257–262.
- [6] L. KOSCHMIEDER, *Vorzeicheneigenschaften der Abschnitte einiger physikalisch bedeutsamer Reihen*, Monatsh. Math. Phys. **39**, 1 (1932), 321–344.
- [7] G. V. MILOVANOVIĆ, D. S. MITRINOVIĆ, TH. M. RASSIAS, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Sci. Publ., Singapore, 1994.
- [8] P. TURÁN, *Über die Partialsummen der Fourierreihe*, J. London. Math. Soc. **13**, 1 (1938), 278–282.