

ASYMPTOTICS FOR GENERATING FUNCTIONS OF THE FUSS–CATALAN NUMBERS

ASSIS AZEVEDO* AND DAVIDE AZEVEDO

(Communicated by S. Varošanec)

Abstract. We consider a certain class of polynomials with coefficients in \mathbb{Z}_M , all of which admit a unique zero. We prove that the zero of each of those can be given by a (multiple) sum involving the coefficients and a vectorial generalization of the Fuss-Catalan numbers.

We also consider the sequence of the partial sums of the generating function of the d -Fuss-Catalan numbers. Using the holonomy of this sequence, we study its asymptotic behaviour. The main difference from the known case $d = 2$ is, in that one, we have a “closed” expression for the generating function.

1. Introduction

The Catalan numbers were studied by Euler, in the context of enumerating triangulations of regular polygons [5]. Their study by the Mongolian mathematician Antu Ming in the eighteenth century was announced in 1988 by Luo in [10] and further discussed by Larcombe in [9].

These numbers have multiple interpretations and applications, several of which can be found, for example, in [18], which also covers different generalizations of them. Throughout this paper we focus on a couple of these, the d -Fuss-Catalan numbers, for $d \in \mathbb{N} \setminus \{1\}$, whose element of order n , $C_d(n)$, is defined by

$$C_d(n) = \frac{1}{(d-1)n+1} \binom{dn}{n}, \quad (1)$$

and a vectorial generalization of the Catalan numbers, which we will define in (4). $C_d(n)$, introduced by Fuss in [6], counts, for example, the number of partitions of a $n(d-1)+2$ -gon into $d+1$ -gons and the number of d -ary trees with n internal nodes (see [7]). Recall that the Catalan numbers are the 2-Fuss-Catalan numbers.

The first problem we are interested in is finding the zeros of some polynomials in \mathbb{Z}_M , the ring of the integers modulo $M \in \mathbb{N}$. Consider a polynomial $Q = Q(x)$ with coefficients in \mathbb{Z}_M of the form

$$a_d x^d + \cdots + a_1 x + a_0, \quad \text{where } a_i \text{ is nilpotent for } i \geq 2 \text{ and } a_1 \text{ invertible.} \quad (2)$$

Mathematics subject classification (2020): 41A60, 05A10.

Keywords and phrases: Fuss-Catalan numbers, asymptotic behavior, holonomic sequence, polynomial congruence, vectorial Catalan numbers.

* Corresponding author.

The Chinese remainder theorem and the Hensel lemma guarantee that there exists exactly one zero of Q in \mathbb{Z}_M . In this work, we will find a polynomial P in $d + 1$ variables such that the zero of any polynomial as in (2) is equal to $P(a_0, a_1^{-1}, a_2, \dots, a_d)$. The coefficients of P are essentially vector generalized Catalan numbers, which are d -Fuss-Catalan numbers if $a_i = 0$ for $1 < i < d$.

The second problem was motivated by sequences presented in OEIS, The On-Line Encyclopedia of Integer Sequences [17]. For $d \in \mathbb{N} \setminus \{1\}$, $r \in \mathbb{R} \setminus \{0\}$, and $n \in \mathbb{N}$, consider the sequence

$$X(d, r, n) = \sum_{k=0}^n C_d(k)r^k. \tag{3}$$

In connection with the first problem, we will see that, if p is a prime number and r a multiple of p then, $X(d, r, n)$ is the zero, in $\mathbb{Z}_{p^{n+1}}$ of the polynomial $rx^d - x + 1$.

OEIS, in the sequence <https://oeis.org/A112696> and onwards, presents recurrence formulas for $(X(2, r, n))_{n \in \mathbb{N}}$ for some values of r , conjecturing them for some others. In this work, we obtain recurrence formulas for all values of d and r .

We also study the asymptotic behaviour of this sequence, when it diverges. For $d = 2$, this was done by Mattarei in [11], using, among other instruments, the generating function of the Catalan numbers $F_2(x) = \frac{1 - \sqrt{1 - 4x}}{2x}$. Elezović, in [3, 4] gives an efficient algorithm for recursive calculations of asymptotic expansions of several sums including $X(2, 1, n)$. If $d > 2$ we do not have a nice expression for $F_d(x)$, apart from the equality $F_d(x) = 1 + xF_d(x)^d$.

We use some well-known results for holonomic sequences such as the Poincaré-Perron Theorem in [13, 12], and Corollary 1.6 of [8] to prove that

$$X(d, r, n) \sim \frac{1}{\sqrt{2\pi}} \frac{\sqrt{d}}{(d-1)^{\frac{3}{2}}} \frac{A(d)r}{A(d)r-1} (A(d)r)^n n^{-\frac{3}{2}},$$

where $A(d) = \frac{d^d}{(d-1)^{d-1}}$, and $A(d)|r| > 1$.

2. Preliminaries

The Catalan numbers have a lot of generalizations. In this work we are interested in the d -Fuss-Catalan numbers, defined in (1), and the natural vectorial generalization, $C_{\vec{v}}(\vec{n})$, seen, for example, in [2] and a more general case in [14]. $C_{\vec{v}}(\vec{n})$ is defined by

$$C_{\vec{v}}(\vec{n}) = \frac{1}{(\vec{v}-\vec{1}) \cdot \vec{n} + 1} \binom{\vec{v} \cdot \vec{n}}{\vec{n}} = \frac{1}{\vec{v} \cdot \vec{n} + 1} \binom{\vec{v} \cdot \vec{n} + 1}{\vec{n}} \tag{4}$$

where, given $s \in \mathbb{N}$, $\vec{n} \in \mathbb{N}_0^s$ and $\vec{v} \in \mathbb{N}^s$, $\vec{v} \cdot \vec{n}$ denotes the inner product of \vec{n} and \vec{v} and $\binom{\vec{v} \cdot \vec{n}}{\vec{n}}$ is the multinomial coefficient $\frac{(\vec{v} \cdot \vec{n})!}{n_1! \dots n_s! (\vec{v} \cdot \vec{n} - (n_1 + \dots + n_s))!}$.

$C_{\vec{v}}(\vec{n})$ is, for example, the number of ways that $\vec{v} \cdot \vec{n}$ people can be seated at a (round) table in such a way that, for all $i = 1, \dots, s$, there exist n_i groups of v_i people giving a v_i -hand shake with no crossings between different groups [2]. Of course, this




Figure 1: This is one of the 92810 possible configuration for 18 people to be seated around a table, as referred to in the text for $\vec{n} = (3, 4)$ and $\vec{v} = (2, 3)$.

is the same as the number of subdivisions of $\vec{v} \cdot \vec{n}$ points on a circumference in n_i sets of v_i point groups without crossing.

$C_{\vec{v}}(\vec{n})$ is also the number of polygonal dissections of an $(\vec{v} - \vec{1}) \cdot \vec{n} + 2$ -gon into $n_1 + \dots + n_s$ polygons with n_i of them having $v_i + 1$ edges, for $i = 1, \dots, s$. This can be found, for example, in [15].

Analogously with what happens with the Catalan numbers [16] and Fuss-Catalan numbers [6], these generalized Catalan numbers satisfy a recurrence relation that is an easy consequence of a result of Rhoades in [14] stating, in particular, that, if $\vec{r} \in \mathbb{N}_0^s$, $\vec{v} \in \mathbb{N}^s$, $m \in \mathbb{N}$ then

$$\sum_{\vec{r}_1 + \dots + \vec{r}_m = \vec{r}} C_{\vec{v}}(\vec{r}_1) \dots C_{\vec{v}}(\vec{r}_m) = \frac{m}{m + \vec{v} \cdot \vec{r}} \binom{m + \vec{v} \cdot \vec{r}}{\vec{r}}. \tag{5}$$

LEMMA 1. For $s \in \mathbb{N}$, $\vec{n} \in \mathbb{N}_0^s$ and $\vec{v} \in \mathbb{N}^s$ we have

$$\forall \vec{n} \in \mathbb{N}_0^s \setminus \{\vec{0}\} \quad C_{\vec{v}}(\vec{n}) = \sum_{i=1}^s \left(\sum_{\vec{r}_1 + \dots + \vec{r}_{v_i} = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \dots C_{\vec{v}}(\vec{r}_{v_i}) \right)^1 \tag{6}$$

where \vec{e}_i is the unit-vector with 1 in its i^{th} coordinate.

Proof. For $i = 1, \dots, s$ such that $n_i > 0$, using (5) for $m = v_i$ and $\vec{r} = \vec{n} - \vec{e}_i$, we obtain

$$\begin{aligned} \sum_{\vec{r}_1 + \dots + \vec{r}_{v_i} = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \dots C_{\vec{v}}(\vec{r}_{v_i}) &= \frac{v_i}{v_i + \vec{v} \cdot (\vec{n} - \vec{e}_i)} \binom{v_i + \vec{v} \cdot (\vec{n} - \vec{e}_i)}{\vec{n} - \vec{e}_i} \\ &= \frac{v_i}{\vec{v} \cdot \vec{n}} \binom{\vec{v} \cdot \vec{n}}{\vec{n} - \vec{e}_i} \\ &= \frac{(\vec{v} \cdot \vec{n})!}{(\vec{v} \cdot \vec{n})n_1! \dots n_s! \left((\vec{v} - \vec{1}) \cdot \vec{n} + 1 \right)!} v_i^{n_i} \end{aligned}$$

¹As $\vec{n} \neq \vec{0}$ the sum is never empty, although the second summation is, if $n_i = 0$.

and then

$$\sum_{i=1}^s \sum_{\vec{r}_1 + \dots + \vec{r}_i = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \cdots C_{\vec{v}}(\vec{r}_i) = \frac{(\vec{v} \cdot \vec{n})!}{n_1! \cdots n_s! \left((\vec{v} - \vec{1}) \cdot \vec{n} + 1 \right)!},$$

completing the proof. \square

Recall that a sequence $(a_n)_{n \in \mathbb{N}}$ is holonomic of order s ($s \in \mathbb{N}$) and degree t ($t \in \mathbb{N}_0$) if there exist p_0, p_1, \dots, p_s polynomials in n such that p_0 never vanishes (to simplify), the maximum of their degrees is t and

$$\forall n \in \mathbb{N} \quad \left[n > s \Rightarrow p_0(n)a_n = \sum_{i=1}^s p_i(n)a_{n-s} \right].$$

It is well known (the proof can be made, for example, using the Stirling approximation) that

$$C_d(n) \sim \frac{1}{\sqrt{2\pi}} \frac{\sqrt{d}}{(d-1)^{\frac{3}{2}}} \left(\frac{d^d}{(d-1)^{d-1}} \right)^n n^{-\frac{3}{2}}. \tag{7}$$

In the article [1] one can find good approximations of binomials of the form $\binom{dn}{n}$.

3. The zero of polynomials of particular kind

As it was said in the Introduction, any polynomial of the form (2) has a unique zero in \mathbb{Z}_M . This is a consequence of the following result, which is just a version of Hensel’s Lemma applied to this kind of polynomials, and of the Chinese Remainder Theorem.

LEMMA 2. *Let p be a prime number and $Q = Q(x)$ a polynomial of the form $a_d x^d + \dots + a_1 x + a_0$, where p divides a_i for $i \geq 2$ and p do not divide a_1 . Then, for all $k \in \mathbb{N}$, the congruence $Q(x) \equiv 0 \pmod{p^k}$ has a unique solution.*

Proof. If $k = 1$ then the result is trivial as $Q(x) \equiv 0 \pmod{p}$ is equivalent to $a_1 x + a_0 \equiv 0 \pmod{p}$ and a_1 is invertible modulo p . For $m \geq 1$, if x_m is the unique solution of $Q(x) \equiv 0 \pmod{p^m}$, then all solutions of $Q(x) \equiv 0 \pmod{p^{m+1}}$ are of the form $x = x_m + sp^m$, with $s \in \mathbb{Z}$. As p divides a_i for $i \geq 2$ and p^m divides $Q(x_m)$,

$$\begin{aligned} Q(x) \equiv 0 \pmod{p^{m+1}} &\Leftrightarrow Q(x_m) + a_1 p^m s \equiv 0 \pmod{p^{m+1}} \\ &\Leftrightarrow \frac{Q(x_m)}{p^m} + a_1 s \equiv 0 \pmod{p} \end{aligned}$$

and the conclusion follows as this last congruence has only one solution modulo p . \square

We now present an expression for the zero of polynomials of the form (2), for $M \in \mathbb{N}$. All the operations in this section are made in the ring \mathbb{Z}_M and it is clear that all the “infinite” sums referred to here only have a finite number of non-zero terms.

Let $d \geq 2$ and $\vec{v} = (v_2, \dots, v_d) \in \mathbb{N}^{d-1}$. Consider, for $\vec{x} = (x_2, \dots, x_d)$ whose coordinates are all nilpotent in \mathbb{Z}_M , the (finite) sum in \mathbb{Z}_M

$$y_{\vec{v}}(\vec{x}) = \sum_{\vec{n} \in \mathbb{N}_0^{d-1}} C_{\vec{v}}(\vec{n}) x_2^{n_2} \cdots x_d^{n_d}, \quad \text{where } \vec{n} = (n_2, \dots, n_d). \tag{8}$$

Notice that $y_{\vec{v}}(\vec{x})$ is always invertible as it is a sum of 1 with a nilpotent element.

LEMMA 3. *With the above notation,*

$$y_{\vec{v}}(\vec{x}) = 1 + x_2 y_{\vec{v}}(\vec{x})^{v_2} + \cdots + x_d y_{\vec{v}}(\vec{x})^{v_d}. \tag{9}$$

Proof. It is easy to see by comparing the terms of the sums that, for $i = 2, \dots, m$,

$$\begin{aligned} x_i \cdot y_{\vec{v}}(\vec{x})^{v_i} &= \sum_{\vec{n} \in \mathbb{N}_0^{d-1}} \left(\sum_{\vec{r}_1 + \cdots + \vec{r}_{v_i} = \vec{n}} C_{\vec{v}}(\vec{r}_1) \cdots C_{\vec{v}}(\vec{r}_{v_i}) \right) x_2^{n_2} \cdots x_d^{n_d} \cdot x_i \\ &= \sum_{\vec{n} \in \mathbb{N}_0^{d-1}, n_i \geq 1} \left(\sum_{\vec{r}_1 + \cdots + \vec{r}_{v_i} = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \cdots C_{\vec{v}}(\vec{r}_{v_i}) \right) x_2^{n_2} \cdots x_d^{n_d} \end{aligned}$$

and then, denoting by z the right side of (9),

$$\begin{aligned} z &= 1 + \sum_{i=2}^d \left(\sum_{\vec{n} \in \mathbb{N}_0^{d-1}, n_i \geq 1} \left(\sum_{\vec{r}_1 + \cdots + \vec{r}_{v_i} = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \cdots C_{\vec{v}}(\vec{r}_{v_i}) \right) x_2^{n_2} \cdots x_d^{n_d} \right) \\ &= 1 + \sum_{\vec{n} \in \mathbb{N}_0^{d-1} \setminus \{\vec{0}\}} \left(\sum_{i=2}^d \sum_{\vec{r}_1 + \cdots + \vec{r}_{v_i} = \vec{n} - \vec{e}_i} C_{\vec{v}}(\vec{r}_1) \cdots C_{\vec{v}}(\vec{r}_{v_i}) \right) x_2^{n_2} \cdots x_d^{n_d} \end{aligned}$$

and the conclusion follows using (6) and the fact that $C_{\vec{v}}(\vec{0}) = 1$. \square

We are now in the conditions to show an (algebraic) expression for the zero of a polynomial as in (2), whose existence and uniqueness are guaranteed by Lemma 2 and the Chinese Remainder Theorem.

THEOREM 1. *Let $M \in \mathbb{N}$ and $P(x) = a_d x^d + \cdots + a_1 x + a_0$ be a polynomial in \mathbb{Z}_M as in (2). Then the unique zero x of the polynomial is equal to the (finite) sum*

$$x_0 = -a_1^{-1} a_0 \sum_{\vec{n} = (n_2, \dots, n_d) \in \mathbb{N}_0^{d-1}} (-1)^{\vec{v} \cdot \vec{n}} C_{\vec{v}}(\vec{n}) a_0^{(\vec{v} - \vec{1}) \cdot \vec{n}} a_1^{-\vec{v} \cdot \vec{n}} a_2^{n_2} \cdots a_d^{n_d}, \tag{10}$$

where $\vec{v} = (2, 3, \dots, d)$ and $\vec{1} = (1, \dots, 1)$.

Moreover x_0 is invertible if and only if a_0 is invertible.

Proof. We find a solution x_0 of the form x_1y , where $y = y_{\bar{v}}(\bar{x})$ is defined in (8) for x_2, \dots, x_d nilpotents. Using equality (9),

$$\begin{aligned} P(x_1y) = 0 &\iff \sum_{i=2}^d a_i x_1^i y^i + a_1 x_1 y + a_0 = 0 \\ &\iff \sum_{i=2}^d a_i x_1^i y^i + a_1 x_1 (1 + x_2 y^2 + \dots + x_d y^d) + a_0 = 0 \\ &\iff \sum_{i=2}^d (a_i x_1^i + a_1 x_1 x_i) y^i + a_1 x_1 + a_0 = 0. \end{aligned}$$

So, if we choose

$$\begin{cases} x_1 = -a_0 a_1^{-1} \\ x_i = -a_i a_1^{-1} x_1^{i-1} = (-1)^i a_0^{i-1} a_1^{-i} a_i, \quad i \geq 2, \end{cases}$$

we obtain the referred solution.

The last observation is an immediate consequence of the fact that y is invertible, as mentioned before. \square

For example, the zero of the polynomial $a_d x^d + a_1 x + a_0$ is, with the previous notation, equal to the sum

$$x_0 = - \sum_{k \in \mathbb{N}_0} (-1)^{dk} C_d(k) a_0^{(d-1)k+1} a_1^{-dk-1} a_d^k.$$

In particular, if p is a prime number and r a multiple of p then, for $n \in \mathbb{N}_0$,

$$\sum_{k=0}^n C_d(k) r^k$$

is a solution of the congruence $rx^d - x + 1 \pmod{p^{n+1}}$.

The rate of growth, in n , of this sum, for all $r \neq 0$, follows from Theorem 3.

REMARK 1. Suppose we have a polynomial $Q(x) = \sum_{i=0}^d a_i x^i$ in \mathbb{Z}_M such that a_i are nilpotent for $i \leq d-2$, and a_{d-1} and a_d are invertible, which can be seen as a kind of reverse form of (2).

Q may have more than one solution, as we can see, for example, if $Q(x) = x^3 + x^2 + 3x + 9$ and $M = 27$, but only one is invertible. To prove this, consider the polynomial $Q^*(y) = \sum_{i=0}^d a_i y^{d-i}$, of the form (2), noticing that $y^d Q(y^{-1}) = Q^*(y)$, for invertible y .

4. Holonomic sequences related to Fuss-Catalan numbers

For $d \in \mathbb{N} \setminus \{1\}$, $r \in \mathbb{R} \setminus \{0\}$ and $n \in \mathbb{N}$, consider $X(d, r, n)$ defined in (3). We intend to obtain a recurrence relation for the sequence $(X(d, r, n))_{n \in \mathbb{N}}$, generalizing some cases referred to in OEIS, as mentioned in the Introduction.

For $n, k \in \mathbb{N}$, we let $(n)_k$ denote the *falling factorial* $\prod_{i=0}^{k-1} (n-i)$ ($= \frac{n!}{(n-k)!}$). Notice that $(n)_k$ is a polynomial in n of degree k .

THEOREM 2. *Let $d \in \mathbb{N} \setminus \{1\}$, and $r \in \mathbb{R} \setminus \{0\}$. Then $(X(d, r, n))_{n \in \mathbb{N}}$ is a holonomic sequence of order 2 and degree $d - 1$. More precisely, for $p_0(n) = ((d - 1)n + 1)_{d-1}$, $p_2(n) = d(dn - 1)_{d-1}$ and $p_1 = p_0 + rp_2$, we have*

$$\forall n \in \mathbb{N} \setminus \{1\} \quad p_0(n)X(d, r, n) = p_1(n)X(d, r, n - 1) - rp_2(n)X(d, r, n - 2).$$

Proof. As

$$\begin{aligned} & p_1(n)X(d, r, n - 1) - rp_2(n)X(d, r, n - 2) \\ &= p_0(n) \sum_{k=0}^{n-1} C_d(k)r^k + p_2(n) \sum_{k=0}^{n-1} C_d(k)r^{k+1} - p_2(n) \sum_{k=0}^{n-2} C_d(k)r^{k+1} \\ &= p_0(n) \sum_{k=0}^{n-1} C_d(k)r^k + p_2(n)C_d(n-1)r^n \\ &= p_0(n)X(d, r, n) - p_0(n)C_d(n)r^n + p_2(n)C_d(n-1)r^n, \end{aligned}$$

we only need to prove that $p_0(n)C_d(n) = p_2(n)C_d(n - 1)$. In fact,

$$\begin{aligned} \frac{C_d(n)}{C_d(n-1)} &= \frac{((d-1)(n-1)+1) \binom{dn}{n}}{((d-1)n+1) \binom{d(n-1)}{n-1}} \\ &= \frac{((d-1)(n-1)+1)}{((d-1)n+1)} \frac{(n-1)!((d-1)(n-1))! (dn)!}{n!((d-1)n)! (d(n-1))!} \\ &= \frac{((d-1)(n-1)+1)! (dn)!}{n((d-1)n+1)! (d(n-1))!} \\ &= \frac{(dn)_d}{n((d-1)n+1)_{d-1}} \\ &= \frac{d(dn-1)_{d-1}}{((d-1)n+1)_{d-1}}, \end{aligned}$$

which concludes the proof. \square

The following observation will be useful in the next section.

REMARK 2. Notice that a constant sequence satisfies the recurrence referred to in the previous theorem. As a consequence, if $(Z_n)_n$ is a non-constant solution of the recurrence, then $\langle (Z_n)_n, (1)_n \rangle$ is a basis of the space of solutions of the recurrence.

Notice also that the characteristic polynomial of the recurrence, $p_0(n)x^2 - p_1(n)x - rp_2(n)$, has the zeros 1 and $\frac{rp_2(n)}{p_0(n)}$ and that

$$\lim_{n \rightarrow \infty} \frac{rp_2(n)}{p_0(n)} = \frac{rd^d}{(d-1)^{d-1}}.$$

5. Asymptotics for generating functions of the Fuss-Catalan numbers

We are now in conditions to establish the asymptotic behaviour of the sequence $(X(d, r, n))_n$, when $\frac{|r|d^d}{(d-1)^{d-1}} > 1$ which, using (7), is when it diverges.

We use the following asymptotic behaviour: if $a, b \in \mathbb{Z}$, with $a \neq 0$, then

$$\prod_{j=2}^{n+1} (aj + b) = a^n \prod_{j=2}^{n+1} (j + \frac{b}{a}) = a^n \frac{\Gamma(n + 2 + \frac{b}{a})}{\Gamma(2 + \frac{b}{a})} \sim \frac{\Gamma(n)}{\Gamma(2 + \frac{b}{a})} a^n n^{2 + \frac{b}{a}} \tag{11}$$

as $\Gamma(x + \alpha) \sim \Gamma(x)x^\alpha$ when $x \rightarrow +\infty$.

REMARK 3. In order to apply Corollary 1.6 of [8] in the next theorem we draw the attention to the fact that, if p and q are two polynomials of the same degree s and q is never zero in \mathbb{N} , then

$$\sum_{n=1}^{\infty} \left| \frac{p(n+1)}{q(n+1)} - \frac{p(n)}{q(n)} \right| < \infty,$$

as the degree of the polynomial, in n , $p(n+1)q(n) - p(n)q(n+1)$ is at most $2s - 2$.

THEOREM 3. With the above notation, if $A(d) = \frac{d^d}{(d-1)^{d-1}}$ and $A(d)|r| > 1$,

$$X(d, r, n) \sim \frac{1}{\sqrt{2\pi}} \frac{\sqrt{d}}{(d-1)^{\frac{3}{2}}} \frac{A(d)r}{A(d)r-1} (A(d)r)^n n^{-\frac{3}{2}}.$$

Proof. By Remark 2, the zeros of the characteristic polynomial of the recurrence equation converge, when n tends to infinity, to different numbers, namely $A(d)r$ and 1. Therefore, and using Remark 3 for $p = p_i, i = 1, 2$ and $q = p_0$, we are in the conditions to apply Corollary 1.6 of [8]. In particular, there exists a solution $(Y_n)_n$ of the recurrence equation such that $Y_n \sim \prod_{j=2}^{n+1} \frac{rp_2(j)}{p_0(j)}$. Notice that, using (11), we have

$$\begin{aligned} \prod_{j=2}^{n+1} \frac{rp_2(j)}{p_0(j)} &= \prod_{j=2}^{n+1} \frac{rd(dj-1)_{d-1}}{((d-1)j+1)_{d-1}} = (rd)^n \prod_{i=1}^{d-1} \prod_{j=2}^{n+1} \frac{dj-i}{(d-1)j+2-i} \\ &\sim r^n d^n \prod_{i=1}^{d-1} \frac{\Gamma(2 + \frac{2-i}{d-1})}{\Gamma(2 - \frac{i}{d})} \left(\frac{d}{d-1}\right)^n n^{-\frac{i}{d} - \frac{2-i}{d-1}} \\ &= k_d r^n d^n \left(\frac{d}{d-1}\right)^{(d-1)n} n^{-\frac{3}{2}}, \quad \text{where } k_d = \left(\prod_{i=1}^{d-1} \frac{\Gamma(2 + \frac{2-i}{d-1})}{\Gamma(2 - \frac{i}{d})}\right) \\ &= k_d \left(\frac{d^d}{(d-1)^{d-1}} r\right)^n n^{-\frac{3}{2}}. \end{aligned}$$

As $\langle (Y_n)_n, (1)_n \rangle$ is a basis of the space of solutions of the recurrence, there exist $a, b \in \mathbb{R}$ such that, letting X_n denote $X(d, r, n)$, $(X_n)_n = a(Y_n)_n + b(1)_n$ and then

$$X_n \sim aY_n \sim ak_d(A(d)r)^n n^{-\frac{3}{2}}. \tag{12}$$

To calculate ak_d , using (7), we have

$$\frac{X_n - X_{n-1}}{Y_n} = \frac{C_d(n)r^n}{Y_n} \xrightarrow{n} \frac{1}{k_d\sqrt{2\pi}} \frac{\sqrt{d}}{(d-1)^{\frac{3}{2}}}$$

and, on the other hand, using (12),

$$\frac{X_n - X_{n-1}}{Y_n} = \frac{Y_n - Y_{n-1}}{Y_n} \xrightarrow{n} a \left(1 - \frac{1}{A(d)r} \right),$$

from where we obtain

$$ak_d = \frac{1}{\sqrt{2\pi}} \frac{\sqrt{d}}{(d-1)^{\frac{3}{2}}} \frac{A(d)r}{A(d)r-1},$$

concluding the proof. \square

REMARK 4. Although it is not relevant, we would like to point out that k_d referred to in the above proof is equal to $\frac{1}{\sqrt{2\pi}} \left(\frac{d}{d-1} \right)^{d+\frac{1}{2}}$.

Acknowledgements. The research of the authors were partially financed by Portuguese Funds through Fundação para a Ciência e a Tecnologia within the Projects UIDB/00013/2020 and UIDP/00013/2020.

REFERENCES

- [1] T. BURIĆ AND N. ELEZOVIĆ, *Asymptotic expansions of the binomial coefficients*, J. Appl. Math. Comput. **46** (2014) 135–145.
- [2] W. C. CHU, *A new combinatorial interpretation for generalized Catalan number*, Discrete Math. **65** (1987), 91–94.
- [3] N. ELEZOVIĆ, *Asymptotic expansions of central binomial coefficients and Catalan numbers*, J. Integer Seq. **17** (2014) Article 14.2.1.
- [4] N. ELEZOVIĆ, *Asymptotic expansions of gamma and related functions, binomial coefficients, inequalities and means*, J. Math. Inequalities **9** (2015) 1001–1054.
- [5] L. EULER, *Leonhardi Euleri—Opera omnia. Series 4 A. Commmercium epistolicum. Vol. 4.1. Leonhardi Euleri commercium epistolicum cum Christiano Goldbach. Pars I/Correspondence of Leonhard Euler with Christian Goldbach. Part I*, Original texts in Latin and German, Edited by Franz Lemmermeyer and Martin Mattmüller, Springer, Basel (2015).
- [6] N. I. FUSS, *Solutio quaestionis, quot modis polygonum n laterum in polygona m laterum, per diagonales resolvi queat*, Nova Acta Academiae Scientiarum Imperialis Petropolitanae **9** (2014) 243–251.
- [7] P. HILTON AND J. PEDERSEN, *Catalan numbers, their generalization, and their uses*, Math. Intelligencer **13** (2) (1991) 64–75.
- [8] R. J. KOOMAN, *Asymptotic behaviour of solutions of linear recurrences and sequences of Möbius-transformations*, J. Approx. Theory **93** (1998) 1–58.
- [9] P. J. LARCOMBE, *The 18th century Chinese discovery of the Catalan numbers*, Math. Spectrum **32** (1999) 5–7.
- [10] J. J. LUO, *Antu Ming, the first inventor of Catalan numbers in the world*, Neimenggu Daxue Xuebao **19** (1988) 239–245 (in Chinese).
- [11] S. MATTAREI, *Asymptotics of partial sums of central binomial coefficients and Catalan numbers*, arXiv preprint arXiv:0906.4290 (2009), available at <https://arxiv.org/abs/0906.4290>.

- [12] O. PERRON, *Über einen Satz des Herrn Poincaré*, J. Reine Angew Math. **136** (1909) 17–37.
- [13] H. POINCARÉ, *Sur les équations linéaires aux différentielles ordinaires et aux différences finies*, Amer. J. Math. **7** (1885) 203–258.
- [14] B. RHOADES, *Enumeration of connected Catalan objects by type*, European J. Combin. **32** (2011) 330–338.
- [15] A. SCHUETZ AND G. WHIELDON, *Polygonal dissections and reversions of series*, Involve **9** (2016) 223–236.
- [16] A. SEGNER, *Enumeratio modorum, quibus figurae planae rectilineae per diagonales dividuntur in triangula*, Nova Acta Academiae Scientiarum Imperialis Petropolitanae, **7** (1761) 203–209.
- [17] N. J. A. SLOANE et al., *The On-Line Encyclopedia of Integer Sequences*, OEIS Foundation Inc., <https://oeis.org>.
- [18] R. P. STANLEY, *Catalan Numbers*, Cambridge University Press, New York (2015).

(Received January 13, 2023)

Assis Azevedo
Center of Mathematics
University of Minho
Campus de Gualtar, 4710-157 Braga, Portugal
e-mail: assis@math.uminho.pt

Davide Azevedo
Center of Mathematics
University of Minho
Campus de Gualtar, 4710-157 Braga, Portugal
e-mail: davidemsa@math.uminho.pt