

GENERALIZED INTEGRATION OPERATORS FROM WEIGHTED BERGMAN SPACES INTO GENERAL FUNCTION SPACES

XIANGLING ZHU

(Communicated by I. Perić)

Abstract. This article studies the boundedness of the inclusion mapping from weighted Bergman spaces A_α^p into a class of tent type space $\mathcal{T}_s^{p,n}(\mu)$. As an application, the boundedness, compactness and essential norm of generalized integral operators $T_g^{n,k}$ and $S_g^{n,0}$ from A_α^p to general function spaces are also investigated.

1. Introduction

We denote by \mathbb{D} and $\partial\mathbb{D}$ the unit disk and its boundary in the complex plane \mathbb{C} , respectively. Let $H(\mathbb{D})$ be the class of functions analytic in \mathbb{D} . For $0 < p < \infty$ and $\alpha > -1$, the weighted Bergman space A_α^p is the set of all $f \in H(\mathbb{D})$ for which

$$\|f\|_{A_\alpha^p}^p = (\alpha + 1) \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dA(z) < \infty.$$

When $\alpha = 0$, we denote A_α^p by A^p . The Bloch space \mathcal{B} is the space of all $f \in H(\mathbb{D})$ such that

$$\|f\|_{\mathcal{B}} = |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

The little Bloch space \mathcal{B}_0 , is the set consisting of all $f \in H(\mathbb{D})$ such that $\lim_{|z| \rightarrow 1^-} (1 - |z|^2) |f'(z)| = 0$. Let H^∞ denote the space of all bounded analytic functions with the supremum norm $\|f\|_{H^\infty} = \sup_{z \in \mathbb{D}} |f(z)|$.

Let $0 < p, s < \infty$, $-2 < q < \infty$. The general function space $F(p, q, s)$, which was introduced by Zhao in [42], consists of all $f \in H(\mathbb{D})$ such that

$$\|f\|_{F(p,q,s)}^p = |f(0)|^p + \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^q (1 - |\sigma_a(z)|^2)^s dA(z) < \infty.$$

Here $\sigma_a(z) = \frac{a-z}{1-\bar{a}z}$. $F(p, q, s)$ is a Banach space under the norm $\|\cdot\|_{F(p,q,s)}$ when $p \geq 1$. It is easy to see that $F(p, p, 0)$ is just the Bergman space. When $p = 2$ and $q = 0$, it gives the Q_s space. Especially, Q_1 is the *BMOA* space, the space of analytic

Mathematics subject classification (2020): 30H20, 47B38.

Keywords and phrases: Weighted Bergman space, generalized integral operator, Carleson measure.

This work was supported by GuangDong Basic and Applied Basic Research Foundation (no. 2022A1515010317, no. 2023A1515010614).

functions in the Hardy space whose boundary functions have bounded mean oscillation. Also, it is known that $F(p, q, s)$ contains only constant functions if $s + q \leq -1$. For some other results on the space their generalizations and operators on them see also [15, 26, 28, 38].

Let $g \in H(\mathbb{D})$. The Volterra integral operator T_g and its companion operator I_g with symbol g are defined by

$$T_g f(z) = \int_0^z g'(w)f(w)dw, \quad I_g f(z) = \int_0^z g(w)f'(w)dw, \quad f \in H(\mathbb{D}),$$

respectively. The multiplication operator M_g is defined by $M_g f(z) = f(z)g(z)$. It is easy to see that

$$M_g f(z) = f(0)g(0) + I_g f(z) + T_g f(z).$$

Pommerenke [23] introduced the operator T_g and showed that T_g is bounded on H^2 if and only if g belongs to the space $BMOA$. For some generalizations on H^p spaces see [1, 2, 6, 8, 14, 29]. For some results on the Bergman-type spaces see [3, 8, 16]. Further results about Volterra integral operators on analytic function spaces on the unit disk, as well as the unit ball and unit polydisk in \mathbb{C}^n can be found [11, 12, 13, 15, 17, 18, 19, 21, 22, 24, 26, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39] as well as in their respective references.

We define the Carleson box, denoted by $S(I)$, based on the arc I which is a subset of $\partial\mathbb{D}$, as follows:

$$S(I) = \left\{ z \in \mathbb{D} : 1 - |I| \leq |z| < 1 \text{ and } \frac{z}{|z|} \in I \right\}.$$

If I equals the unit circle $\partial\mathbb{D}$, we let $S(I)$ be the entire unit disk \mathbb{D} . For a positive Borel measure μ on \mathbb{D} and $0 < s < \infty$, we say that μ is an s -Carleson measure if

$$\sup_{I \subset \partial\mathbb{D}} \frac{\mu(S(I))}{|I|^s} < \infty.$$

The classical Carleson measure is obtained when $s = 1$.

Let \mathbb{N} and \mathbb{N}_0 denote the set of positive integers and nonnegative integers, respectively. Let $0 < p, s < \infty$, $n \in \mathbb{N}_0$ and μ be a positive Borel measure on \mathbb{D} . We define $\mathcal{T}_s^{p,n}(\mu)$ as the set of functions $f \in H(\mathbb{D})$ satisfying the condition (see [25])

$$\sup_{I \subset \partial\mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} \left| f^{(n)}(z)(1 - |z|^2)^n \right|^p d\mu(z) < \infty.$$

Suppose $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$ satisfy $0 \leq k < n$, and $g \in H(\mathbb{D})$. Chalmoukis introduced the operator $T_g^{n,k}$ in [7], defined by

$$T_g^{n,k} f(z) = I^n (f^{(k)}(z)g^{(n-k)}(z)), \quad f \in H(\mathbb{D}),$$

where $I f(z) = \int_0^z f(w)dw$. In particular, $T_g^{1,0} f = T_g f$ for any $f \in H(\mathbb{D})$.

Qian and the author of this paper introduced and studied the operator $S_g^{n,0}$ in [25], where

$$S_g^{n,0} f(z) = I^n(f^{(n)}(z)g(z)).$$

In particular, $S_g^{1,0} f = I_g f$.

Chalmoukis investigated the boundedness of the operator $T_g^{n,k}$ on Hardy spaces H^p in [7]. Specifically, he demonstrated that $T_g^{n,k} : H^p \rightarrow H^p$ is bounded if and only if $g \in \mathcal{B}$ when $k \geq 1$. Meanwhile, $T_g^{n,k} : H^p \rightarrow H^q$ is bounded if and only if

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^{\frac{1}{q} - \frac{1}{p} + n - k} |g^{(n-k)}(z)| < \infty$$

when $0 < p < q < \infty$. In [9], Du, Li, and Qu studied the boundedness, compactness, and essential norm of the operator $T_g^{n,k}$ on weighted Bergman spaces induced by doubling weights. Further details related to the operator $T_g^{n,k}$, see [7, 9, 25].

The purpose of this paper is to establish that the inclusion mapping $I_d : A_\alpha^p \rightarrow \mathcal{T}_s^{p,n}(\mu)$ is bounded if and only if

$$\sup_{I \subset \partial \mathbb{D}} \frac{\int_{S(I)} (1 - |z|^2)^{pn} d\mu(z)}{|I|^{pn+2+\alpha+s}} < \infty. \tag{1.1}$$

This result is then applied to characterize the boundedness of $T_g^{n,k}$ and $S_g^{n,0}$, which act from A_α^p to $F(p, p + \alpha, s)$. Additionally, we investigate the essential norm and compactness of $T_g^{n,k}$ and $S_g^{n,0}$ when act from A_α^p to $F(p, p + \alpha, s)$.

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be Banach spaces, with $T : X \rightarrow Y$ being a bounded linear operator. The essential norm of $T : X \rightarrow Y$, denoted as $\|T\|_{e, X \rightarrow Y}$, can be defined as:

$$\|T\|_{e, X \rightarrow Y} = \inf_K \{\|T - K\|_{X \rightarrow Y} : K \text{ is compact from } X \text{ to } Y\}.$$

It is easy to observe that $T : X \rightarrow Y$ is compact if and only if $\|T\|_{e, X \rightarrow Y} = 0$.

Throughout this paper, we say that $f \lesssim g$ if there exists a constant C such that $f \leq Cg$. The symbol $f \approx g$ means that $f \lesssim g \lesssim f$.

2. Boundedness of $I_d : A_\alpha^p \rightarrow \mathcal{T}_s^{p,n}(\mu)$

In this section, our objective is to investigate the boundedness of the inclusion mapping $I_d : A_\alpha^p \rightarrow \mathcal{T}_s^{p,n}(\mu)$. To accomplish this task, we will introduce several lemmas that will be utilized throughout this paper.

LEMMA 1. [28, Theorem 3.2] *Let $-2 < q < \infty$, $0 < s < \infty$, $1 < p < \infty$ and $n \in \mathbb{N}$. Then the following statements are equivalent.*

- (i) $f \in F(p, q, s)$;
- (ii)

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{p(n-1)+q} (1 - |\sigma_a(z)|^2)^s dA(z) < \infty;$$

(iii)

$$\sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(n)}(z)|^p (1 - |z|^2)^{p(n-1)+q+s} dA(z) < \infty.$$

REMARK 1. Let

$$\|f\|_{F(p,q,s,1)}^p = \sum_{j=0}^{n-1} |f^{(j)}(0)|^p + \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{p(n-1)+q} (1 - |\sigma_a(z)|^2)^s dA(z),$$

$$\|f\|_{F(p,q,s,2)}^p = \sum_{j=0}^{n-1} |f^{(j)}(0)|^p + \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(n)}(z)|^p (1 - |z|^2)^{p(n-1)+q+s} dA(z).$$

From the proof of Theorem 3.2 of [28], we see that

$$\|f\|_{F(p,q,s)} \approx \|f\|_{F(p,q,s,1)} \approx \|f\|_{F(p,q,s,2)}.$$

LEMMA 2. [43, Theorem 4.28] *Let $-1 < \alpha < \infty$, $1 < p < \infty$ and $n \in \mathbb{N}$. Then $f \in A_\alpha^p$ if and only if*

$$\int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha} dA(z) < \infty.$$

Moreover,

$$\|f\|_{A_\alpha^p}^p \approx \sum_{j=0}^{n-1} |f^{(j)}(0)|^p + \int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha} dA(z).$$

LEMMA 3. [43, Theorem 5.4] *If f is analytic in \mathbb{D} and $n \geq 2$, then $f \in \mathcal{B}$ if and only if the function $(1 - |z|^2)^n f^{(n)}(z)$ is bounded in \mathbb{D} . Moreover, there exists a constant $C > 0$ such that*

$$C^{-1} \|f\|_{\mathcal{B}} \leq \sup_{z \in \mathbb{D}} (1 - |z|^2)^n |f^{(n)}(z)| \leq C \|f\|_{\mathcal{B}}$$

for all f with $f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0$.

The following result comes from [27, Theorem 4.1.2]. When $\beta = 2$, it was proved in [40]. When $n = 1$, it was proved in [41].

LEMMA 4. *Let $1 < p < \infty$, $1 < \beta < \infty$ and $n \in \mathbb{N}$. Then $g \in \mathcal{B}$ if and only if*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |g^{(n)}(z)|^p (1 - |z|^2)^{pn-2} (1 - |\sigma_a(z)|^2)^\beta dA(z) < \infty.$$

LEMMA 5. *Let $-1 < \alpha < \infty$, $0 < s < \infty$ and $1 < p < \infty$. Then*

$$\|f\|_{F(p,p+\alpha,s)}^p \lesssim \|f\|_{A_\alpha^p}^p.$$

Proof. By Lemma 2, we get

$$\begin{aligned} & \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{p+\alpha} (1 - |\sigma_a(z)|^2)^s dA(z) \\ & \leq \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{p+\alpha} dA(z) \\ & \lesssim \int_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\alpha dA(z), \end{aligned}$$

as desired. \square

The proof of the following result is standard. See, for example [25, Lemma 2.4]. We omit the proof here.

LEMMA 6. *Let $-1 < \alpha < \infty$, $0 < s < \infty$, $1 < p < \infty$ and $n \in \mathbb{N}_0$. If $f \in F(p, p + \alpha, s)$, then*

$$|f^{(n)}(z)| \lesssim \frac{\|f\|_{F(p, p+\alpha, s)}}{(1 - |z|^2)^{\frac{2+\alpha}{p} + n}}, \quad z \in \mathbb{D}.$$

LEMMA 7. [5, Lemma 2.1] *Let μ be a positive measure on \mathbb{D} and $0 < s < \infty$. Then μ is a bounded s -Carleson measure if and only if*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\frac{1 - |a|^2}{|1 - \bar{a}z|^2} \right)^s d\mu(z) < \infty.$$

Now we are in a position to state and prove our main result in this section.

THEOREM 1. *Let $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$ and $n \in \mathbb{N}_0$. Let μ be a positive Borel measure on \mathbb{D} . Then the inclusion mapping $I_d : A_\alpha^p \rightarrow \mathcal{F}_s^{p, n}(\mu)$ is bounded if and only if (1.1) holds.*

Proof. Assume first that (1.1) holds. Let $dv(z) = (1 - |z|^2)^{pn} d\mu(z)$. We observe that

$$\sup_{I \subset \partial \mathbb{D}} \frac{v(S(I))}{|I|^{pn+2+\alpha+s}} < \infty,$$

which, in combination with [43, Theorem 7.4], implies that the inclusion mapping $I_d : A_{pn+\alpha+s}^p \rightarrow L^p(dv)$ is a bounded operator. For any arc $I \subset \partial \mathbb{D}$, let ξ be the center point of I and $w = (1 - |I|)\xi$. From [10, p. 232], we see that

$$|1 - \bar{w}z| \approx 1 - |w|^2 \approx |I|, \quad z \in S(I). \tag{2.0}$$

Let $f \in A_\alpha^p$. By Lemma 2, we see that

$$f^{(n)} \in A_{pn+\alpha}^p \subset A_{pn+s+\alpha}^p.$$

Let $l > 2s$. Using the above results and Lemma 5, we obtain

$$\begin{aligned} & \frac{1}{|I|^s} \int_{S(I)} |f^{(n)}(z)|^p (1 - |z|^2)^{pn} d\mu(z) \\ & \approx (1 - |w|^2)^{l-s} \int_{S(I)} \left| \frac{f^{(n)}(z)}{(1 - \bar{w}z)^{l/p}} \right|^p dv(z) \quad (\text{using (2.0)}) \\ & \lesssim (1 - |w|^2)^{l-s} \int_{\mathbb{D}} \frac{|f^{(n)}(z)|^p}{|1 - \bar{w}z|^l} (1 - |z|^2)^{np+s+\alpha} dA(z) \\ & \quad (I_d : A_{pn+\alpha+s}^p \rightarrow L^p(dv) \text{ is bounded}) \\ & \leq \int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha} \frac{(1 - |z|^2)^s (1 - |w|^2)^s}{|1 - \bar{w}z|^{2s}} dA(z) \quad (l > 2s) \\ & \lesssim \|f\|_{F(p,p+\alpha,s)}^p \lesssim \|f\|_{A_\alpha^p}^p, \end{aligned}$$

which implies the desired result.

Conversely, assume that the inclusion mapping $I_d : A_\alpha^p \rightarrow \mathcal{F}_s^{p,n}(\mu)$ is bounded. Using this assumption and taking $f(z) = z^n \in A_\alpha^p$, we obtain

$$\int_{\mathbb{D}} (1 - |z|^2)^{pn} d\mu(z) < \infty.$$

For any arc $I \subset \partial\mathbb{D}$, let ξ be the center point of I and $w = (1 - |I|)\xi$. Take

$$f_w(z) = \frac{(1 - |w|^2)}{\bar{w}^n (1 - \bar{w}z)^{1 + \frac{2+\alpha}{p}}}. \tag{2.1}$$

By Lemma 3.10 of [43], we see that $f_w \in A_\alpha^p$. By (2.0),

$$|f_w^{(n)}(z)|^p \approx \frac{1}{|I|^{pn+2+\alpha}}, \quad z \in S(I).$$

By the assumption that the inclusion mapping $I_d : A_\alpha^p \rightarrow \mathcal{F}_s^{p,n}(\mu)$ is bounded, we have

$$\begin{aligned} \frac{1}{|I|^s} \int_{S(I)} |f_w^{(n)}(z)|^p (1 - |z|^2)^{pn} d\mu(z) & \lesssim \|I_d f_w\|_{\mathcal{F}_s^{p,n}(\mu)}^p \\ & \lesssim \|I_d\|^p \|f_w\|_{A_\alpha^p}^p \lesssim \|f_w\|_{A_\alpha^p}^p < \infty, \end{aligned}$$

which implies that

$$\sup_{I \subset \partial\mathbb{D}} \frac{\int_{S(I)} (1 - |z|^2)^{pn} d\mu(z)}{|I|^{pn+2+\alpha+s}} < \infty.$$

So, (1.1) holds. The proof is complete. \square

3. Boundedness

In this section, we provide some characterizations for the boundedness of $T_g^{n,k}$ and $S_g^{n,0}$ from A_α^p to $F(p, p + \alpha, s)$.

THEOREM 2. *Let $g \in H(\mathbb{D})$, $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$, $n \in \mathbb{N}$, $k \in \mathbb{N}_0$ such that $0 \leq k < n$. Then $T_g^{n,k} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded if and only if $g \in \mathcal{B}$. Moreover,*

$$\|T_g^{n,k}\|_{A_\alpha^p \rightarrow F(p,p+\alpha,s)} \approx \sup_{z \in \mathbb{D}} (1 - |z|^2) |g'(z)|.$$

Proof. Assume that $g \in \mathcal{B}$. We first consider the case $k = 0$. Since $g \in \mathcal{B}$ and $s + \alpha + 2 > 1$, by Lemma 4 we see that

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |g^{(n)}(z)|^p (1 - |z|^2)^{pn-2} (1 - |\sigma_a(z)|^2)^{s+\alpha+2} dA(z) < \infty,$$

which implies that

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left(\frac{1 - |a|^2}{1 - \bar{a}z} \right)^{s+2+\alpha} d\mu_g(z) < \infty,$$

where $d\mu_g(z) = |g^{(n)}(z)|^p (1 - |z|^2)^{pn+s+\alpha} dA(z)$. Using Lemma 7, we see that μ_g is an $(s + 2 + \alpha)$ -Carleson measure. Let $f \in A_\alpha^p$. From Theorem 1, we can easily deduce that

$$\begin{aligned} & \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |(T_g^{n,0} f)^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &= \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f(z)|^p |g^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &= \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f(z)|^p d\mu_g(z) \\ &\lesssim \|f\|_{A_\alpha^p}^p. \end{aligned} \tag{3.1}$$

Now we consider the case $k \geq 1$. From Lemmas 1, 3 and 5 we get

$$\begin{aligned} & \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |(T_g^{n,k} f)^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &= \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(k)}(z) g^{(n-k)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &\lesssim \|g\|_{\mathcal{B}}^p \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(k)}(z)|^p (1 - |z|^2)^{pk+\alpha+s} dA(z) \\ &\lesssim \|g\|_{\mathcal{B}}^p \|f\|_{F(p,p+\alpha,s)}^p \\ &\lesssim \|g\|_{\mathcal{B}}^p \|f\|_{A_\alpha^p}^p. \end{aligned} \tag{3.2}$$

From (3.1) and (3.2), we see that $T_g^{n,k} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded.

Conversely, assume that $T_g^{n,k} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded. For $w \in \mathbb{D}$, we define

$$G_w(z) = \frac{(1 - |w|^2)}{(1 - \bar{w}z)^{1 + \frac{2+\alpha}{p}}}.$$

It is easy to check that $G_w \in A_\alpha^p$ using Lemma 3.10 of [43]. Moreover,

$$G_w^{(k)}(z) = \prod_{i=1}^k \left(i + \frac{2 + \alpha}{p} \right) \frac{\bar{w}^k (1 - |w|^2)}{(1 - \bar{w}z)^{1+k + \frac{2+\alpha}{p}}},$$

$$G_w^{(k)}(w) = \prod_{i=1}^k \left(i + \frac{2 + \alpha}{p} \right) \frac{\bar{w}^k}{(1 - |w|^2)^{k + \frac{2+\alpha}{p}}}.$$

Using Lemma 6, we obtain that

$$\frac{\|T_g^{n,k} G_w\|_{F(p,p+\alpha,s)}}{(1 - |w|^2)^{n + \frac{2+\alpha}{p}}} \gtrsim |(T_g^{n,k} G_w)^{(n)}(w)| \gtrsim \frac{|w|^k |g^{(n-k)}(w)|}{(1 - |w|^2)^{k + \frac{2+\alpha}{p}}}.$$

Thus,

$$\sup_{|w| > 1/2} |g^{(n-k)}(w)| (1 - |w|^2)^{n-k} < \infty.$$

It is obvious that

$$\sup_{|w| \leq 1/2} |g^{(n-k)}(w)| (1 - |w|^2)^{n-k} < \infty.$$

Therefore,

$$\sup_{w \in \mathbb{D}} |g^{(n-k)}(w)| (1 - |w|^2)^{n-k} < \infty,$$

which implies that $g \in \mathcal{B}$ by Lemma 3. The proof is complete. \square

THEOREM 3. *Let $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$, $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$ such that $0 \leq k < n$. Then $S_g^{n,0} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded if and only if $g \in H^\infty$. Moreover,*

$$\|S_g^{n,0}\|_{A_\alpha^p \rightarrow F(p,p+\alpha,s)} \approx \|g\|_{H^\infty}. \tag{3.3}$$

Proof. We first assume that $S_g^{n,0} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded. For $b \in \mathbb{D}$ and $r > 0$, let $\mathbb{D}(b, r)$ denote the Bergman metric disk centered at b with radius r . From [43], we see that

$$\frac{(1 - |b|^2)^2}{|1 - \bar{b}z|^4} \approx \frac{1}{(1 - |z|^2)^2} \approx \frac{1}{(1 - |b|^2)^2}, \quad z \in \mathbb{D}(b, r). \tag{3.4}$$

For any $w \in \mathbb{D} \setminus \{0\}$, let f_w be defined in (2.1). We have that $f_w \in A_\alpha^p$ by Lemma 3.10 of [43]. Using (3.4), we obtain

$$|f_w^{(n)}(z)|^p \approx \frac{1}{(1 - |z|)^{np+2+\alpha}}, \quad z \in \mathbb{D}(w, r).$$

Therefore,

$$\begin{aligned} \infty &> \|S_g^{n,0} f\|_{F(p,p+\alpha,s)}^p \\ &= \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |(S_g^{n,0} f_w)^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha} (1 - |\sigma_a(z)|^2)^s dA(z) \\ &= \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f_w^{(n)}(z)g(z)|^p (1 - |z|^2)^{pn+\alpha} (1 - |\sigma_a(z)|^2)^s dA(z) \\ &\gtrsim \int_{\mathbb{D}(w,r)} |f_w^{(n)}(z)g(z)|^p (1 - |z|^2)^{pn+\alpha} (1 - |\sigma_w(z)|^2)^s dA(z) \\ &\gtrsim \int_{\mathbb{D}(w,r)} |g(z)|^p (1 - |z|^2)^{-2} dA(z) \\ &\gtrsim |g(w)|^p, \end{aligned}$$

which implies that $g \in H^\infty$.

Conversely, suppose that $g \in H^\infty$. Let $f \in A_\alpha^p$. Then by Lemma 1 we obtain

$$\begin{aligned} \|S_g^{n,0} f\|_{F(p,p+\alpha,s)}^p &\approx \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |(S_g^{n,0} f)^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &= \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(n)}(z)g(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &\lesssim \|g\|_{H^\infty}^p \sup_{I \subset \partial \mathbb{D}} \frac{1}{|I|^s} \int_{S(I)} |f^{(n)}(z)|^p (1 - |z|^2)^{pn+\alpha+s} dA(z) \\ &\lesssim \|g\|_{H^\infty}^p \|f\|_{F(p,p+\alpha,s)}^p \\ &\lesssim \|g\|_{H^\infty}^p \|f\|_{A_\alpha^p}^p. \end{aligned}$$

Therefore, $S_g^{n,0} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded. From the above proof, we see that (3.3) holds. The proof is complete. \square

4. Essential norm

In this section, we investigate the essential norm of $T_g^{n,k}$ and $S_g^{n,0}$ from A_α^p into $F(p, p + \alpha, s)$. The proof of the following result can be proved similarly as [25, Lemma 5.1]. We omit the proof here.

LEMMA 8. *Let $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$, $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$ such that $0 \leq k < n$. If $0 < r < 1$ and $g \in \mathcal{B}$, then $T_{g_r}^{n,k} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is compact.*

For $0 < r < 1$, $z \in \mathbb{D}$ and $f \in \mathcal{B}$, set $f_r(z) = f(rz)$. Let $\text{dist}_{\mathcal{B}}(f, \mathcal{B}_0)$ denote the distance from the Bloch function to the little Bloch space, that is,

$$\text{dist}_{\mathcal{B}}(f, \mathcal{B}_0) = \inf_{g \in \mathcal{B}_0} \|f - g\|_{\mathcal{B}}.$$

The following result can be found in [4].

LEMMA 9. *If $g \in \mathcal{B}$, then*

$$\limsup_{|z| \rightarrow 1^-} (1 - |z|^2) |g'(z)| \approx \text{dist}_{\mathcal{B}}(g, \mathcal{B}_0) \approx \limsup_{r \rightarrow 1^-} \|g - g_r\|_{\mathcal{B}}.$$

THEOREM 4. *Let $g \in H(\mathbb{D})$, $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$, $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$ such that $0 \leq k < n$. If $T_g^{n,k} : A_{\alpha}^p \rightarrow F(p, p + \alpha, s)$ is bounded, then*

$$\|T_g^{n,k}\|_{e, A_{\alpha}^p \rightarrow F(p, p + \alpha, s)} \approx \limsup_{|z| \rightarrow 1^-} (1 - |z|^2) |g'(z)| \approx \text{dist}_{\mathcal{B}}(g, \mathcal{B}_0).$$

Proof. Let $0 < r < 1$. By Lemma 8, $T_{g_r}^{n,k} : A_{\alpha}^p \rightarrow F(p, p + \alpha, s)$ is compact. Then by Theorem 2,

$$\|T_g^{n,k}\|_{e, A_{\alpha}^p \rightarrow F(p, p + \alpha, s)} \leq \|T_g^{n,k} - T_{g_r}^{n,k}\| = \|T_{g - g_r}^{n,k}\| \approx \|g - g_r\|_{\mathcal{B}}.$$

Using Lemma 9, we have

$$\|T_g^{n,k}\|_{e, A_{\alpha}^p \rightarrow F(p, p + \alpha, s)} \lesssim \limsup_{r \rightarrow 1^-} \|g - g_r\|_{\mathcal{B}} \approx \text{dist}_{\mathcal{B}}(g, \mathcal{B}_0).$$

On the other hand, suppose $\{z_j\}$ is a sequence in \mathbb{D} such that $\lim_{j \rightarrow \infty} |z_j| = 1$. Let G_{z_j} be defined as in the proof of Theorem 2 for each j . Then $\{G_{z_j}\}$ is a bounded sequence in A_{α}^p , and as $j \rightarrow \infty$, it converges uniformly to zero on every compact subset of \mathbb{D} . Let $K : A_{\alpha}^p \rightarrow F(p, p + \alpha, s)$ be a compact operator. Since A_{α}^p is a reflexive space we have that $\lim_{j \rightarrow \infty} \|KG_{z_j}\|_{F(p, p + \alpha, s)} = 0$ (see [20]). From the proof of Theorem 2, we also have

$$\begin{aligned} \|T_g^{n,k} - K\| &\gtrsim \limsup_{j \rightarrow \infty} \|(T_g^{n,k} - K)G_{z_j}\|_{F(p, p + \alpha, s)} \\ &\gtrsim \limsup_{j \rightarrow \infty} \left(\|T_g^{n,k}G_{z_j}\|_{F(p, p + \alpha, s)} - \|KG_{z_j}\|_{F(p, p + \alpha, s)} \right) \\ &\approx \limsup_{j \rightarrow \infty} \|T_g^{n,k}G_{z_j}\|_{F(p, p + \alpha, s)} \\ &\gtrsim \limsup_{j \rightarrow \infty} (1 - |z_j|^2)^{n-k} |g^{(n-k)}(z_j)|. \end{aligned}$$

Hence,

$$\|T_g^{n,k}\|_{e, A_{\alpha}^p \rightarrow F(p, p + \alpha, s)} \gtrsim \limsup_{j \rightarrow \infty} (1 - |z_j|^2)^{n-k} |g^{(n-k)}(z_j)|.$$

It follows from the arbitrariness of $\{z_j\}$ and Lemmas 3 and 9 that

$$\begin{aligned} \|T_g^{n,k}\|_{e,A_\alpha^p \rightarrow F(p,p+\alpha,s)} &\gtrsim \limsup_{|z| \rightarrow 1^-} (1 - |z|^2)^{n-k} |g^{(n-k)}(z)| \\ &\approx \limsup_{|z| \rightarrow 1^-} (1 - |z|^2) |g'(z)| \approx \text{dist}_{\mathcal{B}}(g, \mathcal{B}_0). \end{aligned}$$

The proof is complete. \square

The following result can be deduced by Theorem 4 directly.

COROLLARY 1. *Let $g \in H(\mathbb{D})$, $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$, $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$ such that $0 \leq k < n$. Then $T_g^{n,k} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is compact if and only if $g \in \mathcal{B}_0$.*

THEOREM 5. *Let $g \in H(\mathbb{D})$, $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$ and $n \in \mathbb{N}$. If $S_g^{n,0} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is bounded, then*

$$\|S_g^{n,0}\|_{e,A_\alpha^p \rightarrow F(p,p+\alpha,s)} \approx \limsup_{|z| \rightarrow 1^-} |g(z)|.$$

Proof. By Theorem 3 we have

$$\|S_g^{n,0}\|_{e,A_\alpha^p \rightarrow F(p,p+\alpha,s)} = \inf_K \|S_g^{n,0} - K\| \lesssim \limsup_{|z| \rightarrow 1^-} |g(z)|.$$

On the other hand, suppose $\{z_j\}$ is a sequence in \mathbb{D} such that $\lim_{j \rightarrow \infty} |z_j| = 1$. Let f_{z_j} be defined as in (2.1). Let $K : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ be a compact operator. Then, we similarly have $\lim_{j \rightarrow \infty} \|K f_{z_j}\|_{F(p,p+\alpha,s)} = 0$. Hence,

$$\begin{aligned} \|S_g^{n,0} - K\| &\gtrsim \limsup_{j \rightarrow \infty} \|(S_g^{n,0} - K) f_{z_j}\|_{F(p,p+\alpha,s)} \\ &\gtrsim \limsup_{j \rightarrow \infty} \|S_g^{n,0} f_{z_j}\|_{F(p,p+\alpha,s)} - \limsup_{j \rightarrow \infty} \|K f_{z_j}\|_{F(p,p+\alpha,s)} \\ &= \limsup_{j \rightarrow \infty} \|S_g^{n,0} f_{z_j}\|_{F(p,p+\alpha,s)}. \end{aligned}$$

Therefore, from the proof of Theorem 3,

$$\|S_g^{n,0}\|_{e,A_\alpha^p \rightarrow F(p,p+\alpha,s)} \gtrsim \limsup_{j \rightarrow \infty} \|S_g^{n,0} f_{z_j}\|_{F(p,p+\alpha,s)} \gtrsim \limsup_{j \rightarrow \infty} |g(z_j)|,$$

which implies that

$$\|S_g^{n,0}\|_{e,A_\alpha^p \rightarrow F(p,p+\alpha,s)} \gtrsim \limsup_{|z| \rightarrow 1^-} |g(z)|.$$

The proof is complete. \square

From Theorem 5 we get the following result.

COROLLARY 2. Let $1 < p < \infty$, $-1 < \alpha < \infty$, $0 < s < \infty$ and $n \in \mathbb{N}$. Then $S_g^{n,0} : A_\alpha^p \rightarrow F(p, p + \alpha, s)$ is compact if and only if $g = 0$.

Data Availability. No data were used to support this study.

Conflicts of Interest. The author declare that she has no conflicts of interest.

REFERENCES

- [1] A. ALEMAN AND J. CIMA, *An integral operator on H^p and Hardy's inequality*, J. Anal. Math. 85 (2001), 157–176.
- [2] A. ALEMAN AND A. SISKAKIS, *An integral operator on H^p* , Complex Variables Theory Appl. 28 (1995), 149–158.
- [3] A. ALEMAN AND A. SISKAKIS, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. 46 (1997), 337–356.
- [4] K. ATTELE, *Interpolating sequences for the derivatives of Bloch functions*, Glasgow Math. J. 34 (1992), 35–41.
- [5] R. AULASKARI, D. STEGENGA AND J. XIAO, *Some subclasses of BMOA and their characterization in terms of Carleson measures*, Rocky Mountain J. Math. 26 (1996), 485–506.
- [6] K. AVETISYAN AND S. STEVIĆ, *Extended Cesàro operators between different Hardy spaces*, Appl. Math. Comput. 207 (2009), 346–350.
- [7] N. CHALMOUKIS, *Generalized integration operators on Hardy spaces*, Proc. Amer. Math. Soc. 148 (2020), 3325–3337.
- [8] D. CHANG, S. LI AND S. STEVIĆ, *On some integral operators on the unit polydisk and the unit ball*, Taiwanese J. Math. 11 (5) (2007), 1251–1286.
- [9] J. DU, S. LI AND D. QU, *The generalized Volterra integral operator and Toeplitz operator on weighted Bergman spaces*, Mediterr. J. Math. (2022) 19:263, <https://doi.org/10.1007/s00009-022-02161-9>.
- [10] J. GARNETT, *Bounded Analytic Functions*, Academic Press, New York, 1981.
- [11] P. LI, J. LIU AND Z. LOU, *Integral operators on analytic Morrey spaces*, Sci. China Math. 57 (2014), 1961–1974.
- [12] S. LI, J. LIU AND C. YUAN, *Embedding theorems for Dirichlet type spaces*, Canad. Math. Bull. 63 (2020), 106–117.
- [13] S. LI AND S. STEVIĆ, *Integral type operators from mixed-norm spaces to α -Bloch spaces*, Integral Transforms Spec. Funct. 18 (7) (2007), 485–493.
- [14] S. LI AND S. STEVIĆ, *Riemann-Stieltjes operators on Hardy spaces in the unit ball of \mathbb{C}^n* , Bull. Belg. Math. Soc. Simon Stevin 14 (4) (2007), 621–628.
- [15] S. LI AND S. STEVIĆ, *Compactness of Riemann-Stieltjes operators between $F(p, q, s)$ and α -Bloch spaces*, Publ. Math. Debrecen 72 (1–2) (2008), 111–128.
- [16] S. LI AND S. STEVIĆ, *Riemann-Stieltjes operators between different weighted Bergman spaces*, Bull. Belg. Math. Soc. Simon Stevin 15 (4) (2008), 677–686.
- [17] S. LI AND S. STEVIĆ, *Cesàro type operators on some spaces of analytic functions on the unit ball*, Appl. Math. Comput. 208 (2009), 378–388.
- [18] S. LI AND S. STEVIĆ, *Integral-type operators from Bloch-type spaces to Zygmund-type spaces*, Appl. Math. Comput. 215 (2009), 464–473.
- [19] S. LI AND H. WULAN, *Volterra type operators on Q_K spaces*, Taiwanese J. Math. 14 (2010), 195–211.
- [20] M. LINDSTRÖM, D. NORRBO AND S. STEVIĆ, *On compactness of operators from Banach spaces of holomorphic functions to Banach space*, J. Math. Inequal. 17 (2023) to appear.
- [21] X. LIU, S. LI AND R. QIAN, *Volterra type operators and Carleson embedding on Campanato spaces*, J. Nonlinear Var. Anal. 5 (2021), 141–153.
- [22] J. PAU AND R. ZHAO, *Carleson measures, Riemann-Stieltjes and multiplication operators on a general family of function spaces*, Integral Equations and Operator Theory 78 (2014), 483–514.

- [23] C. POMMERENKE, *Schlichte funktionen und analytische funktionen von beschränkten mittlerer Oszillation*, *Comm. Math. Helv.* 52 (1977), 591–602.
- [24] R. QIAN AND S. LI, *Volterra type operators on Morrey type spaces*, *Math. Inequal. Appl.* 18 (2015), 1589–1599.
- [25] R. QIAN AND X. ZHU, *Embedding Hardy spaces H^p into tent spaces and generalized integration operators*, *Ann. Polon. Math.* 128 (2022), 143–157.
- [26] R. QIAN AND X. ZHU, *Volterra integral operator from weighted Bergman spaces to general function spaces*, *Math. Inequal. Appl.* 25 (2022), 985–998.
- [27] J. RÄTTYÄ, *On some complex function spaces and classes*, *Ann. Acad. Sci. Fenn. Math. Diss.* 124 (2001), 1–73.
- [28] J. RÄTTYÄ, *n -th derivative characterizations, mean growth of derivatives and $F(p, q, s)$* , *Bull. Australian Math. Soc.* 68 (2003), 405–421.
- [29] B. SEHBA AND S. STEVIĆ, *On some product-type operators from Hardy-Orlicz and Bergman-Orlicz spaces to weighted-type spaces*, *Appl. Math. Comput.* 233 (2014), 565–581.
- [30] C. SHEN, Z. LOU AND S. LI, *Embedding of $BMOA_{\log}$ into tent spaces and Volterra integral operators*, *Comput. Methods Funct. Theory* 20 (2020), 217–234.
- [31] Y. SHI AND S. LI, *Essential norm of integral operators on Morrey type spaces*, *Math. Inequal. Appl.* 19 (2016), 385–393.
- [32] S. STEVIĆ, *Boundedness and compactness of an integral operator on a weighted space on the polydisc*, *Indian J. Pure Appl. Math.* 37 (6) (2006), 343–355.
- [33] S. STEVIĆ, *Boundedness and compactness of an integral operator on mixed norm spaces on the polydisc*, *Siberian Math. J.* 48 (3) (2007), 559–569.
- [34] S. STEVIĆ, *Integral-type operators from a mixed norm space to a Bloch-type space on the unit ball*, *Siberian Math. J.* 50 (6) (2009), 1098–1105.
- [35] S. STEVIĆ, *Products of integral-type operators and composition operators from the mixed norm space to Bloch-type spaces*, *Siberian Math. J.* 50 (4) (2009), 726–736.
- [36] S. STEVIĆ, *On an integral-type operator from Zygmund-type spaces to mixed-norm spaces on the unit ball*, *Abstr. Appl. Anal.* 2010, Article ID 198608, (2010), 7 pages.
- [37] S. STEVIĆ, *On operator P_{ϕ}^g from the logarithmic Bloch-type space to the mixed-norm space on unit ball*, *Appl. Math. Comput.* 215 (2010), 4248–4255.
- [38] S. STEVIĆ, *On some integral-type operators between a general space and Bloch-type spaces*, *Appl. Math. Comput.* 218 (2011), 2600–2618.
- [39] S. STEVIĆ AND S. UEKI, *Integral-type operators acting between weighted-type spaces on the unit ball*, *Appl. Math. Comput.* 215 (2009), 2464–2471.
- [40] K. STROETHOFF, *Besov-type characterizations for the Bloch space*, *Bull. Austral. Math. Soc.* 39 (1989), 405–420.
- [41] R. ZHAO, *On α -Bloch functions and $VMOA$* , *Acta. Math. Sci.* 16 (1996), 349–360.
- [42] R. ZHAO, *On a general family of function spaces*, *Ann. Acad. Sci. Fenn. Math. Diss. No. 105* (1996), 1–56.
- [43] K. ZHU, *Operator Theory in Function Spaces*, second edition, *Math. Surveys and Monographs*, 138 (2007).

(Received June 16, 2023)

Xiangling Zhu
 University of Electronic Science and Technology of China
 Zhongshan Institute
 528402, Zhongshan, Guangdong, P. R. China
 e-mail: jyuzxl@163.com