

ON THE MINIMUM RANK OF DISTANCE MATRICES

ZAHRA GACHKOOBAN AND RAHIM ALIZADEH*

Abstract. Let $X = \{x_1, \dots, x_n\}$ be a finite set endowed with a metric d . The matrix $A = (d(x_i, x_j))_{n \times n}$ is called a distance matrix. In this paper we discuss about the minimum rank that can be achieved by an $n \times n$ distance matrix and prove that the rank of every 5×5 and 6×6 distance matrix is not less than 4.

Mathematics subject classification (2020): 15B99, 46B85.

Keywords and phrases: Distance matrix, matrix rank, finite metric space.

REFERENCES

- [1] K. BALL, *Isometric embedding in l_p -spaces*, European Journal of Combinatorics., **11**, 4 (1990), 305–311.
- [2] J. C. GOWER, *Properties of Euclidean and non-Euclidean distance matrices*, Linear Algebra and Its Applications., **67**, (1985), 81–97.
- [3] W. HOLSZTYNSKI, \mathbb{R}^n as universal metric space, Notices of the Amer. Math. Soc., **25**, 3 (1978), A-367.
- [4] C. K. LI, T. MILLIGAN, AND M. TROSSET, *Euclidean and circum-Euclidean distance matrices: Characterizations and linear preservers*, The Electronic Journal of Linear Algebra., **20**, (2010), 739–752.
- [5] H. MAEHARA, *Euclidean embeddings of finite metric spaces*, Discrete Mathematics., **313**, 23 (2013), 2848–2856.
- [6] H. S. WITSENHAUSEN, *Minimum dimension embedding of finite metric spaces*, Journal of Combinatorial Theory Series A., **42**, 2 (1986), 184–199.