

ONE-SIDED MAXIMAL OPERATORS ON HERZ SPACES WITH VARIABLE EXPONENTS

KWOK-PUN HO

(Communicated by S. Varošanec)

Abstract. This paper extends the boundedness of the one-sided maximal operators from the Lebesgue spaces with variable exponents to the one-sided Herz spaces with variable exponents. The main result generalizes the boundedness of the one-sided maximal operators on the classical Herz spaces.

1. Introduction

In this paper, we establish the boundedness of the one-sided maximal operators on the one-sided Herz spaces with variable exponents.

The Herz spaces with variable exponents are extensions of the Herz spaces and the Lebesgue spaces with variable exponents. The Herz spaces were introduced by Herz in [9] to study the Fourier series. The Herz spaces also provide applications on summability of Fourier series, harmonic analysis and partial differential equations, see [6, 15, 22, 30, 31]. The Lebesgue spaces with variable exponent are extensions of the Lebesgue spaces. It provides applications on partial differential equation and fluid dynamics, see [4, Part III]. An important breakthrough for the studies of the Lebesgue spaces with variable exponent is on the boundedness of the Hardy-Littlewood maximal function, see [3, 4, 27, 28].

The one-sided maximal operators are the “original” maximal function studies by Hardy and Littlewood in [8]. They are ancestors of the nowadays well known Hardy-Littlewood maximal function. The study of the one-sided maximal operators has its own independent interest. For example, it offers applications on the ergodic theory. For the mapping properties of the one-sided maximal operators and their applications, see [23, 24, 25, 26, 34].

The boundedness of the one-sided maximal operators has been extended to the Lebesgue spaces with variable exponents in [5, 29]. It motivates us to investigate the boundedness of the one-sided maximal operators on the Herz spaces with variable exponents. We introduce the one-sided Herz spaces with variable exponents to study the boundedness of the one-sided maximal operators. We obtain our main result by using the notion of localized operators introduced in [14].

Mathematics subject classification (2020): 26D10, 42B25, 42B35, 46E40.

Keywords and phrases: Herz spaces, one-sided maximal operators, variable exponent.

This paper is organized as follows. Section 2 contains the definitions of the one-sided maximal operators, the Lebesgue spaces with variable exponents and the one-sided Herz spaces with variable exponents. The boundedness of the one-sided maximal operators on the Lebesgue spaces with variable exponents is also recalled in this section. The main result of this paper, the boundedness of the one-sided maximal operators on the one-sided Herz spaces with variable exponents, is established in Section 3.

2. Definitions and preliminaries

Let \mathcal{M} and L^1_{loc} denote the space of Lebesgue measurable functions and the space of locally integrable functions on \mathbb{R} , respectively.

For any $f \in L^1_{\text{loc}}$, the one-sided Hardy-Littlewood maximal operators M^+f and M^-f are defined as

$$M^+f(x) = \sup_{t>0} \frac{1}{t} \int_x^{x+t} |f(y)| dy, \quad x \in \mathbb{R},$$

$$M^-f(x) = \sup_{t>0} \frac{1}{t} \int_{x-t}^x |f(y)| dy, \quad x \in \mathbb{R},$$

respectively.

We now recall the definition of the Lebesgue spaces with variable exponents on \mathbb{R} [3, 4].

DEFINITION 1. Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. The Lebesgue space with variable exponent $L^{p(\cdot)}$ consists of all Lebesgue measurable functions $f : \mathbb{R} \rightarrow \mathbb{C}$ so that

$$\|f\|_{L^{p(\cdot)}} = \inf \{ \lambda > 0 : \rho(f/\lambda) \leq 1 \} < \infty$$

where

$$\rho(f) = \int_{\mathbb{R}} |f(x)|^{p(x)} dx.$$

We call $p(\cdot)$ the exponent function of $L^{p(\cdot)}$.

The Lebesgue space with variable exponent is a Banach function space. For simplicity, we refer the reader to [2, Chapter 1] for the definition of Banach function space and its properties.

For any exponent function $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$, write

$$p_+ = \operatorname{ess\,sup}_{x \in (0, \infty)} p(x) \quad \text{and} \quad p_- = \operatorname{ess\,inf}_{x \in (0, \infty)} p(x).$$

For any Lebesgue measurable function $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$, define $p'(x) = \frac{p(x)}{p(x)-1}$, $x \in \mathbb{R}$. For the details of the Lebesgue spaces with variable exponents, the reader is referred to [3, 4].

DEFINITION 2. Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. We write $p(\cdot) \in \mathcal{M}_+$ if $M^+ : L^{p(\cdot)} \rightarrow L^{p(\cdot)}$ is bounded. We write $p(\cdot) \in \mathcal{M}_-$ if $M^- : L^{p(\cdot)} \rightarrow L^{p(\cdot)}$ is bounded.

We now give some concrete conditions that guarantee $p(\cdot) \in \mathcal{M}_+ \cup \mathcal{M}_-$ from [29].

DEFINITION 3. Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function with $1 < p_- \leq p_+ < \infty$. We write $p(\cdot) \in \mathcal{L}_+$ if there is a constant $C > 0$ such that

$$p(y) \geq p(x) + \frac{C}{\ln(y-x)}$$

for all $x, y \in \mathbb{R}$ with $0 < y - x \leq \frac{1}{2}$.

We write $p(\cdot) \in \mathcal{L}_-$ if $\tilde{p}(\cdot) \in \mathcal{L}_+$ where $\tilde{p}(x) = p(-x)$, $x \in \mathbb{R}$.

DEFINITION 4. Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. We write $p(\cdot) \in \mathcal{P}$ if there is a constant $c > 0$ such that

$$\int_{\{x: p(x) \neq 0\}} c^{\frac{1}{p(x)}} dx < \infty.$$

We have the following result from [29, Theorem 1].

THEOREM 1. *Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function.*

1. *If $p(\cdot) \in \mathcal{L}_+$ and there exists a non-increasing function $q(\cdot)$ satisfying $1 < q_- \leq q_+ < \infty$, $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $p(\cdot) \in \mathcal{M}_+$.*
2. *If $p(\cdot) \in \mathcal{L}_-$ and there exists a non-decreasing function $q(\cdot)$ satisfying $1 < q_- \leq q_+ < \infty$, $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $p(\cdot) \in \mathcal{M}_-$.*

We have another condition that guarantees $p(\cdot) \in \mathcal{M}_+ \cup \mathcal{M}_-$. We need to recall some notations from [29] to present this condition.

Let $e_0 = 1$ and $e_{k+1} = e^{e_k}$, $k \in \mathbb{N} \setminus \{0\}$. Write $\ln_0 x = x$. Let $k \in \mathbb{N} \setminus \{0\}$. For any $x \in (e_k, \infty)$, write $\ln_k x = \ln(\ln_{k-1} x)$. For any $\alpha > 0$, write

$$b_{k,\alpha}(x) = -\frac{1}{\alpha} \frac{d}{dx} (\ln_k^{-\alpha} x), \quad x \in (e_k, \infty).$$

DEFINITION 5. Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function with $1 < p_- \leq p_+ < \infty$. We write $p(\cdot) \in \mathfrak{Q}$ if

1. $p(x) = p(-x)$ for all $x \in \mathbb{R}$,
2. there are $\alpha, C > 0$ and $k \in \mathbb{N} \setminus \{0\}$ such that $p(\cdot)$ is monotone on (e_k, ∞) and

$$\left| \frac{dp(x)}{dx} \right| \leq C b_{k,\alpha}(x), \quad x \in (e_k, \infty).$$

The following result is from [29, Theorem 2].

THEOREM 2. *Let $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function.*

1. *If $p(\cdot) \in \mathcal{L}_+$ and there exists a $q(\cdot) \in \mathfrak{Q}$ such that $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $p(\cdot) \in \mathcal{M}_+$.*
2. *If $p(\cdot) \in \mathcal{L}_-$ and there exists a $q(\cdot) \in \mathfrak{Q}$ such that $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $p(\cdot) \in \mathcal{M}_-$.*

We now present the definition of the one-sided Herz spaces with variable exponents studied in this paper. For any $m \in \mathbb{N} \setminus \{0\}$, define

$$D_0 = (-1, 1), \quad D_m = [2^{m-1}, 2^m), \quad D_{-m} = (-2^m, -2^{m-1}].$$

DEFINITION 6. Let $\theta \in (0, \infty)$, $\alpha \in (-\infty, \infty)$ and $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. The one-sided Herz space with variable exponent $\dot{K}_{p(\cdot), \theta}^\alpha$ consists of all Lebesgue measurable functions f satisfying

$$\|f\|_{\dot{K}_{p(\cdot), \theta}^\alpha} = \left(\sum_{k \in \mathbb{Z}} \|2^{k\alpha} f \chi_{D_k}\|_{L^{p(\cdot)}}^\theta \right)^{\frac{1}{\theta}} < \infty.$$

When $\alpha > 0$, the Herz space with variable exponent studied in [1, 10, 11] is a subspace of $\dot{K}_{p(\cdot), \theta}^\alpha$. Whenever $p(\cdot) = p$, $p \in (1, \infty)$, is a constant function, we denote $\dot{K}_{p(\cdot), \theta}^\alpha$ by $\dot{K}_{p, \theta}^\alpha$. It is an extension of the classical Herz spaces. For the studies of the classical Herz space and its generalizations, see [6, 9, 14, 16, 20, 21, 22, 30, 31, 35].

A number of important results in harmonic analysis, such as the boundedness of the Hardy-Littlewood maximal function, the singular integral operators and bilinear operators had been extended to the classical Herz spaces and Herz spaces with variable exponents studied in [1, 10, 11], see [12, 13, 17, 18, 19].

The following result assures that for any $f \in \dot{K}_{p(\cdot), \theta}^\alpha$, $M^+ f$ and $M^- f$ are Lebesgue measurable.

PROPOSITION 1. *Let $\theta \in (0, \infty)$, $\alpha \in (-\infty, \infty)$ and $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. For any $f \in \dot{K}_{p(\cdot), \theta}^\alpha$, $M^+ f, M^- f$ are Lebesgue measurable functions.*

Proof. We show that for any $m \in \mathbb{N}$ and $f \in \dot{K}_{p(\cdot), \theta}^\alpha$, $\int_{-2^m}^{2^m} |f(y)| dy$ is finite.

The Hölder inequality for $L^{p(\cdot)}$ [3, Theorem 2.36] gives

$$\int_{-2^m}^{2^m} |f(y)| dy = \sum_{k=-m}^m \int_{\mathbb{R}} \chi_{D_k}(y) |f(y)| dy \leq C \sum_{k=-m}^m \|\chi_{D_k}\|_{L^{p'(\cdot)}} \|\chi_{D_k} f\|_{L^{p(\cdot)}}.$$

When $\theta \in (1, \infty)$, the Hölder inequality and the inequality $\|\chi_{D_k}\|_{L^{p'(\cdot)}} \leq \|\chi_{(-2^m, 2^m)}\|_{L^{p'(\cdot)}}$ yield

$$\begin{aligned}
 & \int_{-2^m}^{2^m} |f(y)| dy \\
 & \leq C \left(\sum_{k=-m}^m (2^{-k\alpha} \|\chi_{D_k}\|_{L^{p'(\cdot)}})^{\theta'} \right)^{\frac{1}{\theta'}} \left(\sum_{k=-m}^m (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^{\theta} \right)^{\frac{1}{\theta}} \\
 & \leq C \left(\sum_{k=-m}^m (2^{-k\alpha} \|\chi_{(-2^m, 2^m)}\|_{L^{p'(\cdot)}})^{\theta'} \right)^{\frac{1}{\theta'}} \|f\|_{\dot{K}_{p(\cdot), \theta}^{\alpha}} \\
 & \leq C (2^m)^{\frac{1}{\theta'}} 2^{m|\alpha|} \|\chi_{(-2^m, 2^m)}\|_{L^{p'(\cdot)}} \|f\|_{\dot{K}_{p(\cdot), \theta}^{\alpha}} < \infty.
 \end{aligned}$$

When $\theta \in (0, 1]$, the θ -inequality gives

$$\begin{aligned}
 & \int_{-2^m}^{2^m} |f(y)| dy \leq C \sum_{k=-m}^m \|\chi_{D_k}\|_{L^{p'(\cdot)}} \|\chi_{D_k} f\|_{L^{p(\cdot)}} \\
 & \leq C 2^{m|\alpha|} \|\chi_{(-2^m, 2^m)}\|_{L^{p'(\cdot)}} \left(\sum_{k=-m}^m (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^{\theta} \right)^{\frac{1}{\theta}} \\
 & \leq C 2^{m|\alpha|} \|\chi_{(-2^m, 2^m)}\|_{L^{p'(\cdot)}} \|f\|_{\dot{K}_{p(\cdot), \theta}^{\alpha}} < \infty.
 \end{aligned}$$

Thus, $\dot{K}_{p(\cdot), \theta}^{\alpha} \subset L_{\text{loc}}^1$. As the one-sided maximal operators M^+ and M^- are defined for locally integrable functions, M^+ and M^- are defined on $\dot{K}_{p(\cdot), \theta}^{\alpha}$.

We now use the idea from [7, p.91] to show that for any $f \in \dot{K}_{p(\cdot), \theta}^{\alpha}$, $M^+ f$ and $M^- f$ are Lebesgue measurable.

For any $x \in \mathbb{R}$ and $t > 0$, $F_+(x, t) = \frac{1}{t} \int_x^{x+t} |f(y)| dy$ and $F_-(x, t) = \frac{1}{t} \int_{x-t}^x |f(y)| dy$ are continuous in t for each x . In view of the Fubini's theorem, $F_+(x, t)$ and $F_-(x, t)$ are measurable in x for each t . As $F_+(x, t)$ and $F_-(x, t)$ are continuous in r , we see that

$$\begin{aligned}
 \sup_{t \in \mathbb{Q} \cap (0, \infty)} F_+(x, t) &= \sup_{t > 0} F_+(x, t) = M^+ f(x) \\
 \sup_{t \in \mathbb{Q} \cap (0, \infty)} F_-(x, t) &= \sup_{t > 0} F_-(x, t) = M^- f(x).
 \end{aligned}$$

Therefore, for any $f \in \dot{K}_{p(\cdot), \theta}^{\alpha}$, $M^+ f$ and $M^- f$ are Lebesgue measurable. \square

The above proposition assures that it does make sense to study the boundedness of the one-sided maximal operators M^+ and M^- on the one-sided Herz space with variable exponent $\dot{K}_{p(\cdot), \theta}^{\alpha}$.

3. Main results

The main result of this paper, the boundedness of the one-sided maximal operators on the one-sided Herz spaces with variable exponent $\dot{K}_{p(\cdot),\theta}^\alpha$, is established in this section.

THEOREM 3. *Let $\theta \in (0, \infty)$, $\alpha \in (0, \infty)$, $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. If $1 < p_- \leq p_+ < \infty$ and $p(\cdot) \in \mathcal{M}_+$, then $M^+ : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ is bounded.*

Proof. We first establish that for any $f \in \dot{K}_{p(\cdot),\theta}^\alpha$ and $x \in D_m$, $m \in \mathbb{Z}$, we have

$$\chi_{D_m}(x)M^+f(x) \leq M^+(\chi_{\cup_{k=m}^\infty D_k}f)(x), \quad x \in \mathbb{R}. \quad (1)$$

When $x \notin D_m$, we have

$$\chi_{D_m}(x)M^+f(x) = 0 \leq M^+(\chi_{\cup_{k=m}^\infty D_k}f)(x). \quad (2)$$

Whenever $x \in D_m$, we find that for any $h > 0$

$$\begin{aligned} \frac{1}{h} \int_x^{x+h} |f(y)| dy &= \frac{1}{h} \int_x^{x+h} \chi_{\cup_{k=m}^\infty D_k}(y) |f(y)| dy \\ &\leq M^+(\chi_{\cup_{k=m}^\infty D_k}f)(x) \end{aligned} \quad (3)$$

because $x \in D_m$ asserts that $(x, x+h) \subset (x, \infty) \subset \cup_{k=m}^\infty D_k$. By taking the supremum over $h > 0$ on (3), we find that for any $x \in D_m$

$$\chi_{D_m}(x)M^+f(x) \leq M^+(\chi_{\cup_{k=m}^\infty D_k}f)(x). \quad (4)$$

Consequently, (2) and (4) yield (1).

Next, we apply the norm $\|\cdot\|_{L^{p(\cdot)}}$ on both sides of (1), we obtain

$$\|\chi_{D_m}M^+f\|_{L^{p(\cdot)}} \leq \|M^+(\chi_{\cup_{k=m}^\infty D_k}f)\|_{L^{p(\cdot)}}.$$

As $\alpha > 0$, the Hölder inequality on ℓ^θ for $\theta \geq 1$ and the θ -inequality for $0 < \theta < 1$ yield $\chi_{\cup_{k=m}^\infty D_k}f \in L^{p(\cdot)}$. Consequently, $p(\cdot) \in \mathcal{M}_+$ gives

$$\|\chi_{D_m}M^+f\|_{L^{p(\cdot)}} \leq C \|\chi_{\cup_{k=m}^\infty D_k}f\|_{L^{p(\cdot)}} \leq C \sum_{k=m}^\infty \|\chi_{D_k}f\|_{L^{p(\cdot)}} \quad (5)$$

for some $C > 0$.

We now consider the case when $\theta \in (1, \infty)$. (5) gives

$$\begin{aligned} \sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m}M^+f\|_{L^{p(\cdot)}})^\theta &\leq C \sum_{m \in \mathbb{Z}} \left(\sum_{k=m}^\infty 2^{m\alpha} \|\chi_{D_k}f\|_{L^{p(\cdot)}} \right)^\theta \\ &= C \sum_{m \in \mathbb{Z}} \left(\sum_{k=m}^\infty 2^{(m-k)\alpha} 2^{k\alpha} \|\chi_{D_k}f\|_{L^{p(\cdot)}} \right)^\theta. \end{aligned}$$

We now use the idea in [32, Proposition 1.2]. The Hölder inequality guarantees that

$$\begin{aligned} & \sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta \\ & \leq C \sum_{m \in \mathbb{Z}} \left(\sum_{k=m}^{\infty} 2^{\frac{\theta}{2}(m-k)\alpha} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \right) \left(\sum_{k=m}^{\infty} 2^{\frac{\theta'}{2}(m-k)\alpha} \right)^{\frac{\theta}{\theta'}}. \end{aligned}$$

As $\alpha > 0$, we find that $\sum_{k=m}^{\infty} 2^{\frac{\theta'}{2}(m-k)\alpha} < C$ for some $C > 0$ independent of m . Therefore, by interchanging the summations, we obtain

$$\begin{aligned} \sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta & \leq C \sum_{m \in \mathbb{Z}} \sum_{k=m}^{\infty} 2^{\frac{\theta}{2}(m-k)\alpha} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \\ & \leq C \sum_{k \in \mathbb{Z}} \sum_{m=-\infty}^k 2^{\frac{\theta}{2}(m-k)\alpha} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \\ & = C \sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \sum_{m=-\infty}^k 2^{\frac{\theta}{2}(m-k)\alpha} \\ & \leq C \sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \end{aligned}$$

where we have the last inequality because $\alpha > 0$. Hence,

$$\begin{aligned} \|M^+ f\|_{\dot{K}_{p(\cdot),\theta}^\alpha} & = \left(\sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta \right)^{\frac{1}{\theta}} \\ & \leq C \left(\sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \right)^{\frac{1}{\theta}} = C \|f\|_{\dot{K}_{p(\cdot),\theta}^\alpha}. \end{aligned}$$

Therefore, we establish the boundedness of $M^+ : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ when $\theta \in (1, \infty)$.

We consider the case $\theta \in (0, 1]$. The θ -inequality and (5) yield

$$\begin{aligned} \sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta & \leq C \sum_{m \in \mathbb{Z}} \left(\sum_{k=m}^{\infty} 2^{m\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}} \right)^\theta \\ & = C \sum_{m \in \mathbb{Z}} \left(\sum_{k=m}^{\infty} 2^{(m-k)\alpha} 2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}} \right)^\theta \\ & \leq C \sum_{m \in \mathbb{Z}} \sum_{k=m}^{\infty} 2^{\theta(m-k)\alpha} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta. \end{aligned}$$

By interchanging the summations, we get

$$\begin{aligned} \sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta &\leq C \sum_{k \in \mathbb{Z}} \sum_{m=-\infty}^k 2^{\theta(m-k)\alpha} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \\ &= C \sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \sum_{m=-\infty}^k 2^{\theta(m-k)\alpha}. \end{aligned}$$

Since $\alpha > 0$, we find that $\sum_{m=-\infty}^k 2^{\theta(m-k)\alpha} < C$ for some $C > 0$ independent of k . Consequently,

$$\begin{aligned} \|M^+ f\|_{\dot{K}_{p(\cdot),\theta}^\alpha} &= \left(\sum_{m \in \mathbb{Z}} (2^{m\alpha} \|\chi_{D_m} M^+ f\|_{L^{p(\cdot)}})^\theta \right)^{\frac{1}{\theta}} \\ &\leq C \left(\sum_{k \in \mathbb{Z}} (2^{k\alpha} \|\chi_{D_k} f\|_{L^{p(\cdot)}})^\theta \right)^{\frac{1}{\theta}} = C \|f\|_{\dot{K}_{p(\cdot),\theta}^\alpha}. \end{aligned}$$

Thus, we establish the boundedness of $M^+ : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ when $\theta \in (0, 1]$. \square

We also have the boundedness of M^- on the one-sided Herz spaces with variable exponents $\dot{K}_{p(\cdot),\theta}^\alpha$.

THEOREM 4. *Let $\theta \in (0, \infty)$, $\alpha \in (-\infty, 0)$, $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function. If $1 < p_- \leq p_+ < \infty$ and $p(\cdot) \in \mathcal{M}_-$, then $M^- : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ is bounded.*

As the proof of the above result is similar to the proof of Theorem 3, for simplicity, we omit the proof.

In view of Theorems 1, 2, 3 and 4, we have the following boundedness results for the one-sided maximal operators M^+ and M^- on the one-sided Herz space with variable exponent $\dot{K}_{p(\cdot),\theta}^\alpha$.

COROLLARY 1. *Let $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function.*

1. *If $\alpha \in (0, \infty)$, $p(\cdot) \in \mathcal{L}_+$ and there exists a non-increasing function $q(\cdot)$ satisfying $1 < q_- \leq q_+ < \infty$, $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $M^+ : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ is bounded.*
2. *If $\alpha \in (-\infty, 0)$, $p(\cdot) \in \mathcal{L}_-$ and there exists a non-decreasing function $q(\cdot)$ satisfying $1 < q_- \leq q_+ < \infty$, $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $M^- : \dot{K}_{p(\cdot),\theta}^\alpha \rightarrow \dot{K}_{p(\cdot),\theta}^\alpha$ is bounded.*

COROLLARY 2. *Let $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $p(\cdot) : \mathbb{R} \rightarrow (1, \infty)$ be a Lebesgue measurable function.*

1. If $\alpha \in (0, \infty)$, $p(\cdot) \in \mathcal{L}_+$ and there exists a $q(\cdot) \in \mathfrak{Q}$ such that $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $M^+ : \dot{K}_{p(\cdot), \theta}^\alpha \rightarrow \dot{K}_{p(\cdot), \theta}^\alpha$ is bounded.
2. If $\alpha \in (-\infty, 0)$, $p(\cdot) \in \mathcal{L}_-$ and there exists a $q(\cdot) \in \mathfrak{Q}$ such that $|p(\cdot) - q(\cdot)| \in \mathcal{P}$, then $M^- : \dot{K}_{p(\cdot), \theta}^\alpha \rightarrow \dot{K}_{p(\cdot), \theta}^\alpha$ is bounded.

Moreover, Theorems 3 and 4 also give the boundedness result for M^+ and M^- on the one-sided Herz spaces $\dot{K}_{p, \theta}^\alpha$.

COROLLARY 3. Let $\theta \in (0, \infty)$, $\alpha \in \mathbb{R}$ and $p \in (1, \infty)$.

1. If $\alpha \in (0, \infty)$, then $M^+ : \dot{K}_{p, \theta}^\alpha \rightarrow \dot{K}_{p, \theta}^\alpha$ is bounded.
2. If $\alpha \in (-\infty, 0)$, then $M^- : \dot{K}_{p, \theta}^\alpha \rightarrow \dot{K}_{p, \theta}^\alpha$ is bounded.

The mapping properties of M^- on the weighted Herz spaces were given in [14].

Declaration

Funding. Not applicable.

Conflicts of interest/Competing interests. Not applicable.

Availability of data and material. Not applicable.

Code availability. Not applicable.

Acknowledgements. The author thanks the reviewers for their valuable suggestions which improve the presentation of this paper.

REFERENCES

- [1] A. ALMEIDA AND D. DRIHEM, *Maximal, potential and singular type operators on Herz spaces with variable exponents*, J. Math. Anal. Appl., **394**, (2012), 781–795.
- [2] C. BENNETT AND R. SHARPLEY, *Interpolation of Operators*, Academic Press (1988).
- [3] D. CRUZ-URIBE AND A. FIORENZA, *Variable Lebesgue Spaces*, Birkhäuser, 2013.
- [4] L. DIENING, P. HARJULEHTO, P. HÄSTÖ AND M. RUŽIČKA, *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer, 2011.
- [5] D. E. EDMUNDS, V. KOKILASHVILI AND A. MESKHI, *On one-sided operators in variable exponent Lebesgue spaces*, Proc. A. Razmadze Math. Inst., **144** (2007), 126–131.
- [6] H. FEICHTINGER AND F. WEISZ, *Herz spaces and summability of Fourier transforms*, Math. Nachr., **281** (2008), 309–324.
- [7] G. FOLLAND, *Real Analysis: Modern Techniques And Their Applications*, John Wiley & Sons, New York, (1984).
- [8] G. HARDY AND J. LITTLEWOOD, *A maximal theorem with function-theoretic applications*, Acta Math., **54** (1930), 81–116.
- [9] C. HERZ, *Lipschitz spaces and Bernstein's theorem on absolutely convergent Fourier transforms*, J. Math. Mech., **18** (1968/69), 283–323.

- [10] K.-P. HO, *Extrapolation to Herz spaces with variable exponents and applications*, Revista Mat. Complut., **33**, (2020), 437–463.
- [11] K.-P. HO, *Sublinear operators on Herz-Hardy spaces with variable exponents*, Math. Nachr., **295** (2022), 876–889.
- [12] K.-P. HO, *Bilinear operators on ball Banach function spaces*, J. Geom. Anal. **34** (2024), Article no. 337.
- [13] K.-P. HO, *Herz spaces and integral operators on Morrey spaces*, (preprint).
- [14] K.-P. HO, *Localized operators on weighted Herz spaces*, Math. Nachr. **297** (2024) 4067–4080.
- [15] L. HUANG, F. WEISZ, D. YANG AND W. YUAN, *Summability of Fourier transforms on mixed-norm Lebesgue spaces via associated Herz spaces*, Anal. Appl. (Singap.) **21** (2023), 279–328.
- [16] M. IZUKI AND Y. SAWANO, *The Haar wavelet characterization of weighted Herz spaces and greediness of the Haar wavelet basis*, J. Math. Anal. Appl. **362** (2010), 140–155.
- [17] M. IZUKI, *Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization*, Anal. Math., **36** (2010), 33–50.
- [18] M. IZUKI AND T. NOI, *Boundedness of some integral operators and commutators on generalized Herz spaces with variable exponents*, http://sci.osaka-cu.ac.jp/math/OCAMI/preprint/2011/11_15.pdf.
- [19] M. IZUKI, T. NOI AND Y. SAWANO, *Extrapolation to two-weighted Herz spaces with three variable exponents*, Adv. Oper. Theory **9** (2024) Article No. 44.
- [20] Y. LI, D. YANG AND L. HUANG, *Real-variable theory of Hardy spaces associated with generalized Herz spaces of Rafeiro and Samko*, Lecture Notes in Mathematics, No. 2320, Springer (2022).
- [21] S. LU AND D. YANG, *Hardy-Littlewood-Sobolev theorem of fractional integration on Herz-type spaces and its applications*, Canad. J. Math., **48** (1996), 363–380.
- [22] S. LU, D. YANG AND G. HU, *Herz type spaces and their applications*, Science Press, Beijing, 2008.
- [23] S. LU, Z. FU, F. ZHAO AND S. SHI, *Hardy Operators on Euclidean Spaces and Related Topics*, World Scientific Press, Singapore (2023).
- [24] F. MARTÍN-REYES, P. ORTEGA AND A. DE LA TORRE, *Weighted inequalities for one-sided maximal functions*, Trans. Amer. Math. Soc. **319** (1990), 517–534.
- [25] F. MARTÍN-REYES, *Weights, one-sided operators, singular integrals and ergodic theorems*, Nonlinear Analysis, Function Spaces and Applications, Praha: Prometheus Publishing House, (1994), 103–137.
- [26] F. MARTÍN-REYES AND A. DE LA TORRE, *Some weighted inequalities for general one-sided maximal operators*, Studia Math. **122** (1997), 1–14.
- [27] A. NEKVINDA, *Hardy-Littlewood maximal operator on $L^{p(\cdot)}(\mathbb{R}^n)$* , Math. Inequal. Appl. **7** (2004), 255–265.
- [28] A. NEKVINDA, *Maximal operator on variable Lebesgue spaces for almost monotone radial exponent*, J. Math. Anal. Appl. **337** (2008), 1345–1365.
- [29] A. NEKVINDA, *A note on one-sided maximal operator in $L^{p(\cdot)}(\mathbb{R})$* , Math. Inequal. Appl. **13** (2010), 887–897.
- [30] M. A. RAGUSA, *Homogeneous Herz spaces and regularity results*, Nonlinear Anal. **71** (2009), 1909–1914.
- [31] M. A. RAGUSA, *Parabolic Herz spaces and their applications*, Appl. Math. Lett. **25** (2012), 1270–1273.
- [32] Y. SAWANO, *Theory of Besov Spaces*, Developments in Mathematics, vol. 56, Berlin, Springer.
- [33] Y. SAWANO, K.-P. HO, D. YANG AND S. YANG, *Hardy spaces for ball quasi-Banach function spaces*, Dissertationes Math. (Rozprawy Mat.) **525** (2017), 1–102.
- [34] E. SAWYER, *Weighted inequalities for the one-sided Hardy-Littlewood maximal functions*, Trans. Amer. Math. Soc. **297** (1986), 53–61.
- [35] Y. ZHAO, D. YANG AND Y. ZHANG, *Mixed-norm Herz spaces and their applications in related Hardy spaces*, Anal. Appl. (Singap.) **21** (2023), 1131–1222.

(Received July 18, 2024)

Kwok-Pun Ho
 Department of Mathematics and Information Technology
 The Education University of Hong Kong
 10 Lo Ping Road, Tai Po, Hong Kong, China
 e-mail: vkpho@eduhk.hk