

SOME PROPERTIES OF GENERALIZATION CLASSES OF ANALYTIC FUNCTIONS

HATUN ÖZLEM GÜNEY*, DANIEL BREAZ AND SHIGEYOSHI OWA

(Communicated by S. Varošanec)

Abstract. Let $\overline{\mathcal{A}}(n)$ be the class of functions $f(z)$ of the form

$$f(z) = z + \sum_{k=1}^{\infty} a_{1+\frac{k}{n}} z^{1+\frac{k}{n}} \quad (n = 1, 2, 3, \dots)$$

which are analytic in the open unit disc \mathbb{U} . If $a_{1+\frac{k}{n}} = 0$ for $k \neq n, 2n, 3n, \dots$, then $f(z)$ is given by $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$. For such functions $f(z) \in \overline{\mathcal{A}}(n)$, some generalization classes $\overline{\mathcal{S}}^*(n, \alpha)$, $\overline{\mathcal{C}}(n, \alpha)$ and $\overline{\mathcal{B}}(n, \alpha)$ are defined. The object of present paper is to discuss some interesting properties of $f(z) \in \overline{\mathcal{A}}(n)$ concerning with subordinations and strongly functions.

1. Introduction

Let n be a natural number and $\overline{\mathcal{A}}(n)$ be the class of functions $f(z)$ of the form

$$f(z) = z + \sum_{k=1}^{\infty} a_{1+\frac{k}{n}} z^{1+\frac{k}{n}} \quad (n = 1, 2, 3, \dots) \quad (1)$$

that are analytic in the open unit disc $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. Here, we take the principal value for $\sqrt[n]{z}$. Let $\overline{\mathcal{S}}^*(n, \alpha)$ denote the subclass of $\overline{\mathcal{A}}(n)$ consisting of $f(z)$ that satisfies

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \alpha, \quad (z \in \mathbb{U}) \quad (2)$$

for some α ($0 \leq \alpha < 1$). Also, $\overline{\mathcal{C}}(n, \alpha)$ is the subclass of $\overline{\mathcal{A}}(n)$ consisting of $f(z)$ that satisfies

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha, \quad (z \in \mathbb{U}) \quad (3)$$

for some α ($0 \leq \alpha < 1$). By the definitions for $\overline{\mathcal{S}}^*(n, \alpha)$ and $\overline{\mathcal{C}}(n, \alpha)$, we see that $f(z) \in \overline{\mathcal{C}}(n, \alpha)$ if and only if $zf'(z) \in \overline{\mathcal{S}}^*(n, \alpha)$, and that $f(z) \in \overline{\mathcal{S}}^*(n, \alpha)$ if and only

Mathematics subject classification (2020): Primary 30C45; Secondary 30C80.

Keywords and phrases: Analytic function, subordination, argument property, strongly function.

* Corresponding author.

if $\int_0^z \frac{f(t)}{t} dt \in \overline{\mathcal{C}}(n, \alpha)$. If $a_{1+\frac{k}{n}} = 0$ for $k \neq n, 2n, 3n, \dots$, then $f(z) \in \overline{\mathcal{A}}(n)$ can be written by

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (4)$$

and write that $f(z) \in \mathcal{A}$.

Let us consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$\begin{aligned} f(z) &= \frac{z}{(1 - \sqrt[n]{z})^{2n(1-\alpha)}} \\ &= z + \sum_{k=1}^{\infty} \frac{\prod_{j=1}^k (j + 2n(1-\alpha) - 1)}{k!} z^{1+\frac{k}{n}} \end{aligned} \quad (5)$$

with $0 \leq \alpha < 1$. If $n = 2$, then

$$\begin{aligned} f(z) &= \frac{z}{(1 - \sqrt{z})^{4(1-\alpha)}} \\ &= z + \sum_{k=1}^{\infty} \frac{\prod_{j=1}^k (j + 3 - 4\alpha)}{k!} z^{1+\frac{k}{2}} \end{aligned} \quad (6)$$

and $f(z) \in \overline{\mathcal{S}}^*(2, \alpha)$ in Güney, Breaz and Owa [3]. Also, $n = 3$, then

$$\begin{aligned} f(z) &= \frac{z}{(1 - \sqrt[3]{z})^{6(1-\alpha)}} \\ &= z + \sum_{k=1}^{\infty} \frac{\prod_{j=1}^k (j + 5 - 6\alpha)}{k!} z^{1+\frac{k}{3}} \end{aligned} \quad (7)$$

and $f(z) \in \overline{\mathcal{S}}^*(3, \alpha)$ in Güney and Owa [2]. Also, $f(z)$ given by (5) satisfies

$$Re \left\{ \frac{zf'(z)}{f(z)} \right\} = Re \left(\frac{1 + (1-2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \right) > \alpha, (z \in \mathbb{U}). \quad (8)$$

Thus we know that $f(z)$ given by (5) is in the class $\overline{\mathcal{S}}^*(n, \alpha)$. If we consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f'(z) = \frac{1}{(1 - \sqrt[n]{z})^{2n(1-\alpha)}} \quad (0 \leq \alpha < 1), \quad (9)$$

then $f(z)$ satisfies

$$Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} = Re \left(\frac{1 + (1-2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \right) > \alpha, (z \in \mathbb{U}) \quad (10)$$

and $f(z) \in \overline{\mathcal{C}}(n, \alpha)$. Further, if $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f(z) = (2\alpha - 1)z + 2(1 - \alpha) \int_0^z \frac{1}{1 - \sqrt[n]{t}} dt, \quad (11)$$

then $f(z)$ satisfies

$$Re f'(z) = Re \left(\frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \right) > \alpha, (z \in \mathbb{U}). \quad (12)$$

We denote by $f(z) \in \overline{\mathcal{R}}(n, \alpha)$ if $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$Re f'(z) > \alpha, (z \in \mathbb{U}) \quad (13)$$

for $0 \leq \alpha < 1$.

For analytic functions $f(z)$ and $F(z)$ in \mathbb{U} , we introduce that $f(z)$ is subordinate to $F(z)$, written $f(z) \prec F(z)$ ($z \in \mathbb{U}$), if there exists a function $w(z)$ analytic in \mathbb{U} with $w(0) = 0$ and $|w(z)| < 1$ ($z \in \mathbb{U}$) and such that $f(z) = F(w(z))$ (see [6]). With the definition for subordinations, if $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\frac{zf'(z)}{f(z)} \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, (z \in \mathbb{U}), \quad (14)$$

with $0 \leq \alpha < 1$, then $f(z) \in \overline{\mathcal{S}}^*(n, \alpha)$, and if $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$1 + \frac{zf''(z)}{f'(z)} \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, (z \in \mathbb{U}), \quad (15)$$

with $0 \leq \alpha < 1$, then $f(z) \in \overline{\mathcal{C}}(n, \alpha)$. Further, if $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$f'(z) \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, (z \in \mathbb{U}), \quad (16)$$

with $0 \leq \alpha < 1$, then $f(z) \in \overline{\mathcal{R}}(n, \alpha)$.

EXAMPLE 1. We consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f(z) = \frac{z}{1 - \sqrt[n]{z}} = z + \sum_{k=1}^{\infty} z^{1+\frac{k}{n}}. \quad (17)$$

Letting $z = e^{i\theta}$ ($0 \leq \theta < 2\pi$), we have

$$\begin{aligned} Re f(z) &= Re \left(\frac{e^{i\theta}}{1 - e^{i\frac{\theta}{n}}} \right) \\ &= Re \left(\frac{e^{i(1-\frac{1}{n})\theta}}{e^{-i\frac{\theta}{2n}} - e^{i\frac{\theta}{2n}}} \right) \\ &= -\frac{1}{2} \left(\frac{\sin \left(\frac{2n-1}{2n} \theta \right)}{\sin \left(\frac{1}{2n} \theta \right)} \right). \end{aligned} \quad (18)$$

If $n = 1$, then

$$Re f(z) = -\frac{1}{2} \left(\frac{\sin \frac{\theta}{2}}{\sin \frac{\theta}{n}} \right) = -\frac{1}{2}. \quad (19)$$

If $n = 2$, then

$$\begin{aligned} \operatorname{Ref}(z) &= -\frac{1}{2} \left(\frac{\sin \frac{3}{4}\theta}{\sin \frac{\theta}{4}} \right) = -\frac{1}{2} \left(\frac{\sin \left(\frac{\theta}{4} + \frac{\theta}{2} \right)}{\sin \frac{\theta}{4}} \right) \\ &= -\frac{1}{2} \left(4 \cos^2 \frac{\theta}{4} - 1 \right) \\ &\geq -\frac{3}{2}. \end{aligned} \quad (20)$$

If $n = 3$, then

$$\begin{aligned} \operatorname{Ref}(z) &= -\frac{1}{2} \left(\frac{\sin \frac{5}{6}\theta}{\sin \frac{\theta}{6}} \right) = -\frac{1}{2} \left(\frac{\sin \left(\frac{\theta}{6} + \frac{2}{3}\theta \right)}{\sin \frac{\theta}{6}} \right) \\ &= -\frac{1}{2} \left(4 \cos^2 \frac{\theta}{3} + 2 \cos \frac{\theta}{3} - 1 \right) \\ &\geq -\frac{5}{2}. \end{aligned} \quad (21)$$

If $n = 4$, then

$$\begin{aligned} \operatorname{Ref}(z) &= -\frac{1}{2} \left(\frac{\sin \frac{7}{8}\theta}{\sin \frac{\theta}{8}} \right) = -\frac{1}{2} \left(\frac{\sin \left(\frac{\theta}{8} + \frac{3}{4}\theta \right)}{\sin \frac{\theta}{8}} \right) \\ &= -\frac{1}{2} \left(4 \cos^2 \frac{\theta}{4} \left(2 \cos^2 \frac{\theta}{4} - 1 \right) + (4 \cos^2 \theta - 1) \right) \\ &\geq -\frac{7}{2}. \end{aligned} \quad (22)$$

If $n = 5$, then

$$\begin{aligned} \operatorname{Ref}(z) &= -\frac{1}{2} \left(\frac{\sin \frac{9}{10}\theta}{\sin \frac{\theta}{10}} \right) = -\frac{1}{2} \left(\frac{\sin \left(\frac{\theta}{10} + \frac{4}{5}\theta \right)}{\sin \frac{\theta}{10}} \right) \\ &= -\frac{1}{2} \left(4 \cos^2 \frac{\theta}{5} \left(4 \cos^2 \frac{\theta}{5} - 3 \right) + 4 \cos \frac{\theta}{5} \left(2 \cos^2 \frac{\theta}{5} - 1 \right) + 1 \right) \\ &\geq -\frac{9}{2}. \end{aligned} \quad (23)$$

Therefore, we can be expected that (18) gives us

$$\begin{aligned} \operatorname{Ref}(z) &= -\frac{1}{2} \left(\frac{\sin \left(\frac{\theta}{2n} + \frac{n-1}{n}\theta \right)}{\sin \frac{\theta}{2n}} \right) \\ &\geq -\frac{2n-1}{2}. \end{aligned} \quad (24)$$

2. Some applications of subordinations

To consider some applications of subordinations, we need the following lemma due to Suffridge [10].

LEMMA 1. Let $p(z)$ be analytic in \mathbb{U} with $p(0) = 1$ and $h(z)$ be analytic and starlike in \mathbb{U} . If

$$zp'(z) \prec h(z), (z \in \mathbb{U}), \quad (25)$$

then

$$p(z) \prec \int_0^z \frac{h(t)}{t} dt, (z \in \mathbb{U}). \quad (26)$$

Applying the above lemma, we have the following theorem.

THEOREM 1. Let $f(z) \in \overline{\mathcal{A}}(n)$ and $h(z)$ be given by

$$h(z) = \frac{z}{(1 - \sqrt[n]{z})^{2n(1-\alpha)}}, (z \in \mathbb{U}) \quad (27)$$

with $0 \leq \alpha < 1$. If $f(z)$ satisfies

$$zf''(z) \prec h(z) \quad (28)$$

then

$$f'(z) \prec \int_0^z \frac{h(t)}{t} dt = \int_0^z \frac{1}{(1 - \sqrt[n]{t})^{2n(1-\alpha)}} dt, (z \in \mathbb{U}). \quad (29)$$

Proof. We define a function $p(z) = f'(z)$ for $f(z) \in \overline{\mathcal{A}}(n)$. Then $p(z)$ is analytic in \mathbb{U} and $p(0) = 1$. Also, $h(z)$ given by (27) satisfies

$$\operatorname{Re} \left(\frac{zh'(z)}{h(z)} \right) = \operatorname{Re} \left(\frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \right) > \alpha, (z \in \mathbb{U}). \quad (30)$$

Using Lemma 1, we see if

$$zp'(z) = zf''(z) \prec h(z), (z \in \mathbb{U}), \quad (31)$$

then

$$p(z) = f'(z) \prec \int_0^z \frac{h(t)}{t} dt = \int_0^z \frac{1}{(1 - \sqrt[n]{t})^{2n(1-\alpha)}} dt, (z \in \mathbb{U}). \quad \square \quad (32)$$

Letting $n = 2$ in Theorem 1, we see the following corollary.

COROLLARY 1. If $f(z) \in \overline{\mathcal{A}}(2)$ satisfies

$$zf''(z) \prec \frac{z}{(1 - \sqrt{z})^{4(1-\alpha)}}, \quad (z \in \mathbb{U}) \quad (33)$$

with $0 \leq \alpha < 1$, then

$$\begin{aligned} f'(z) &\prec \int_0^z \frac{1}{(1 - \sqrt{t})^{4(1-\alpha)}} dt \\ &= \frac{1}{(2\alpha - 1)(4\alpha - 3)} \left(1 - (1 - \sqrt{z})^{4\alpha-3} (1 - (3 - 4\alpha)\sqrt{z}) \right), \quad (z \in \mathbb{U}). \end{aligned} \quad (34)$$

Next, we have the following theorem.

THEOREM 2. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\frac{zf'(z) - f(z)}{z} \prec \frac{z}{(1 - \sqrt[n]{z})^{2n(1-\alpha)}}, \quad (z \in \mathbb{U}) \quad (35)$$

with $0 \leq \alpha < 1$, then

$$\frac{f(z)}{z} \prec \int_0^z \frac{1}{(1 - \sqrt[n]{t})^{2n(1-\alpha)}} dt, \quad (z \in \mathbb{U}). \quad (36)$$

Proof. Let us consider a function $p(z)$ by $p(z) = \frac{f(z)}{z}$ for $f(z) \in \overline{\mathcal{A}}(n)$. Then $p(z)$ is analytic in \mathbb{U} and $p(0) = 1$. Since

$$zp'(z) = \frac{zf'(z) - f(z)}{z} \prec \frac{z}{(1 - \sqrt[n]{z})^{2n(1-\alpha)}}, \quad (z \in \mathbb{U}), \quad (37)$$

Lemma 1 implies that

$$p(z) = \frac{f(z)}{z} \prec \int_0^z \frac{1}{(1 - \sqrt[n]{t})^{2n(1-\alpha)}} dt, \quad (z \in \mathbb{U}). \quad \square \quad (38)$$

Next, we have to introduce the following lemma by Hallenbeck and Ruscheweyh [5].

LEMMA 2. Let a function $p(z)$ be analytic in \mathbb{U} with $p(0) = 1$ and $h(z)$ be analytic and convex in \mathbb{U} . If $p(z)$ satisfies

$$p(z) + zp'(z) \prec h(z), \quad (z \in \mathbb{U}), \quad (39)$$

then

$$p(z) \prec \frac{1}{z} \int_0^z h(t) dt, \quad (z \in \mathbb{U}). \quad (40)$$

Applying Lemma 2, we derive the following theorem.

THEOREM 3. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$f'(z) + zf''(z) \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, \quad (z \in \mathbb{U}) \quad (41)$$

for some real α ($0 \leq \alpha < 1$), then

$$f'(z) \prec \frac{1}{z} \int_0^z \frac{1 + (1 - 2\alpha)\sqrt[n]{t}}{1 - \sqrt[n]{t}} dt, \quad (z \in \mathbb{U}). \quad (42)$$

Proof. We consider a function $p(z) = f'(z)$ for $f(z) \in \overline{\mathcal{A}}(n)$. Then $p(z)$ is analytic in \mathbb{U} with $p(0) = 1$. Note that the function $h(z)$ given by

$$h(z) = \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, \quad (z \in \mathbb{U}) \quad (43)$$

is convex in \mathbb{U} . By the condition (41), we say that

$$p(z) + zp'(z) = f'(z) + zf''(z) \prec h(z) = \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}. \quad (44)$$

Using Lemma 2, we see that

$$p(z) = f'(z) \prec \frac{1}{z} \int_0^z h(t) dt = \frac{1}{z} \int_0^z \frac{1 + (1 - 2\alpha)\sqrt[n]{t}}{1 - \sqrt[n]{t}} dt, \quad (z \in \mathbb{U}). \quad \square \quad (45)$$

Taking $n = 2$ in Theorem 3, we obtain the following corollary.

COROLLARY 2. If $f(z) \in \overline{\mathcal{A}}(2)$ satisfies

$$f'(z) + zf''(z) \prec \frac{1 + (1 - 2\alpha)\sqrt{z}}{1 - \sqrt{z}}, \quad (z \in \mathbb{U}) \quad (46)$$

with $0 \leq \alpha < 1$, then

$$f'(z) \prec (2\alpha - 1) - 4(1 - \alpha) \frac{\log(1 - \sqrt{z})}{z} + \frac{4(1 - \alpha)}{\sqrt{z}}. \quad (47)$$

Considering $p(z) = \frac{f(z)}{z}$ for $f(z) \in \overline{\mathcal{A}}(n)$ and

$$h(z) = \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (48)$$

with $0 \leq \alpha < 1$ in Lemma 2, we have the following theorem.

THEOREM 4. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$f'(z) \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}}, \quad (z \in \mathbb{U}) \quad (49)$$

with $0 \leq \alpha < 1$, then

$$f(z) \prec \int_0^z \frac{1 + (1 - 2\alpha)\sqrt[n]{t}}{1 - \sqrt[n]{t}} dt, \quad (z \in \mathbb{U}). \quad (50)$$

REMARK 1. If we consider $f(z) \in \overline{\mathcal{A}}(2)$ given by

$$f(z) = (2\alpha - 1)z - 4(1 - \alpha)\log(1 - \sqrt{z}) + 4(1 - \alpha)\sqrt{z}. \quad (51)$$

with $0 \leq \alpha < 1$, then

$$f'(z) = \frac{1 + (1 - 2\alpha)\sqrt{z}}{1 - \sqrt{z}}, \quad (z \in \mathbb{U}), \quad (52)$$

that is $f(z) \in \overline{\mathcal{R}}(2, \alpha)$.

Next, we introduce the lemma due to Miller and Mocanu [7].

LEMMA 3. Let $F(z)$ be analytic in \mathbb{U} and $G(z)$ be analytic in \mathbb{U} and the boundary of \mathbb{U} with $F(0) = G(0)$. If $F(z)$ is not subordinate to $G(z)$, then there exist points $z_0 \in \mathbb{U}$ and $\xi_0 \in \partial\mathbb{U}$, and a real $m \geq 1$ for which $F(|z| < |z_0|) \subset G(\mathbb{U})$,

$$\begin{aligned} (i) \quad & F(z_0) = G(\xi_0) \\ (ii) \quad & z_0 F'(z_0) = m \xi_0 G'(\xi_0). \end{aligned}$$

Using the above lemma, we prove the following theorem.

THEOREM 5. Let β_0 be the solution of

$$\beta\pi = \frac{3}{2}\pi - \tan^{-1}\left(\frac{\beta}{n}\right) \quad (53)$$

and let

$$\alpha = \beta + \frac{2}{\pi} \tan^{-1}\left(\frac{\beta}{n}\right), \quad (0 < \beta \leq \beta_0) \quad (54)$$

with $n = 1, 2, 3, \dots$. If $p(z)$ is analytic in \mathbb{U} with $p(0) = 1$ and

$$p(z) + zp'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}}\right)^\alpha \quad (z \in \mathbb{U}) \quad (55)$$

then

$$p(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}}\right)^\beta \quad (z \in \mathbb{U}). \quad (56)$$

Proof. We use the same method of the proof by Miller and Mocanu (Theorem 5 in [8]). Note that (53) implies

$$\frac{2}{\pi} \tan^{-1} \left(\frac{\beta}{n} \right) = 3 - 2\beta \quad (57)$$

and

$$\beta \leq 3 - \alpha \quad (0 < \beta \leq \beta_0) \quad (58)$$

by (54). We consider the functions

$$h(z) = \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\alpha \quad (59)$$

and

$$g(z) = \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta. \quad (60)$$

Then we see that $|\arg h(z)| < \frac{\pi}{2}\alpha$ and $|\arg g(z)| < \frac{\pi}{2}\beta$ in \mathbb{U} . For such $h(z)$ and $g(z)$, we need to show that $p(z) \prec g(z)$ ($z \in \mathbb{U}$). Since $p(0) = g(0) = 1$, by Lemma 3, if $p(z)$ is not subordinate to $g(z)$, then there exist points $z_0 \in \mathbb{U}$ and $\xi_0 \in \partial\mathbb{U}$, and a real $m \geq 1$ for which $p(|z| < |z_0|) \subset g(\mathbb{U})$, $p(z_0) = g(\xi_0)$ and $z_0 p'(z_0) = m \xi_0 g'(\xi_0)$. Let $p(z_0) \neq 0$, then $g(\xi_0) \neq 0$ and $\sqrt[n]{\xi_0} \neq \pm 1$. Noting that

$$|\arg g(\xi_0)| = \beta \left| \arg \left(\frac{1 + \sqrt[n]{\xi_0}}{1 - \sqrt[n]{\xi_0}} \right) \right| \leq \frac{\pi}{2} \beta \quad (\xi_0 \in \partial\mathbb{U}), \quad (61)$$

we consider

$$ir = \frac{1 + \sqrt[n]{\xi_0}}{1 - \sqrt[n]{\xi_0}} \quad (\xi_0 \in \partial\mathbb{U}). \quad (62)$$

Then, we obtain that

$$\begin{aligned} p(z_0) + z_0 p'(z_0) &= g(\xi_0) + m \xi_0 g'(\xi_0) \\ &= g(\xi_0) \left(1 + m \frac{\xi_0 g'(\xi_0)}{g(\xi_0)} \right) \\ &= (ir)^\beta \left(1 + \frac{2m\beta}{n} \frac{\sqrt[n]{\xi_0}}{1 - \left(\sqrt[n]{\xi_0} \right)^2} \right) \\ &= (ir)^\beta \left(1 + i \frac{m\beta}{2n} \left(r + \frac{1}{r} \right) \right). \end{aligned} \quad (63)$$

It follows from (63), that

$$\arg(p(z_0) + z_0 p'(z_0)) = \frac{\pi}{2} \beta + \tan^{-1} \left(\frac{m\beta \left(r + \frac{1}{r} \right)}{2n} \right) \quad (64)$$

and

$$\frac{\pi}{2}\beta + \tan^{-1}\left(\frac{\beta}{n}\right) \leq |\arg(p(z_0) + z_0 p'(z_0))| \leq \frac{\pi}{2}\beta + \frac{\pi}{2}. \quad (65)$$

Thus, using (57) and (58), we have

$$\frac{\pi}{2}\alpha \leq |\arg(p(z_0) + z_0 p'(z_0))| \leq 2\pi - \frac{\pi}{2}\alpha. \quad (66)$$

Since $|\arg h(z)| < \frac{\pi}{2}\alpha$ ($z \in \mathbb{U}$), (66) contradicts (55). Therefore, the subordination (56) is true for $p(z_0) \neq 0$ ($z \in \mathbb{U}$). If $p(z_0) = 0$ ($z \in \mathbb{U}$), with the same reason of the proof by Miller and Mocanu (Theorem 5 in [8]), we have

$$\frac{\pi}{2}\alpha \leq |\arg(p(z_0) + z_0 p'(z_0))| \leq 2\pi - \frac{\pi}{2}\alpha. \quad \square \quad (67)$$

Taking $\beta = n$ in Theorem 5, then we have the following corollary.

COROLLARY 3. *If $p(z)$ is analytic in \mathbb{U} with $p(0) = 1$ and*

$$p(z) + zp'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^{\frac{2n+1}{2}} \quad (z \in \mathbb{U}) \quad (68)$$

for $n = 1, 2, 3, \dots$, then

$$p(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^n \quad (z \in \mathbb{U}). \quad (69)$$

Taking $\beta = \sqrt{3}n$ in Theorem 5, then we obtain the following corollary.

COROLLARY 4. *If $p(z)$ is analytic in \mathbb{U} with $p(0) = 1$ and*

$$p(z) + zp'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^{\frac{2}{3} + \sqrt{3}n} \quad (z \in \mathbb{U}) \quad (70)$$

for $n = 1, 2, 3, \dots$, then

$$p(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^{\sqrt{3}n} \quad (z \in \mathbb{U}). \quad (71)$$

Taking $p(z) = f'(z)$ in Theorem 5, then we have the following corollary.

COROLLARY 5. *Let α and β be define (53) and (54). If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies*

$$f'(z) + zf''(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\alpha \quad (z \in \mathbb{U}) \quad (72)$$

for $n = 1, 2, 3, \dots$, then

$$f'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta \quad (z \in \mathbb{U}), \quad (73)$$

that is, $f(z) \in \overline{\mathcal{H}}(n, 0)$.

If we consider $p(z) = \frac{f(z)}{z}$ for $f(z) \in \overline{\mathcal{A}}(n)$ in Theorem 5, then we have the following corollary.

COROLLARY 6. Let α and β be define (53) and (54). If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$f'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\alpha \quad (z \in \mathbb{U}) \quad (74)$$

for $n = 1, 2, 3, \dots$, then

$$\frac{f(z)}{z} \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta \quad (z \in \mathbb{U}). \quad (75)$$

Next, we show the following theorem.

THEOREM 6. Let $p(z)$ be analytic in \mathbb{U} with $p(0) = 1$. If $p(z)$ satisfies

$$p(z) + zp'(z) \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}) \quad (76)$$

for some real α ($0 \leq \alpha < 1$), then

$$p(z) \prec \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}) \quad (77)$$

where $n = 1, 2, 3, \dots$.

Proof. If we suppose that $p(z)$ is not subordinate to a function $g(z)$ given by

$$g(z) = \frac{1 + (1 - 2\alpha)\sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}), \quad (78)$$

Lemma 3 gives us that there exist points $z_0 \in \mathbb{U}$ and $\xi_0 \in \partial\mathbb{U}$, and a real $m \geq 1$ for which $p(|z| < |z_0|) \subset g(\mathbb{U})$, $p(z_0) = g(\xi_0)$ and $z_0 p'(z_0) = m \xi_0 g'(\xi_0)$. It follows that

$$\begin{aligned} p(z_0) + z_0 p'(z_0) &= g(\xi_0) + m \xi_0 g'(\xi_0) \\ &= \frac{1 + (1 - 2\alpha)\sqrt[n]{\xi_0}}{1 - \sqrt[n]{\xi_0}} \left\{ 1 + \frac{m}{n} \left(\frac{(1 - 2\alpha)\sqrt[n]{\xi_0}}{1 + (1 - 2\alpha)\sqrt[n]{\xi_0}} + \frac{\sqrt[n]{\xi_0}}{1 - \sqrt[n]{\xi_0}} \right) \right\}. \end{aligned} \quad (79)$$

Letting $\xi_0 = e^{i\theta}$ ($0 \leq \theta < 2\pi$), we have

$$\begin{aligned} p(z_0) + z_0 p'(z_0) &= \frac{1 + (1 - 2\alpha)e^{i\frac{\theta}{n}}}{1 - e^{i\frac{\theta}{n}}} \\ &\quad + \frac{m}{n} \left(\frac{(1 - 2\alpha)e^{i\frac{\theta}{n}}}{1 - e^{i\frac{\theta}{n}}} + \frac{e^{i\frac{\theta}{n}} \left(1 + (1 - 2\alpha)e^{i\frac{\theta}{n}} \right)}{\left(1 - e^{i\frac{\theta}{n}} \right)^2} \right). \end{aligned} \quad (80)$$

This implies that

$$Re(p(z_0) + z_0 p'(z_0)) = \alpha - \frac{m}{2n} \left(1 - 2\alpha + \frac{1 + (1 - 2\alpha) \cos \frac{\theta}{n}}{1 - \cos \frac{\theta}{n}} \right). \quad (81)$$

If we define a function $k(t)$ given by

$$k(t) = \frac{1 + (1 - 2\alpha)t}{1 - t} \quad \left(t = \cos \frac{\theta}{n} \right), \quad (82)$$

then $k(t) > \alpha$ and

$$k'(t) = \frac{2(1 - \alpha)}{(1 - t)^2} > 0. \quad (83)$$

Thus, we have

$$Re(p(z_0) + z_0 p'(z_0)) < \alpha - \frac{m(1 - \alpha)}{2n} < \alpha. \quad (84)$$

Noting that $Reg(z) > \alpha$ ($z \in \mathbb{U}$), (84) contradicts the condition (76). Therefore, $p(z)$ satisfies the subordination (77). \square

Taking $p(z) = f'(z)$ in Theorem 6, then we have the following corollary.

COROLLARY 7. *If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies*

$$f'(z) + z f''(z) \prec \frac{1 + (1 - 2\alpha) \sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}) \quad (85)$$

for some real α ($0 \leq \alpha < 1$), then

$$f'(z) \prec \frac{1 + (1 - 2\alpha) \sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}), \quad (86)$$

that is, $f(z) \in \overline{\mathcal{R}}(n, \alpha)$.

Further, we have the following corollary.

COROLLARY 8. *If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies*

$$f'(z) \prec \frac{1 + (1 - 2\alpha) \sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}) \quad (87)$$

for some real α ($0 \leq \alpha < 1$), then

$$\frac{f(z)}{z} \prec \frac{1 + (1 - 2\alpha) \sqrt[n]{z}}{1 - \sqrt[n]{z}} \quad (z \in \mathbb{U}). \quad (88)$$

To discuss the next our results for subordinations, we have to introduce the lemma due to Nunokawa, Owa and Sokol [9].

LEMMA 4. Let $p(z)$ be analytic in \mathbb{U} with $p(0) = 1$ and $p(z) \neq 0$ in \mathbb{U} . If there exists a point $z_0 \in \mathbb{U}$ such that

$$|\arg p(z)| < \frac{\pi}{2}\beta \quad (|z| < |z_0| < 1) \quad (89)$$

and

$$|\arg p(z_0)| = \frac{\pi}{2}\beta \quad (90)$$

for some $\beta > 0$, then

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik\beta, \quad (91)$$

where

$$k \geq \frac{a^2 + 1}{2a} \geq 1 \quad (\arg p(z_0) = \frac{\pi}{2}\beta) \quad (92)$$

and

$$k \leq -\frac{a^2 + 1}{2a} \leq -1 \quad (\arg p(z_0) = -\frac{\pi}{2}\beta), \quad (93)$$

where

$$(p(z_0))^{\frac{1}{\beta}} = \pm ia \quad (a > 0). \quad (94)$$

Now, we derive the following theorem.

THEOREM 7. Let $p(z)$ be analytic with $p(0) = 1$ and $p(z) \neq 0$ in \mathbb{U} . If $p(z)$ satisfies

$$p(z) + \frac{zp'(z)}{p(z)} \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta + \frac{2\beta \sqrt[n]{z}}{1 - (\sqrt[n]{z})^2} \quad (z \in \mathbb{U}) \quad (95)$$

with $0 < \beta \leq 1$ and $n = 1, 2, 3, \dots$, then

$$p(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta \quad (z \in \mathbb{U}). \quad (96)$$

Proof. We consider a function $p(z)$ which is not satisfy the subordination (96). Then there exists a point $z_0 \in \mathbb{U}$ such that

$$|\arg p(z)| < \frac{\pi}{2}\beta \quad (|z| < |z_0| < 1) \quad (97)$$

and

$$|\arg p(z_0)| = \frac{\pi}{2}\beta. \quad (98)$$

Applying Lemma 4, we have

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik\beta, \quad (99)$$

where k satisfies (92) and (93) in Lemma 4. If $p(z_0) = \frac{\pi}{2}\beta$, then

$$p(z_0) + \frac{z_0 p'(z_0)}{p(z_0)} = (ia)^\beta + ik\beta \quad (a > 0). \quad (100)$$

Considering a boundary point $z = e^{i\theta}$ in \mathbb{U} , we see that

$$\left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta + \frac{2\beta \sqrt[n]{z}}{1 - (\sqrt[n]{z})^2} = \left(i \frac{\sin \frac{\theta}{n}}{1 - \cos \frac{\theta}{n}} \right)^\beta + i \frac{\beta}{\sin \frac{\theta}{n}}. \quad (101)$$

Thus, if the subordination (95) is satisfied, then

$$a = \frac{\sin \frac{\theta}{n}}{1 - \cos \frac{\theta}{n}} > 0 \quad (102)$$

and

$$k \geq \frac{a^2 + 1}{2a} = \frac{1}{\sin \frac{\theta}{n}} \geq 1. \quad (103)$$

This implies that $p(z)$ is not satisfy the condition (95). For the case $\arg p(z_0) = -\frac{\pi}{2}\beta$, using the same way for the case $\arg p(z_0) = \frac{\pi}{2}\beta$, we say that $p(z)$ is not satisfy the condition (95). Therefore, we complete the proof of the theorem. \square

Letting $p(z) = f'(z)$ in Theorem 7, we have the following corollary.

COROLLARY 9. *If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies $f'(z) \neq 0$ ($z \in \mathbb{U}$) and*

$$f'(z) + \frac{zf''(z)}{f'(z)} \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta + \frac{2\beta \sqrt[n]{z}}{1 - (\sqrt[n]{z})^2} \quad (z \in \mathbb{U}) \quad (104)$$

with $0 < \beta \leq 1$ then

$$f'(z) \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta \quad (z \in \mathbb{U}), \quad (105)$$

that is, $f(z) \in \overline{\mathcal{R}}(n, 0)$.

COROLLARY 10. *If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies $\frac{zf'(z)}{f(z)} \neq 0$ ($z \in \mathbb{U}$) and*

$$1 + \frac{zf''(z)}{f'(z)} \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta + \frac{2\beta \sqrt[n]{z}}{1 - (\sqrt[n]{z})^2} \quad (z \in \mathbb{U}) \quad (106)$$

with $0 < \beta \leq 1$ then

$$\frac{zf'(z)}{f(z)} \prec \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right)^\beta \quad (z \in \mathbb{U}), \quad (107)$$

that is, $f(z) \in \overline{\mathcal{S}}^*(n, 0)$.

3. Consideration for strongly functions

Let us consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f(z) = \int_0^z \left(\frac{1 + \sqrt[n]{t}}{1 - \sqrt[n]{t}} \right)^\alpha dt, (z \in \mathbb{U}). \quad (108)$$

with $0 \leq \alpha < 1$. Then $f(z)$ satisfies

$$|\arg f'(z)| = \alpha \left| \arg \left(\frac{1 + \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right) \right| < \frac{\pi}{2} \alpha, (z \in \mathbb{U}). \quad (109)$$

We say that $f(z)$ is strongly of order α in \mathbb{U} if $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$|\arg f'(z)| < \frac{\pi}{2} \alpha, (z \in \mathbb{U}) \quad (110)$$

for $0 \leq \alpha < 1$.

To discuss about strongly functions in \mathbb{U} we need the following lemma due to Fejér and Riesz [1] (also by Tsuji [11].)

LEMMA 5. *Let a function $f(z)$ be analytic in $|z| \leq 1$. Then, $f(z)$ satisfies*

$$\int_{-1}^1 |f(z)|^\rho |dz| \leq \frac{1}{2} \int_{|z|=1} |f(z)|^\rho |dz|, (\rho > 0), \quad (111)$$

where the above integral on the left hand side is considered along the real axis.

REMARK 2. If we make a change of variables in Lemma 5, then the inequality (111) becomes

$$\int_{-r}^r |f(\rho e^{i\theta})|^\rho d\rho \leq \frac{r}{2} \int_0^{2\pi} |f(re^{i\theta})|^\rho d\theta. \quad (112)$$

Also, we use the following lemma by Gwynne [4].

LEMMA 6. *Let $f(z)$ be a complex valued harmonic function defined on a neighborhood of a closed disk of radius 1 and center 0 in the complex plane. Then*

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\rho}) \frac{1 - r^2}{1 + r^2 - 2r\cos(\theta - \rho)} d\rho \quad (113)$$

and

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{1 - r^2}{1 + r^2 - 2r\cos\rho} d\rho = 1. \quad (114)$$

Now, we derive the following theorem.

THEOREM 8. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{\alpha}{2}, \quad (z \in \mathbb{U}) \quad (115)$$

for some real α ($0 < \alpha \leq 1$), then

$$|\arg f'(z)| \leq \frac{\pi}{2} \alpha, \quad (z \in \mathbb{U}). \quad (116)$$

Proof. We note that

$$\log f'(z) = \log |f'(z)| + i \arg f'(z) \quad (117)$$

and

$$\log f'(z) = \int_0^z (\log f'(t))' dt = \int_0^z \frac{f''(t)}{f'(t)} dt. \quad (118)$$

Thus we have

$$\begin{aligned} |\arg f'(z)| &= |\operatorname{Im}(\log f'(z))| \\ &= \left| \operatorname{Im} \int_0^z \frac{f''(t)}{f'(t)} dt \right| \\ &\leq \left| \operatorname{Im} \int_0^r \frac{f''(\rho e^{i\theta})}{f'(\rho e^{i\theta})} e^{i\theta} d\rho \right| \\ &= \int_0^r \left| \operatorname{Im} \left(\frac{e^{i\theta} f''(\rho e^{i\theta})}{f'(\rho e^{i\theta})} \right) \right| d\rho \\ &\leq \int_{-r}^r \left| \operatorname{Im} \left(\frac{e^{i\theta} f''(\rho e^{i\theta})}{f'(\rho e^{i\theta})} \right) \right| d\rho \\ &\leq \int_{-r}^r \left| \frac{f''(\rho e^{i\theta})}{f'(\rho e^{i\theta})} \right| d\rho, \end{aligned} \quad (119)$$

where $z = re^{i\theta}$ ($0 \leq \theta < 2\pi$), $0 \leq r < 1$ and $0 \leq \rho \leq r$. Using the inequality (112) with $\rho = 1$, we obtain

$$\begin{aligned} |\arg f'(z)| &\leq \frac{r}{2} \int_0^{2\pi} \left| \frac{f''(re^{i\theta})}{f'(re^{i\theta})} \right| d\theta \\ &= \frac{1}{2} \int_0^{2\pi} \left| \frac{re^{i\theta} f''(re^{i\theta})}{f'(re^{i\theta})} \right| d\theta \\ &\leq \frac{\alpha}{4} \int_0^{2\pi} d\theta = \frac{\pi}{2} \alpha. \end{aligned} \quad (120)$$

This completes the proof of theorem. \square

Taking $\alpha = 1$ in Theorem 8, we obtain the following corollary.

COROLLARY 11. Let $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{1}{2}, (z \in \mathbb{U}) \quad (121)$$

then

$$|\arg f'(z)| < \frac{\pi}{2}, (z \in \mathbb{U}). \quad (122)$$

REMARK 3. If we take $n = 3$ in Theorem 8, then we have the result by Güney and Owa [2].

Next, we derive the following theorem.

THEOREM 9. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\left| \frac{zf''(z)}{f'(z)} \right| < \frac{\alpha}{2} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right), (z \in \mathbb{U}) \quad (123)$$

for some real α ($0 < \alpha \leq 1$) and some real β ($\beta \neq -1$), then

$$|\arg f'(z)| < \frac{\pi}{2} \alpha, (z \in \mathbb{U}). \quad (124)$$

Proof. If we use the same method of the proof in Theorem 8, we have that

$$|\arg f'(z)| < \frac{1}{2} \int_0^{2\pi} \left| \frac{re^{i\theta} f''(re^{i\theta})}{f'(re^{i\theta})} \right| d\theta. \quad (125)$$

Thus, using the inequality (123), we know that

$$\begin{aligned} |\arg f'(z)| &< \frac{\alpha}{4} \int_0^{2\pi} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{r} e^{i\frac{\theta}{n}}}{1 - \sqrt[n]{r} e^{i\frac{\theta}{n}}} \right) d\theta \\ &= \frac{\alpha}{4} \int_0^{2\pi} \left(\frac{1 + (\beta - 1) \sqrt[n]{r} \cos \left(\frac{\theta}{n} \right) - \beta (\sqrt[n]{r})^2}{1 - 2 \sqrt[n]{r} \cos \left(\frac{\theta}{n} \right) + (\sqrt[n]{r})^2} \right) d\theta \\ &= \frac{\alpha}{4} \int_0^{2\pi} \left\{ \frac{1 - \beta}{2} + \frac{1 + \beta}{2} \frac{1 - (\sqrt[n]{r})^2}{1 + (\sqrt[n]{r})^2 - 2 \sqrt[n]{r} \cos \left(\frac{\theta}{n} \right)} \right\} d\theta. \end{aligned} \quad (126)$$

Further, we note that

$$\int_0^{2\pi} \frac{1 - (\sqrt[n]{r})^2}{1 + (\sqrt[n]{r})^2 - 2 \sqrt[n]{r} \cos \frac{\theta}{n}} d\theta = n \int_0^{\frac{2\pi}{n}} \frac{1 - (\sqrt[n]{r})^2}{1 + (\sqrt[n]{r})^2 - 2 \sqrt[n]{r} \cos \rho} d\rho \leq 2\pi \quad (127)$$

by Lemma 6. Therefore, we obtain that

$$|\arg f'(z)| < \frac{\pi}{2} \alpha, (z \in \mathbb{U}). \quad \square \quad (128)$$

REMARK 4. If we consider $n = 3$ in Theorem 9, then we obtain the result for $f(z) \in \overline{\mathcal{A}}(3)$ in Güney and Owa [2].

EXAMPLE 2. We consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f'(z) = \left(\frac{2}{2 - \sqrt[n]{z}} \right)^{3\alpha}, \quad (z \in \mathbb{U}) \quad (129)$$

with α ($0 < \alpha \leq 1$). It follows that

$$|\arg f'(z)| = 3\alpha |\arg(2 - \sqrt[n]{z})| < 3\alpha \frac{\pi}{6} = \frac{\pi}{2}\alpha, \quad (z \in \mathbb{U}) \quad (130)$$

and

$$\left| \frac{zf''(z)}{f'(z)} \right| = \frac{3\alpha}{n} \left| \frac{\sqrt[n]{z}}{2 - \sqrt[n]{z}} \right| < \frac{3}{n}\alpha, \quad (z \in \mathbb{U}). \quad (131)$$

If we consider some real β such that $\beta \leq \frac{n-12}{n}$, then we have

$$\left| \frac{zf''(z)}{f'(z)} \right| < \frac{3}{n}\alpha \leq \frac{\alpha(1-\beta)}{4} < \frac{\alpha}{2} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right), \quad (z \in \mathbb{U}). \quad (132)$$

Next, we derive the following theorem.

THEOREM 10. If $f(z) \in \overline{\mathcal{A}}(n)$ satisfies

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{\alpha}{2} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right), \quad (z \in \mathbb{U}) \quad (133)$$

for some real α ($0 < \alpha \leq 1$) and some real β ($\beta \neq -1$), then

$$\left| \arg \frac{f(z)}{z} \right| < \frac{\pi}{2}\alpha, \quad (z \in \mathbb{U}). \quad (134)$$

Proof. We note that

$$\log \left(\frac{f(z)}{z} \right) = \log \left| \frac{f(z)}{z} \right| + i \arg \left(\frac{f(z)}{z} \right) \quad (135)$$

and

$$\begin{aligned} \log \left(\frac{f(z)}{z} \right) &= \int_0^z \left(\log \left(\frac{f(t)}{t} \right) \right)' dt \\ &= \int_0^z \left(\frac{f'(t)}{f(t)} - \frac{1}{t} \right) dt. \end{aligned} \quad (136)$$

Thus, we have

$$\begin{aligned}
 \left| \arg \left(\frac{f(z)}{z} \right) \right| &= \left| \operatorname{Im} \left(\log \left(\frac{f(z)}{z} \right) \right) \right| \\
 &= \left| \operatorname{Im} \int_0^z \left(\frac{f'(t)}{f(t)} - \frac{1}{t} \right) dt \right| \\
 &= \left| \operatorname{Im} \int_0^r \left(\frac{f'(\rho e^{i\theta})}{f(\rho e^{i\theta})} - \frac{1}{\rho e^{i\theta}} \right) e^{i\theta} d\rho \right| \\
 &\leq \int_0^r \left| \operatorname{Im} \left(\frac{f'(\rho e^{i\theta})}{f(\rho e^{i\theta})} - \frac{1}{\rho e^{i\theta}} \right) e^{i\theta} \right| d\rho \\
 &< \int_{-r}^r \left| \operatorname{Im} \left(\frac{e^{i\theta} f'(\rho e^{i\theta})}{f(\rho e^{i\theta})} - \frac{1}{\rho} \right) \right| d\rho \\
 &\leq \frac{1}{2} \int_0^{2\pi} \left| \frac{re^{i\theta} f'(re^{i\theta})}{f(re^{i\theta})} - 1 \right| d\theta \\
 &< \frac{\alpha}{4} \int_0^{2\pi} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{r} e^{i\frac{\theta}{n}}}{1 - \sqrt[n]{r} e^{i\frac{\theta}{n}}} \right) d\theta \\
 &\leq \frac{\pi}{2} \alpha.
 \end{aligned} \tag{137}$$

Thus, we complete the proof of the theorem. \square

EXAMPLE 3. We consider a function $f(z) \in \overline{\mathcal{A}}(n)$ given by

$$f(z) = z \left(\frac{2}{2 - \sqrt[n]{z}} \right)^{3\alpha}, \quad (z \in \mathbb{U}) \tag{138}$$

with α ($0 < \alpha \leq 1$). Then

$$w(z) = \frac{2}{2 - \sqrt[n]{z}} \tag{139}$$

satisfies

$$\left| w(z) - \frac{4}{3} \right| < \frac{2}{3}, \quad (z \in \mathbb{U}). \tag{140}$$

Thus $w(z)$ satisfies

$$|\arg w(z)| = \left| \arg \left(\frac{2}{2 - \sqrt[n]{z}} \right) \right| < \frac{\pi}{6}, \quad (z \in \mathbb{U}). \tag{141}$$

This gives us that

$$\left| \arg \left(\frac{f(z)}{z} \right) \right| = 3\alpha \left| \arg \left(\frac{2}{2 - \sqrt[n]{z}} \right) \right| < \frac{\pi}{2} \alpha, \quad (z \in \mathbb{U}). \tag{142}$$

Also, we have

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| = \frac{3\alpha}{n} \left| \frac{\sqrt[n]{z}}{2 - \sqrt[n]{z}} \right| < \frac{3}{n}\alpha, (z \in \mathbb{U}). \quad (143)$$

Therefore, considering β such that $\beta \leq \frac{n-12}{n}$, we see

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \frac{3}{n}\alpha \leq \frac{\alpha(1-\beta)}{4} < \frac{\alpha}{2} \operatorname{Re} \left(\frac{1 + \beta \sqrt[n]{z}}{1 - \sqrt[n]{z}} \right), (z \in \mathbb{U}). \quad (144)$$

Author contributions. Conceptualization, H. Ö. G., D. B. and S. O.; Investigation, H. Ö. G., D. B. and S. O.; Methodology, H. Ö. G., D. B. and S. O.; Writing—original draft, S. O.; Writing—review and editing, H. Ö. G. and D. B. All authors have read and agreed to the published version of the manuscript.

Funding. This research received no external funding.

Competing interests. The authors state that they have no conflicts of interest.

Data Availability Statement. No data were used to support this study.

REFERENCES

- [1] L. FEJÉR, F. RIESZ, *Über einige funktionentheoretische Ungleichungen*, Mathematische Zeitschrift **11** (3–4) (1921) 305–314.
- [2] H. Ö. GÜNEY, S. OWA, *New properties of analytic functions*, Mathematics **12**, (2024) 1469.
- [3] H. Ö. GÜNEY, D. BREAZ, S. OWA, *On some interesting classes of analytic functions related to univalent functions*, Mathematics **12**, (2024) 513.
- [4] E. GWYNME, *The Poisson Integral Formula and Representations of $SU(1,1)$* , Rose-Hulman Undergraduate Mathematics Journal **12.2** (2011) 1.
- [5] D. J. HALLENBECK, S. RUSCHEWEYH, *Subordination by convex functions*, Proceedings of the American Mathematical Society **52.1** (1975) 191–195.
- [6] S. S. MILLER, P. T. MOCANU, *Differential Subordinations: Theory and Applications*, Marcel Dekker Inc., New York, (2000).
- [7] S. S. MILLER, P. T. MOCANU, *Differential subordinations and univalent functions*, Michigan Mathematical Journal **28** (2), (1981) 157–171.
- [8] S. S. MILLER, P. T. MOCANU, *Marx-Strohhäcker differential subordination systems*, Proceedings of the American Mathematical Society **99** (3), (1987) 527–534.
- [9] M. NUNOKAWA, S. OWA, J. SOKÓL, *A criterion for bounded functions*, Bulletin of the Korean Mathematical Society **53** (1), (2016) 215–225.

- [10] T. J. SUFFRIDGE, *Some remarks on convex maps of the unit disk*, Duke Mathematical Journal **37**, (1970) 775–777.
- [11] M. TSUJI, *Complex Function Theory*, Tokyo, Japanese, Maki Book Comp., 1968.

(Received October 8, 2024)

Hatun Özlem Güney
Department of Mathematics
Faculty of Science, Dicle University
21280 Diyarbakır, Türkiye
e-mail: ozlemg@dicle.edu.tr

Daniel Breaz
Department of Exact Sciences and Engineering
“1 Decembrie 1918” University
Alba Iulia, 510009 Alba Iulia, Romania
e-mail: dbreaz@uab.ro

Shigeyoshi Owa
Honorary Professor 1 Decembrie 1918 University
Alba Iulia, Romania
e-mail: shige21@ican.zaq.ne.jp