

## REMARKS ON EXTREMAL FUNCTIONS FOR THE ANISOTROPIC TRUDINGER–MOSER INEQUALITIES INVOLVING $L^p$ NORM

XIANFENG SU, XIAOMENG LI\*, RULONG XIE AND MENG QU

**Abstract.** Let  $W^{1,n}(\mathbb{R}^n)$  ( $n \geq 2$ ) be the standard Sobolev space, and denote, for  $p > n$

$$\gamma = \inf_{u \in W^{1,n}(\mathbb{R}^n), u \neq 0} \frac{\int_{\mathbb{R}^n} (F^n(\nabla u) + |u|^n) dx}{\left( \int_{\mathbb{R}^n} |u|^p dx \right)^{\frac{n}{p}}},$$

where  $F : \mathbb{R}^n \rightarrow [0, \infty)$  be a convex function of class  $C^2(\mathbb{R}^n \setminus \{0\})$ , which is even and positively homogeneous of degree 1. For  $\gamma \in [0, \gamma_1)$ , we define a norm in  $W^{1,n}(\mathbb{R}^n)$  by

$$\|u\|_{F,n,\gamma,p} = \left( \int_{\mathbb{R}^n} (F^n(\nabla u) + |u|^n) dx - \gamma \left( \int_{\mathbb{R}^n} |u|^p dx \right)^{\frac{n}{p}} \right)^{\frac{1}{n}}.$$

By performing a blow-up analysis, we prove that for real numbers  $0 \leq \gamma < \gamma_1$  and  $p > n$ , the following anisotropic Trudinger–Moser inequality

$$\sup_{u \in W^{1,n}(\mathbb{R}^n), \|u\|_{F,n,\gamma,p} \leq 1} \int_{\mathbb{R}^n} \Phi(\lambda_n |u|^{\frac{n}{n-1}}) dx$$

can be attained by some function  $u_0 \in W^{1,n}(\mathbb{R}^n)$  with  $\|u_0\|_{F,n,\gamma,p} = 1$ , where  $\Phi(t) = e^t - \sum_{j=0}^{n-1} \frac{t^j}{j!}$ ,  $\lambda_n = n^{\frac{n}{n-1}} \kappa_n^{\frac{1}{n-1}}$  and  $\kappa_n$  is the volume of the unit Wulff ball. In the case  $\gamma = 0$ , this is reduced to a result of Zhou–Zhou [19].

*Mathematics subject classification (2020):* 46E35.

*Keywords and phrases:* Extremal function, Trudinger–Moser inequality, blow-up analysis.

## REFERENCES

- [1] A. ALVINO, V. FERONE, G. TROMBETTI, P. L. LIONS, *Convex symmetrization and applications*, Ann. Inst. H. Poincaré Anal. Non Linéaire **14**, 2 (1997), 275–293.
- [2] L. CARLESON, A. CHANG, *On the existence of an extremal function for an inequality of J. Moser*, Bull. Sci. Math., **110**, 2 (1986), 113–127.
- [3] M. FLUCHER, *Extremal functions for Trudinger–Moser inequality in 2 dimensions*, Comment. Math. Helv. **67**, 1 (1992), 471–497.
- [4] X. LI, Y. YANG, *Extremal functions for singular Trudinger–Moser inequalities in the entire Euclidean space*, Journal of Differential Equations, **264**, 8 (2018), 4901–4943.
- [5] Y. LI, *Moser–Trudinger inequality on compact Riemannian manifolds of dimension two*, J. Partial Differential Equations, **14**, 2 (2001), 163–192.
- [6] K. LIN, *Extremal functions for Moser’s inequality*, Trans. Amer. Math. Soc. **348**, (1996), 2663–2671.
- [7] Y. LIU, *An improved Trudinger–Moser inequality involving  $N$ -Finsler–Laplacian and  $L^p$  norm*, Potential Anal. **60**, 2 (2024), 673–701.
- [8] J. MOSER, *A sharp form of an inequality by N. Trudinger*, Indiana. Univ. Math. J., **20**, 11 (1971), 1077–1091.
- [9] J. PEETRE, *Espaces d’interpolation et theoreme de Soboleff*, Ann. Inst. Fourier (Grenoble), **16**, (1966), 279–317.

- [10] S. POHOZAEV, *The Sobolev embedding in the special case  $pl = n$* , Proceedings of the technical scientific conference on advances of scientific research 1964–1965, Mathematics sections, 158–170, Moscov. Energet. Inst., Moscow, 1965.
- [11] N. S. TRUDINGER, *On embeddings into Orlicz space and some applications*, J. Math. Mech., **17**, (1967), 473–483.
- [12] J. SERRIN, *Local behavior of solutions of quasilinear equations*, Acta Math., **111**, (1964), 247–302.
- [13] P. TOLKSDORF, *Regularity for a more general class of quasilinear elliptic equations*, J. Differential Equations, **51**, (1984), 126–150.
- [14] G. F. WANG, C. XIA, *Blow-up analysis of a Finsler-Liouville equation in two dimensions*, J. Differential Equations, **252**, 2 (2012), 1668–1700.
- [15] Y. YANG, *A sharp form of Moser-Trudinger inequality in high dimension*, J. Funct. Anal., **239**, 1 (2006), 100–126.
- [16] Y. YANG, *Existence of positive solutions to quasi-linear elliptic equations with exponential growth in the whole Euclidean space*, J. Funct. Anal., **262**, 4 (2012), 1679–1704.
- [17] Y. YANG, *Extremal functions for Trudinger-Moser inequalities of Adimurthi-Druet type in dimension two*, J. Differential Equations, **258**, 9 (2015), 3161–3193.
- [18] V. I. YUDOVICH, *Some estimates connected with integral operators and with solutions of elliptic equations*, Sov. Math. Doct., **2**, 4 (1961), 746–749.
- [19] C. L. ZHOU, C. Q. ZHOU, *Moser-Trudinger inequality involving the anisotropic Dirichlet norm  $(\int_{\Omega} F^N(\nabla u) dx)^{\frac{1}{N}}$  on  $W_0^{1,N}(\Omega)$* , J. Funct. Anal., **276**, (2019), 2901–2935.
- [20] C. L. ZHOU, C. Q. ZHOU, *On the anisotropic Moser-Trudinger inequality for unbounded domains in  $\mathbb{R}^n$* , Discrete and Continuous Dynamical Systems, **40**, 2 (2020), 847–881.