

EXTREMAL MARGINAL TRACIAL STATES IN COUPLED SYSTEMS

GEOFFREY L. PRICE * AND SHÔICHIRO SAKAI

Abstract. Let Γ be the convex set consisting of all states ϕ on the tensor product $B \otimes B$ of the algebra $B = M_n(\mathbb{C})$ of all $n \times n$ matrices over the complex numbers \mathbb{C} with the property that the restrictions $\phi|_{B \otimes I}$ and $\phi|_{I \otimes B}$ are the unique tracial states on $B \otimes I$ and $I \otimes B$. We find necessary and sufficient conditions for such a state, called a marginal tracial state, to be extremal in Γ . We also give a characterization of those extreme points in Γ which are pure states. We conjecture that all extremal marginal tracial states are pure states.

Mathematics subject classification (2000): 46L30, 46L06.

Key words and phrases: Pure state, trace, marginal tracial state, Schmidt decomposition.

REFERENCES

- [EK] A. EKERT AND P. L. KNIGHT, *Entangled quantum systems and the Schmidt decomposition*, Am. J. Phys. **63** (1995), 415–423.
- [P1] K. R. PARTHASARATHY, *Extremal quantum states in coupled systems*, Ann. Inst. H. Poincaré **41** (2005), 257–268.
- [P2] K. R. PARTHASARATHY, *On extremal correlations*, J. Stat. Planning and Inf. **103** (2002), 73–80.
- [Sa] S. SAKAI, *C^* -algebras and W^* -algebras*, Springer-Verlag, 1971.
- [Sc] E. SCHMIDT, *Zur theorie der linearen und nichtlinearen integralgleichungen*, Math. Annalen **63** (1906), 433–476.
- [Sz] B. SZ.-NAGY, *Spektraldarstellung linearer transformationen des Hilbertschen raumes*, Springer-Verlag, 1942.
- [T] J. TOMIYAMA, *Applications of Fubini type theorem to the tensor products of C^* -algebras*, Tôhoku Math. J. (2) **19** (1967), 340–344.
- [VW] K. G. H. VOLLBRECHT AND R. F. WERNER, *Why two qubits are special*, J. Math. Phys. **41** (2000), 6772–6782.