

## THE NEW $v$ -METRIC INDUCES THE CLASSICAL GAP TOPOLOGY

AMOL SASANE

(Communicated by J. William Helton)

*Abstract.* Let  $\mathcal{A}_+$  denote the set of Laplace transforms of complex Borel measures  $\mu$  on  $[0, +\infty)$  such that  $\mu$  does not have a singular non-atomic part. In [1], an extension of the classical  $v$ -metric of Vinnicombe was given, which allowed one to address robust stabilization problems for unstable plants over  $\mathcal{A}_+$ . In this article, we show that this new  $v$ -metric gives a topology on unstable plants which coincides with the classical gap topology for unstable plants over  $\mathcal{A}_+$  with a single input and a single output.

### 1. Introduction

We recall the general *stabilization problem* in control theory. Suppose that  $R$  is a commutative integral domain with identity (thought of as the class of stable transfer functions) and let  $\mathbb{F}(R)$  denote the field of fractions of  $R$ . Then the stabilization problem is:

$p \in \mathbb{F}(R)$  (an unstable plant transfer function),  
find  $c \in \mathbb{F}(R)$  (a stabilizing controller transfer function),  
such that (the closed loop transfer function)

$$H(p, c) := \begin{bmatrix} p \\ 1 \end{bmatrix} (1 - cp)^{-1} \begin{bmatrix} -c & 1 \end{bmatrix}$$

belongs to  $R^{2 \times 2}$  (that is, it is stable).

In the *robust stabilization problem*, one goes a step further. One knows that the plant is just an approximation of reality, and so one would really like the controller  $c$  to not only stabilize the *nominal* plant  $p$ , but also all sufficiently close plants  $p'$  to  $p$ . The question of what one means by “closeness” of plants thus arises naturally. So one needs a function  $d$  defined on pairs of stabilizable plants such that

1.  $d$  is a metric on the set of all stabilizable plants,
2.  $d$  is amenable to computation, and
3. stabilizability is a robust property of the plant with respect to  $d$ .

---

*Mathematics subject classification* (2010): Primary 93B36; Secondary 93D15, 46J15.

*Keywords and phrases:*  $v$ -metric, robust control, Banach algebras.

Such a desirable metric, was introduced by Glenn Vinnicombe in [15] and is called the  $v$ -metric. In that paper, essentially  $R$  was taken to be the rational functions without poles in the closed unit disk, and it was also shown that the topology obtained was equivalent to the one obtained from the gap-metric (introduced by Zames and El-Sakkary [16],[6], which in turn is equivalent to the graph metric of Vidyasagar [14]).

The problem of what happens when  $R$  is some other ring of stable transfer functions of infinite-dimensional systems was left open in [15]. This problem of extending the  $v$ -metric from the rational case to transfer function classes of infinite-dimensional systems was addressed in [1]. There the starting point in the approach was abstract. It was assumed that  $R$  is any commutative integral domain with identity which is a subset of a Banach algebra  $S$  satisfying certain assumptions, and then an “abstract”  $v$ -metric was defined in this setup, and it was shown in [1] that it does define a metric on the class of all stabilizable plants. It was also shown there that stabilizability is a robust property of the plant. In particular, this gave a metric on unstable plants over  $\mathcal{A}_+$ , where  $\mathcal{A}_+$  denotes the set of Laplace transforms of complex Borel measures  $\mu$  on  $[0, +\infty)$  such that  $\mu$  does not have a singular non-atomic part.

One can also define a gap-metric for unstable plants over  $\mathcal{A}_+$ , and so it is natural to ask if the  $v$ -metric and the gap-metric induce the same topologies on unstable plants over  $\mathcal{A}_+$ . In this article we address this issue, and prove the following result.

**THEOREM 1.1.** *On the set  $\mathbb{S}(\mathcal{A}_+)$ , the topologies induced by the  $v$ -metric  $d_v$  and the gap-metric  $d_g$  are identical.*

The notation  $\mathbb{S}(\mathcal{A}_+)$  will be explained carefully in Section 3, but roughly speaking, it is to be thought of as the class of unstable plants over  $\mathcal{A}_+$  with a single input and a single output. Owing to a technical difficulty, we restrict ourselves to single input and single output systems. We end this article with an open problem, namely the validity of our main result for systems with multiple inputs and multiple outputs, while pointing out the precise nature of the technical difficulty.

The paper is organized as follows:

1. In Section 3, we recall from [1] the  $v$ -metric in the context of unstable plants over  $\mathcal{A}_+$ , and also derive an alternative expression for it in Proposition 3.6, reminiscent of Georgiou’s formula for the gap-metric from [7].
2. In Section 4, we give the definition of the gap-metric in the context of unstable plants over  $\mathcal{A}_+$ . An alternative expression for the gap-metric is given in Proposition 4.9, which will be used in order to show the equivalence of  $d_v$  and  $d_g$ .
3. Finally, in Section 5, we will prove our main result (Theorem 1.1). At the end of this section, we also highlight the main obstacle towards extending Theorem 1.1 to systems with multiple inputs and outputs.

## 2. Notation index

For the convenience of the reader, we have included a table here which shows the page numbers of the places where the corresponding notation is first defined.

Notation	Page number
$\widehat{\cdot}$	Laplace transform (page 514) or Fourier transform (page 514)
$\cdot^*$	pages 514, 516, 520
$\mathcal{A}$	page 514
$\mathcal{A}_+$	page 514
$AP$	almost periodic functions (page 514)
$C_0$	continuous functions on $\mathbb{R}$ vanishing at $\pm\infty$ (page 522)
$\mathbb{C}_+$	right half of the complex plane (page 514)
$\overline{\delta}$	directed gap (page 519)
$d_g$	gap-metric (page 519)
$d_\nu$	$\nu$ -metric (page 517)
$\mathbb{F}(\mathcal{A}_+)$	field of fractions over $\mathcal{A}_+$ (page 516)
$\mathcal{G}$	graph of a system (page 519)
$G, \widetilde{G}, K, \widetilde{K}$	matrices built from coprime factorizations (page 516)
$\text{inv} \cdot$	invertible elements of a ring (page 513)
$\mu_{p,c}$	stability margin of the pair $(p, c)$ (page 517)
$P_{\mathcal{G}}$	projection onto $\mathcal{G}$ (page 519)
$P_{\mathcal{G}_1} _{\mathcal{G}_2}$	restriction of $P_{\mathcal{G}_1}$ to $\mathcal{G}_2$ (page 520)
$\mathbb{S}(\mathcal{A}_+)$	plants with a normalized coprime factorization (page 516)
$T_X$	Toeplitz operator (page 521)
$w$	winding number for continuous closed curves avoiding 0 (page 515)
$w$	average winding number for invertible $AP$ functions (page 515)
$W$	index for invertible elements in $\mathcal{A}$ (page 515)

## 3. The $\nu$ -metric

In this section we will recall the new  $\nu$ -metric for unstable plants over the ring  $\mathcal{A}_+$  (defined below), which was listed as a particular example in [1, Subsection 5.3] of the abstract  $\nu$ -metric introduced in that paper. At the end of this section, we will also give an alternate expression for the  $\nu$ -metric, which will be used later in order to show the equivalence of the  $\nu$ -metric topology with the classical gap topology.

If  $R$  is a commutative integral domain with identity 1, we use the symbol  $\text{inv } R$  for the set of invertible elements of  $R$ .

We denote by  $\mathcal{A}_+$  the set of Laplace transforms of complex Borel measures  $\mu$  on  $[0, +\infty)$  such that  $\mu$  does not have a singular non-atomic part. A more explicit

description of the elements of  $\mathcal{A}_+$  can be given as follows. Let

$$\mathbb{C}_+ := \{s \in \mathbb{C} : \operatorname{Re}(s) \geq 0\}.$$

Then

$$\mathcal{A}_+ = \left\{ s \in \mathbb{C}_+ \mapsto \widehat{f}_a(s) + \sum_{k \geq 0} f_k e^{-st_k} \mid \begin{array}{l} f_a \in L^1(0, \infty), (f_k)_{k \geq 0} \in \ell^1, \\ 0 = t_0 < t_1, t_2, t_3, \dots \end{array} \right\},$$

and equipped with pointwise operations and the norm:

$$\|F\| = \|f_a\|_{L^1} + \|(f_k)_{k \geq 0}\|_{\ell^1}, \quad F(s) = \widehat{f}_a(s) + \sum_{k \geq 0} f_k e^{-st_k} \quad (s \in \mathbb{C}_+),$$

$\mathcal{A}_+$  is a Banach algebra. Here  $\widehat{f}_a$  denotes the *Laplace transform* of  $f_a$ :

$$\widehat{f}_a(s) = \int_0^\infty e^{-st} f_a(t) dt, \quad s \in \mathbb{C}_+.$$

Similarly, define  $\mathcal{A}$  as follows:

$$\mathcal{A} = \left\{ iy \in i\mathbb{R} \mapsto \widehat{f}_a(iy) + \sum_{k \in \mathbb{Z}} f_k e^{-iyt_k} \mid \begin{array}{l} f_a \in L^1(\mathbb{R}), (f_k)_{k \in \mathbb{Z}} \in \ell^1, \\ \dots, t_{-2}, t_{-1} < 0 = t_0 < t_1, t_2, \dots \end{array} \right\}.$$

Then, equipped with pointwise operations and the norm:

$$\|F\| = \|f_a\|_{L^1} + \|(f_k)_{k \in \mathbb{Z}}\|_{\ell^1}, \quad F(iy) := \widehat{f}_a(iy) + \sum_{k \in \mathbb{Z}} f_k e^{-iyt_k} \quad (y \in \mathbb{R}),$$

$\mathcal{A}$  is a unital commutative complex semisimple Banach algebra. Here  $\widehat{f}_a$  is the *Fourier transform* of  $f_a$ ,

$$\widehat{f}_a(iy) = \int_{-\infty}^\infty e^{-iyt} f_a(t) dt \quad (y \in \mathbb{R}).$$

One can also define an involution  $\cdot^*$  on  $\mathcal{A}$ , given by

$$F^*(iy) = \overline{F(iy)}, \quad y \in \mathbb{R},$$

for  $F \in \mathcal{A}$ . Clearly,  $\mathcal{A}_+ \subset \mathcal{A}$ .

The algebra  $AP$  of complex valued (uniformly) *almost periodic functions* is the smallest closed subalgebra of  $L^\infty(\mathbb{R})$  that contains all the functions  $e_\lambda := e^{i\lambda y}$ . Here the parameter  $\lambda$  belongs to  $\mathbb{R}$ . For any  $f \in AP$ , its *Bohr-Fourier series* is defined by the formal sum

$$\sum_{\lambda} f_\lambda e^{i\lambda y}, \quad y \in \mathbb{R}, \tag{3.1}$$

where

$$f_\lambda := \lim_{N \rightarrow \infty} \frac{1}{2N} \int_{[-N, N]} e^{-i\lambda y} f(y) dy, \quad \lambda \in \mathbb{R},$$

and the sum in (3.1) is taken over the set  $\sigma(f) := \{\lambda \in \mathbb{R} \mid f_\lambda \neq 0\}$ , called the *Bohr-Fourier spectrum* of  $f$ . The Bohr-Fourier spectrum of every  $f \in AP$  is at most a countable set. For each  $f \in \text{inv } AP$ , we can define the *average winding number*  $w(f) \in \mathbb{R}$  of  $f$  as follows [9, Theorem 1, p. 167]:

$$w(f) = \lim_{T \rightarrow \infty} \frac{1}{2T} \left( \arg(f(T)) - \arg(f(-T)) \right).$$

We set

$$F_{AP}(iy) = \sum_{k \in \mathbb{Z}} f_k e^{-iyt_k} \quad (y \in \mathbb{R}) \quad \text{for} \quad F = \widehat{f}_a + \sum_{k \in \mathbb{Z}} f_k e^{-it_k} \in \mathcal{A}.$$

If  $F = \widehat{f}_a + F_{AP} \in \text{inv } \mathcal{A}$ , then it can be shown that ([1, Subsection 5.3])  $F_{AP}(i \cdot) \in \text{inv } AP$ . Moreover,  $F = \widehat{f}_a + F_{AP} \in \mathcal{A}$  is invertible if and only if for all  $y \in \mathbb{R}$ ,  $F(iy) \neq 0$  and  $\inf_{y \in \mathbb{R}} |F_{AP}(iy)| > 0$ .

Since  $L^1(\mathbb{R})$  is an ideal in  $\mathcal{A}$ , it follows that  $F_{AP}^{-1} \widehat{f}_a$  is the Fourier transform of a function in  $L^1(\mathbb{R})$ , and so the map

$$y \mapsto 1 + (F_{AP}(iy))^{-1} \widehat{f}_a(iy) = \frac{F(iy)}{F_{AP}(iy)}$$

has a well-defined winding number  $w$  around 0. Geometrically,  $w(f)$  is the number of times the curve  $t \mapsto f(t)$  winds around the origin in a counterclockwise direction.

Define the *index*  $W : \text{inv } \mathcal{A} \rightarrow \mathbb{R} \times \mathbb{Z}$  by

$$W(F) = \left( w(F_{AP}), w(1 + F_{AP}^{-1} \widehat{f}_a) \right), \tag{3.2}$$

where  $F = \widehat{f}_a + F_{AP} \in \text{inv } \mathcal{A}$ , and

$$\begin{aligned} w(F_{AP}) &:= \lim_{R \rightarrow \infty} \frac{1}{2R} \left( \arg(F_{AP}(iR)) - \arg(F_{AP}(-iR)) \right), \\ w(1 + F_{AP}^{-1} \widehat{f}_a) &:= \frac{1}{2\pi} \left( \arg(1 + (F_{AP}(iy))^{-1} \widehat{f}_a(iy)) \Big|_{y=-\infty}^{y=+\infty} \right). \end{aligned}$$

The map  $W : \text{inv } \mathcal{A} \rightarrow \mathbb{R} \times \mathbb{Z}$  satisfies:

- (I1)  $W(ab) = W(a) + W(b)$  ( $a, b \in \text{inv } \mathcal{A}$ ).
- (I2)  $W(a^*) = -W(a)$  ( $a \in \text{inv } \mathcal{A}$ ).
- (I3)  $W$  is locally constant, that is,  $W$  continuous when  $\mathbb{R} \times \mathbb{Z}$  is equipped with the discrete topology.
- (I4)  $x \in \mathcal{A}_+ \cap (\text{inv } \mathcal{A})$  is invertible as an element of  $\mathcal{A}_+$  if and only if  $W(x) = (0, 0)$ .

A consequence of (I3) is the following ‘‘homotopic invariance of the index’’ (see [1, Proposition 2.1]): if  $H : [0, 1] \rightarrow \text{inv } \mathcal{A}$  is a continuous map, then  $W(H(0)) = W(H(1))$ .

We recall the following standard notation and definitions from the factorization approach to control theory.

### 3.1. The notation $\mathbb{F}(\mathcal{A}_+)$ :

$\mathbb{F}(\mathcal{A}_+)$  denotes the field of fractions of  $\mathcal{A}_+$ .

### 3.2. The notation $F^*$ :

If  $F \in \mathcal{A}_+^{p \times m}$ , then  $F^* \in \mathcal{A}_+^{m \times p}$  is the matrix with the entry in the  $i$ th row and  $j$ th column given by  $F_{ji}^*$ , for all  $1 \leq i \leq p$ , and all  $1 \leq j \leq m$ .

### 3.3. Coprime/normalized coprime factorization:

Given  $p \in \mathbb{F}(R)$ , a factorization  $p = nd^{-1}$ , where  $n, d \in R$ , is called a *coprime factorization of  $P$*  if there exist  $x, y \in R$  such that  $xn + yd = 1$ . If moreover there holds that  $n^*n + d^*d = 1$ , then the coprime factorization is referred to as a *normalized coprime factorization of  $p$* .

### 3.4. The notation $G, \tilde{G}, K, \tilde{K}$ :

Given  $p \in \mathbb{F}(\mathcal{A}_+)$  with a normalized coprime factorization  $p = nd^{-1}$ , we introduce the following matrices with entries from  $\mathcal{A}_+$ :

$$G = \begin{bmatrix} n \\ d \end{bmatrix} \quad \text{and} \quad \tilde{G} = \begin{bmatrix} -d & n \end{bmatrix}.$$

Similarly, given  $c \in \mathbb{F}(\mathcal{A}_+)$  with normalized coprime factorization  $c = xy^{-1}$ , we introduce the following matrices with entries from  $\mathcal{A}_+$ :

$$K = \begin{bmatrix} y \\ x \end{bmatrix} \quad \text{and} \quad \tilde{K} = \begin{bmatrix} -x & y \end{bmatrix}.$$

### 3.5. The notation $\mathbb{S}(\mathcal{A}_+)$ :

We denote by  $\mathbb{S}(\mathcal{A}_+)$  the set of all elements  $p \in \mathbb{F}(\mathcal{A}_+)$  that possess a normalized coprime factorization.

REMARK 3.1.

1. It can be shown (see for example [14, Chapter 8]) that if  $p \in \mathbb{S}(\mathcal{A}_+)$ , then  $p$  is a *stabilizable plant over  $\mathcal{A}_+$* , that is, there exists a  $c \in \mathbb{F}(\mathcal{A}_+)$  such that  $H(p, c) \in \mathbb{R}^{2 \times 2}$ .
2. [2, Subsection 3.5] shows that every stabilizable plant  $p \in \mathbb{F}(\mathcal{A}_+)$  admits a coprime factorization over  $\mathcal{A}_+$ .
3. It follows from the proof of [10, Lemma 6.5.6.(e)] and [10, Theorem 5.2.8] that whenever  $p \in \mathbb{F}(\mathcal{A}_+)$  has a coprime factorization over  $\mathcal{A}_+$ , it also has a *normalized coprime factorization over  $\mathcal{A}_+$* .

Putting these remarks together, we see that  $\mathbb{S}(\mathcal{A}_+)$  is exactly the set of all plants in  $\mathbb{F}(\mathcal{A}_+)$  that are stabilizable over  $\mathcal{A}_+$ .

DEFINITION 3.2. ( $\nu$ -metric  $d_\nu$  on  $\mathbb{S}(\mathcal{A}_+)$ ) For  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , with the normalized coprime factorizations  $p_1 = n_1 d_1^{-1}$  and  $p_2 = n_2 d_2^{-1}$ , we define

$$d_\nu(p_1, p_2) := \begin{cases} \|\tilde{G}_2 G_1\|_\infty & \text{if } G_1^* G_2 \in \text{inv } \mathcal{A} \text{ and } W(G_1^* G_2) = (0, 0), \\ 1 & \text{otherwise.} \end{cases} \tag{3.3}$$

where the notation is as in Subsections 3.1-3.5.

We have the following; see [1]:

THEOREM 3.3.  $d_\nu$  given by (3.3) is a metric on  $\mathbb{S}(\mathcal{A}_+)$ .

Moreover, stabilizability is a robust property of the plant in this new  $\nu$ -metric. In order to see this, we first introduce the notion of stability margin for a pair comprising a plant and its controller.

DEFINITION 3.4. Given  $p, c \in \mathbb{F}(\mathcal{A}_+)$ , the *stability margin* of the pair  $(p, c)$  is defined by

$$\mu_{p,c} = \begin{cases} \|H(p, c)\|_\infty^{-1} & \text{if } p \text{ is stabilized by } c, \\ 0 & \text{otherwise.} \end{cases}$$

The number  $\mu_{p,c}$  can be interpreted as a measure of the performance of the closed loop system comprising  $p$  and  $c$ : larger values of  $\mu_{p,c}$  correspond to better performance, with  $\mu_{p,c} > 0$  if  $c$  stabilizes  $p$ .

The following was proved in [1]:

THEOREM 3.5. If  $p, p' \in \mathbb{S}(\mathcal{A}_+)$  and  $c \in \mathbb{S}(\mathcal{A}_+)$ , then  $\mu_{p',c} \geq \mu_{p,c} - d_\nu(p, p')$ .

The above result says that stabilizability is a robust property of the plant, since if  $c$  stabilizes  $p$  with a stability margin  $\mu_{p,c} > m$ , and  $p'$  is another plant which is close to  $p$  in the sense that  $d_\nu(p', p) \leq m$ , then  $c$  is also guaranteed to stabilize  $p'$ .

We will now derive an alternative expression for the  $\nu$ -metric, which is reminiscent of Georgiou’s formula for the gap-metric from [7].

PROPOSITION 3.6. If  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , then

$$d_\nu(p_1, p_2) = \inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q) = (0,0)}} \|G_1 - G_2 q\|_\infty.$$

*Proof.* Let  $q \in \text{inv } \mathcal{A}$  and  $W(q) = (0, 0)$ . We have

$$\begin{aligned} \|G_1 - G_2q\|_\infty &= \left\| \begin{bmatrix} G_2^* \\ \tilde{G}_2 \end{bmatrix} (G_1 - G_2q) \right\|_\infty \quad (\text{as } [G_2 \ \tilde{G}_2^*] \begin{bmatrix} G_2^* \\ \tilde{G}_2 \end{bmatrix} = I) \\ &= \left\| \begin{bmatrix} G_2^*G_1 - q \\ \tilde{G}_2G_1 \end{bmatrix} \right\|_\infty \quad (\text{since } \tilde{G}_2G_2 = 0 \text{ and } G_2^*G_2 = I) \\ &\geq \|\tilde{G}_2G_1\|_\infty. \end{aligned}$$

So if  $G_2^*G_1 \in \text{inv } \mathcal{A}$  and  $W(G_2^*G_1) = (0, 0)$ , then from the above it follows that  $\|G_1 - G_2q\|_\infty \geq \|\tilde{G}_2G_1\|_\infty = d_V(p_1, p_2)$ . As the choice of  $q$  above was arbitrary, we obtain

$$\inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q)=(0,0)}} \|G_1 - G_2q\|_\infty \geq d_V(p_1, p_2). \tag{3.4}$$

If we define  $q_0 := G_2^*G_1 \in \mathcal{A}$ , then  $q_0 \in \text{inv } \mathcal{A}$  and  $W(q_0) = (0, 0)$ , and so

$$\begin{aligned} \inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q)=(0,0)}} \|G_1 - G_2q\|_\infty &\leq \|G_1 - G_2q_0\|_\infty = \left\| \begin{bmatrix} G_2^*G_1 - q_0 \\ \tilde{G}_2G_1 \end{bmatrix} \right\|_\infty \\ &= \left\| \begin{bmatrix} 0 \\ \tilde{G}_2G_1 \end{bmatrix} \right\|_\infty = \|\tilde{G}_2G_1\|_\infty = d_V(p_1, p_2). \end{aligned}$$

From this and (3.4), the claim in the proposition follows for the case when  $G_2^*G_1 \in \text{inv } \mathcal{A}$  and  $W(G_2^*G_1) = (0, 0)$ .

Now let  $q \in \text{inv } \mathcal{A}$  be such that  $W(q) = (0, 0)$  and  $\|G_1 - G_2q\|_\infty < 1$ . Using  $G_1^*G_1 = 1$ , we see that

$$\|1 - G_1^*G_2q\|_\infty = \|G_1^*(G_1 - G_2q)\|_\infty \leq \|G_1^*\|_\infty \|G_1 - G_2q\|_\infty < 1 \cdot 1 = 1.$$

So  $G_1^*G_2q = 1 - (1 - G_1^*G_2q)$  is invertible as an element of  $\mathcal{A}$ . Consider the map  $H : [0, 1] \rightarrow \text{inv } \mathcal{A}$  given by  $H(t) = 1 - t(1 - G_1^*G_2q)$ ,  $t \in [0, 1]$ . By the homotopic invariance of the index,

$$(0, 0) = W(1) = W(H(0)) = W(H(1)) = W(G_1^*G_2q).$$

As  $W(q) = (0, 0)$ , we obtain that  $W(G_1^*G_2) = (0, 0)$ . So we have shown that if there is a  $q \in \mathcal{A}$  such that  $q \in \text{inv } \mathcal{A}$ ,  $W(q) = (0, 0)$  and  $\|G_1 - G_2q\|_\infty < 1$ , then  $G_1^*G_2 \in \text{inv } \mathcal{A}$  and  $W(G_1^*G_2) = (0, 0)$ . Thus if either  $G_1^*G_2 \notin \text{inv } \mathcal{A}$  or  $G_1^*G_2 \in \text{inv } \mathcal{A}$  but  $W(G_1^*G_2) \neq (0, 0)$ , then for all  $q \in \mathcal{A}$  such that  $q \in \text{inv } \mathcal{A}$ ,  $W(q) = (0, 0)$ , we have that  $\|G_1 - G_2q\|_\infty \geq 1$ , and so

$$\inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q)=(0,0)}} \|G_1 - G_2q\|_\infty \geq 1 = d_V(p_1, p_2).$$

Also, with  $q_n := \frac{1}{n}I$ ,  $q_n \in \text{inv } \mathcal{A}$  and  $W(q_n) = (0, 0)$ . We have

$$\|G_1 - G_2q_n\|_\infty \leq \|G_1\|_\infty + \|G_2\|_\infty \|q_n\|_\infty \leq 1 + 1 \cdot \frac{1}{n}.$$

Hence

$$\inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q)=(0,0)}} \|G_1 - G_2q\|_\infty \leq \inf_n \|G_1 - G_2q_n\|_\infty \leq \inf_n \left(1 + \frac{1}{n}\right) = 1 = d_v(p_1, p_2).$$

Consequently,  $\inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q)=(0,0)}} \|G_1 - G_2q\|_\infty = 1 = d_v(p_1, p_2)$ .  $\square$

### 4. The gap-metric

In this section we will recall the gap-metric topology for unstable plants over the ring  $\mathcal{A}_+$ . We will also prove a few technical lemmas which will be used in the next section in order to prove our main result.

DEFINITION 4.1. (Graph of a system) For  $p \in \mathbb{S}(\mathcal{A}_+)$ , with the normalized co-prime factorization  $p = nd^{-1}$ , we define the *graph of  $p$* , denoted by  $\mathcal{G}$ , to be the following subspace of the Hardy space  $H^2(\mathbb{C}^2)$ :

$$\mathcal{G} = GH^2 = \left\{ \begin{bmatrix} n\varphi \\ d\varphi \end{bmatrix} : \varphi \in H^2 \right\}.$$

Using the fact that there exist  $x, y \in \mathcal{A}_+$  such that  $xn + yd = 1$ , it is easy to see that the graph  $\mathcal{G}$  is a *closed* subspace of  $H^2 \times H^2$ . We denote the orthogonal projection from  $H^2 \times H^2$  onto  $\mathcal{G}$  by  $P_{\mathcal{G}}$ .

DEFINITION 4.2. (Gap-metric  $d_g$ ) For  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , with the normalized co-prime factorizations  $p_1 = n_1d_1^{-1}$  and  $p_2 = n_2d_2^{-1}$ , we define

$$d_g(p_1, p_2) := \|P_{\mathcal{G}_1} - P_{\mathcal{G}_2}\|_{\mathcal{L}(H^2 \times H^2)}. \tag{4.1}$$

We will need a few technical results on the gap-metric  $d_g$ . For a self-contained account of these results, we refer the reader to [13]. It can be checked that  $d_g$  given by (4.1) is well-defined. Since the gap-metric is a metric on the set of closed subspaces of a Hilbert space, it follows that  $d_g$  given by (4.1) is a metric on  $\mathbb{S}(\mathcal{A}_+)$ .

For  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ ,  $d_g(p_1, p_2) = \max\{\vec{\delta}(p_1, p_2), \vec{\delta}(p_2, p_1)\}$ , where  $\vec{\delta}(\cdot, \cdot)$  denotes the *directed gap*, defined by

$$\vec{\delta}(p_1, p_2) := \|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}\|_{\mathcal{L}(H^2 \times H^2)}.$$

If  $d_g(p_1, p_2) < 1$ , then  $d_g(p_1, p_2) = \vec{\delta}(p_1, p_2) = \vec{\delta}(p_2, p_1)$  [8, Prop. 3, p. 675]. In [7], it was shown that

$$d_g(p_1, p_2) = \max \left\{ \inf_{q \in H^\infty} \|G_1 - G_2q\|_\infty, \inf_{q \in H^\infty} \|G_2 - G_1q\|_\infty \right\}.$$

For  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , the infimums above can be taken over  $\mathcal{A}_+$  instead of  $H^\infty$ , and this follows from [10, Theorem 11.3.3].

LEMMA 4.3. *If  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , then*

$$\inf_{q \in H^\infty} \|G_1 - G_2q\|_\infty = \inf_{q \in \mathcal{A}_+} \|G_1 - G_2q\|_\infty.$$

*Proof.* Clearly  $m := \inf_{q \in H^\infty} \|G_1 - G_2q\|_\infty \leq \inf_{q \in \mathcal{A}_+} \|G_1 - G_2q\|_\infty =: M$ . Define

$$V = \begin{bmatrix} G_2 & G_1 \\ 0 & 1 \end{bmatrix}, \quad W := V^* \begin{bmatrix} I & 0 \\ 0 & -M^2 \end{bmatrix} V.$$

(For  $X \in (H^\infty)^{p \times m}$ ,  $X^* \in (L^\infty)^{m \times p}$  is defined by  $X^*(iy) = (X(iy))^*$ ,  $y \in \mathbb{R}$ .) Suppose that  $m < M$ . Then there exists a  $q \in H^\infty$  such that  $\|G_1 - G_2q\|_\infty < M$ . Now we apply [10, Theorem 11.3.3, p. 654] to conclude that the  $q$  can in fact be chosen in  $\mathcal{A}_+$ . For this, a few technical assumptions have to be verified first, and we give these details in the following paragraph for the interested reader.

(First of all, the Standing Hypothesis [10, 11.0.1, p. 611] is satisfied, since  $\mathcal{A}_+$  does satisfy [10, Hypothesis 8.4.7., p. 384], by [10, Theorem 8.4.9( $\beta$ ), p. 385]. Secondly, the Standing Hypothesis [10, 11.3.1, p. 654] is satisfied, since  $G_2^*G_2 = 1$ . Actually, there are two extraneous assumptions in 11.3.1, but neither is used in the part of the proofs required here, and these extraneous assumptions are anyway satisfied in our case. Now as the Assumption (FI1  $\frac{1}{2}$ s) of [10, Theorem 11.3.3, p. 654] holds, also (FI1s) holds. By the last sentence of [10, Theorem 11.3.6, p. 659], as  $W$  has entries from  $\mathcal{A}_+$ , there exists a  $q \in \mathcal{A}_+$  such that  $\|G_1 - G_2q\|_\infty < M$ .)

Consequently,  $m = M$ .  $\square$

We use the notation  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  to mean the restriction of  $P_{\mathcal{G}_1}$  to  $\mathcal{G}_2$ , namely, the operator from  $\mathcal{G}_2$  to  $\mathcal{G}_1$ , given by

$$P_{\mathcal{G}_1}|_{\mathcal{G}_2}g_2 = P_{\mathcal{G}_1}g_2, \quad g_2 \in \mathcal{G}_2.$$

Then  $\ker(P_{\mathcal{G}_1}|_{\mathcal{G}_2}) = \{g_2 \in \mathcal{G}_2 : P_{\mathcal{G}_1}g_2 = 0\} = \mathcal{G}_2 \cap (\ker P_{\mathcal{G}_1}) = \mathcal{G}_2 \cap \mathcal{G}_1^\perp$ . Also, for  $g_1 \in \mathcal{G}_1$  and  $g_2 \in \mathcal{G}_2$ , we have

$$\begin{aligned} \langle P_{\mathcal{G}_1}|_{\mathcal{G}_2}g_2, g_1 \rangle_{\mathcal{G}_1} &= \langle P_{\mathcal{G}_1}g_2, g_1 \rangle_{\mathcal{G}_1} = \langle g_2, g_1 \rangle_{H^2(\mathbb{C}^2)} \\ &= \langle g_2, P_{\mathcal{G}_2}g_1 \rangle_{H^2(\mathbb{C}^2)} = \langle g_2, P_{\mathcal{G}_2}g_1 \rangle_{\mathcal{G}_2} \\ &= \langle g_2, P_{\mathcal{G}_2}|_{\mathcal{G}_1}g_1 \rangle_{\mathcal{G}_2}, \end{aligned}$$

and so  $(P_{\mathcal{G}_1}|_{\mathcal{G}_2})^* = P_{\mathcal{G}_2}|_{\mathcal{G}_1}$ . Thus  $\ker((P_{\mathcal{G}_1}|_{\mathcal{G}_2})^*) = \ker(P_{\mathcal{G}_2}|_{\mathcal{G}_1}) = \mathcal{G}_1 \cap \mathcal{G}_2^\perp$ . So if  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is a Fredholm operator [12, §2.5.1, p. 218], then its Fredholm index is given by  $\dim(\mathcal{G}_2 \cap \mathcal{G}_1^\perp) - \dim(\mathcal{G}_1 \cap \mathcal{G}_2^\perp)$ .

We will use the following result from [11, p. 201].

LEMMA 4.4. (Lemma on Closed Subspaces) *Let  $H$  be a Hilbert space and let  $U, V$  be subspaces of  $H$ . Then the following are equivalent:*

(SI)  $U \cap V^\perp = \{0\}$ .

(S2) Closure of  $P_U V$  is  $U$ .

Also, the following are equivalent:

(S3)  $P_U V = U$  and  $V \cap U^\perp = \{0\}$ .

(S4)  $\|(I - P_U)P_U\| < 1$  and  $\|(I - P_U)P_V\| < 1$ .

LEMMA 4.5. Let  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ . Then  $d_g(p_1, p_2) < 1$  if and only if the following three conditions hold:

1.  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is Fredholm,
2.  $\mathcal{G}_1 \cap \mathcal{G}_2^\perp = \{0\}$ , and
3.  $\mathcal{G}_2 \cap \mathcal{G}_1^\perp = \{0\}$ .

*Proof.* (Only if) As  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is Fredholm, its range is closed, that is,  $P_{\mathcal{G}_1}\mathcal{G}_2$  is a closed subspace. Hence from the equivalence of (S1) with (S2) in Lemma 4.4 above, we have that the closure of  $P_{\mathcal{G}_1}\mathcal{G}_2$ , which is the same as  $P_{\mathcal{G}_1}\mathcal{G}_2$ , is equal to  $\mathcal{G}_1$ . Now from the equivalence of (S3) with (S4) in Lemma 4.4, we obtain that  $\vec{\delta}(p_1, p_2) = \|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}\| < 1$  and  $\vec{\delta}(p_2, p_1) = \|(I - P_{\mathcal{G}_1})P_{\mathcal{G}_2}\| < 1$ . Hence  $d_g(p_1, p_2) < 1$ .

(If) As  $\vec{\delta}(p_1, p_2) = \|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}\| < 1$  and  $\vec{\delta}(p_2, p_1) = \|(I - P_{\mathcal{G}_1})P_{\mathcal{G}_2}\| < 1$ , by the equivalence of (S3) with (S4) in Lemma 4.4, we obtain  $P_{\mathcal{G}_1}\mathcal{G}_2 = \mathcal{G}_1$ , and so the range of  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is closed. Moreover,  $\mathcal{G}_2 \cap \mathcal{G}_1^\perp = \{0\}$ . By interchanging the roles of  $p_1$  and  $p_2$ , we also get that  $\mathcal{G}_1 \cap \mathcal{G}_2^\perp = \{0\}$ .  $\square$

The following is easy to check.

LEMMA 4.6. Let  $H_1, H_2$  be Hilbert spaces and  $T \in \mathcal{L}(H_1, H_2)$ ,  $S \in \mathcal{L}(H_2, H_1)$  be such that  $ST = I$ . Suppose that  $U$  is a subspace of  $H_1$ . Then we have that  $TU$  is closed if and only if  $U$  is closed.

*Proof.* (If) Since  $T$  is left invertible,  $\|x\| = \|STx\| \leq \|S\|\|Tx\|$  ( $x \in H_1$ ). Suppose  $(y_n) = (Tx_n)$  ( $x_n \in U$ ) is a sequence that converges in  $H_2$ . Thus  $\|y_n - y_m\| \geq \frac{1}{\|S\|} \|x_n - x_m\|$ , showing that  $(x_n)$  must converge to some  $x \in H_1$ . As  $U$  is closed,  $x \in U$ . Thus  $y_n = Tx_n \rightarrow Tx \in TU$ . Hence  $TU$  is closed.

(Only if) Now suppose that  $TU$  is closed. If  $(x_n)$  is a sequence in  $U$  that converges to  $x$  in  $H_1$ , then clearly  $Tx_n \rightarrow Tx$ . But  $TU$  is closed, and so  $Tx \in TU$ . Hence  $Tx = Tx'$  for some  $x' \in U$ . Operating by  $S$ , we have  $x = STx = STx' = x'$ , and so  $x = x' \in U$ . Thus  $U$  is closed.  $\square$

For  $X \in (L^\infty)^{p \times m}$ ,  $T_X$  denotes the Toeplitz operator from  $(H^2)^m$  to  $(H^2)^p$ , given by  $T_X \varphi = \Pi_{(H^2)^p}(X\varphi)$  ( $\varphi \in (H^2)^m$ ), where  $X\varphi$  is considered as an element of  $(L^2)^p$  and  $\Pi_{(H^2)^p}$  denotes the canonical orthogonal projection from  $(L^2)^p$  onto  $(H^2)^p$ .

LEMMA 4.7. *Let  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ . Then  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is Fredholm if and only if  $T_{G_1^*G_2}$  is Fredholm. Moreover, their Fredholm indices coincide.*

*Proof.* First of all, we note that  $T_{G_1^*G_2} = T_{G_1^*}T_{G_2}$  (since  $G_2$  has  $H^\infty$  entries). Also, it can be checked that for a matrix  $X$  with  $L^\infty$  entries  $(T_X)^* = T_{X^*}$ . Thus  $(T_{G_1^*G_2})^* = T_{G_2^*G_1}$ .

As  $T_{G_1}$  is an isometry, it follows that the orthogonal projection onto the range of  $T_{G_1}$ , namely the subspace  $\mathcal{G}_1$ , is given by  $T_{G_1}(T_{G_1})^* = T_{G_1}T_{G_1^*}$ . Indeed, with  $P := T_{G_1}T_{G_1^*}$ , and using  $G_1^*G_1 = 1$ , we can check that  $P^2 = P$ , that  $P^* = P$  and that  $P$  maps onto the range of  $T_{G_1}$ :

$$\text{ran}(T_{G_1}T_{G_1^*}) \subset \text{ran} T_{G_1} = \text{ran}(T_{G_1}T_{G_1^*}T_{G_1}) \subset \text{ran}(T_{G_1}T_{G_1^*}).$$

We have that

$$\begin{aligned} \ker(T_{G_1^*}T_{G_2}) &= \{\varphi \in H^2 : T_{G_1^*}T_{G_2}\varphi = 0\} \\ &= \{\varphi \in H^2 : T_{G_1}T_{G_1^*}T_{G_2}\varphi = 0\} \quad (\text{since } [x_1 \ y_1]G_1 = 1) \\ &= \{\varphi \in H^2 : P_{\mathcal{G}_1}T_{G_2}\varphi = 0\} = \{\varphi \in H^2 : T_{G_2}\varphi \in \mathcal{G}_1^\perp\}. \end{aligned}$$

Consider the map  $\iota : \ker(T_{G_1^*}T_{G_2}) \rightarrow \mathcal{G}_1^\perp \cap \mathcal{G}_2$  defined by  $\iota(\varphi) = T_{G_2}\varphi$  for  $\varphi \in \ker(T_{G_1^*}T_{G_2})$ . From the above calculation, we see that  $\iota$  is onto. Also, since  $[x_2 \ y_2]G_2 = 1$  it follows that  $\iota$  is one-to-one. So  $\iota$  is invertible.

The above shows that in case that  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  and  $T_{G_1^*G_2}$  are both Fredholm operators, their Fredholm indices will coincide.

In light of the above, we just need to show that the range of  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is closed if and only if the range of  $T_{G_1^*G_2}$  is closed. The range of  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is

$$P_{\mathcal{G}_1}\mathcal{G}_2 = P_{\mathcal{G}_1}\text{ran} T_{G_2} = T_{G_1}T_{G_1^*}\text{ran} T_{G_2} = T_{G_1}\text{ran} T_{G_1^*G_2}.$$

Since  $G_1$  has a left inverse  $[x_1 \ y_1] \in \mathcal{A}_+^2$ , it follows that  $T_{G_1}$  is left-invertible. By Lemma 4.6, the range of  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is closed if and only if the range of  $\text{ran} T_{G_1^*G_2}$  is closed.  $\square$

We will need the following result, which follows from [4, Thm. 3, p. 150]. Here  $C_0$  denotes the set of continuous functions on  $\mathbb{R}$  that vanish at  $\pm\infty$ .

PROPOSITION 4.8. *Let  $F = f + g$ , where  $f \in AP$  and  $g \in C_0$  be such that  $T_F$  is Fredholm. Then the following hold:*

1.  $T_f$  is invertible.
2.  $F \in \text{inv}(AP + C_0)$ .
3. The Fredholm index of  $T_F$  is the winding number of  $1 + f^{-1}g$ .

*Proof.* Since  $T_F$  is invertible modulo the compacts, it is invertible modulo any bigger ideal which we can take to be the kernel of the symbol map from the Toeplitz  $C^*$ -algebra  $\mathcal{T}(AP + C_0)$  (generated by  $T_\varphi$  for  $\varphi \in AP + C_0$ ) to  $AP + C_0$ . Thus  $F \in \text{inv}(AP + C_0)$ .

It follows from the invertibility of  $F = f + g$  in  $AP + C_0$  that there are  $\tilde{f} \in AP$  and  $\tilde{g} \in C_0$  such that

$$1 = (f + g)(\tilde{f} + \tilde{g}) = f\tilde{f} + (f\tilde{g} + g\tilde{f} + g\tilde{g}).$$

But  $f\tilde{g} + g\tilde{f} + g\tilde{g} \in C_0$  because  $C_0$  is an ideal in  $AP + C_0$ . So

$$f\tilde{f} - 1 \in AP \cap C_0.$$

By [3, 9.19, p. 407],  $AP \cap C_0 = \{0\}$ . Consequently,  $f\tilde{f} = 1$ , that is,  $f \in \text{inv } AP$ . Moreover, using [4, Theorem 3, p. 150] (see also [3, Theorem 9.22]), one knows that the generalized index of  $F$  is  $(0, n)$  for some integer  $n$  and hence the average winding number of  $f$  is zero. Thus  $T_f$  is invertible [5, Theorem 11, p. 25].

Again using [4, Theorem 3, p. 150], one can see that the generalized index of  $T_F$  equals the sum of the generalized indices of  $T_f$  and  $T_{1+f^{-1}g}$ . But the generalized index of  $T_f$  is  $(0, 0)$  which completes the proof.  $\square$

**PROPOSITION 4.9.** *If  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ , then*

$$d_g(p_1, p_2) = \inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty$$

*Proof.*  $\underline{1}^\circ$  Consider first the case when  $d_g(p_1, p_2) < 1$ . From Lemma 4.5, it follows that  $P_{\mathcal{G}_1|_{\mathcal{G}_2}}$  is Fredholm,  $\mathcal{G}_1 \cap \mathcal{G}_2^\perp = \{0\}$  and  $\mathcal{G}_2 \cap \mathcal{G}_1^\perp = \{0\}$ . Furthermore, the Fredholm index of  $P_{\mathcal{G}_1|_{\mathcal{G}_2}}$  is 0. By Lemma 4.7,  $T_{G_1^*G_2}$  is Fredholm, with Fredholm index 0 too. From Proposition 4.8, it follows that  $G_1^*G_2$  is invertible as an element of  $AP + C_0$ . Thus it is also invertible as an element of  $\mathcal{A}$ . Also,  $W(G_1^*G_2) = (0, 0)$ . Now suppose that there is a  $q_0 \in \mathcal{A}_+$  such that  $\|G_1 - G_2q_0\|_\infty < 1$ . Then  $\|I - G_1^*G_2q_0\|_\infty < 1$  and so  $G_1^*G_2q_0 = 1 - (I - G_1^*G_2q_0)$  is invertible in  $\mathcal{A}$ . Hence  $G_1^*G_2q_0 \in \text{inv } \mathcal{A}$ . In particular,  $q_0 \in \text{inv } \mathcal{A}$ . Consider the map  $H : [0, 1] \rightarrow \text{inv } \mathcal{A}$  given by  $H(t) = 1 - t(I - G_1^*G_2q_0)$ ,  $t \in [0, 1]$ . By the homotopic invariance,

$$(0, 0) = W(1) = W(H(0)) = W(H(1)) = W(G_1^*G_2q_0).$$

Since  $W(G_1^*G_2) = (0, 0)$ , it follows that  $W(q_0) = (0, 0)$ . Thus by (I4), we obtain that  $q_0 \in \text{inv } \mathcal{A}_+$ . Consequently,

$$1 > d_g(p_1, p_2) = \tilde{\delta}(p_1, p_2) = \inf_{q \in \mathcal{A}_+} \|G_1 - G_2q\|_\infty = \inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty.$$

$\underline{2}^\circ$  Now suppose that  $d_g(p_1, p_2) = 1$ , but that  $\tilde{\delta}(p_1, p_2) < 1$ . Since we have  $\tilde{\delta}(p_1, p_2) = \|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}\|$ , we obtain  $\mathcal{G}_1 \cap \mathcal{G}_2^\perp = \{0\}$ . For otherwise, if  $0 \neq v \in \mathcal{G}_1 \cap \mathcal{G}_2^\perp$ , then we have  $(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}v = v$ , and so we would obtain that  $\|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}\| \geq$

$\|(I - P_{\mathcal{G}_2})P_{\mathcal{G}_1}v\|/\|v\| = 1$ , a contradiction. From Lemma 4.5, it now follows that either  $\mathcal{G}_2 \cap \mathcal{G}_1^\perp \neq \{0\}$  or  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is not Fredholm.

Suppose first that  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is Fredholm. Then we must have  $\mathcal{G}_2 \cap \mathcal{G}_1^\perp \neq \{0\}$ . This gives that the Fredholm index of  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$ , namely

$$\dim(\mathcal{G}_2 \cap \mathcal{G}_1^\perp) - \dim(\mathcal{G}_1 \cap \mathcal{G}_2^\perp) = \dim(\mathcal{G}_2 \cap \mathcal{G}_1^\perp) - 0 = \dim(\mathcal{G}_2 \cap \mathcal{G}_1^\perp),$$

is nonzero. By Lemma 4.7,  $T_{G_1^*G_2}$  is Fredholm, with Fredholm index nonzero too. It now follows from Proposition 4.8, that  $W(G_1^*G_2) = (*, n)$  with the integer  $n \neq 0$ . By the definition of  $d_v$ ,  $d_v(p_1, p_2) = 1$ .

Next assume that  $P_{\mathcal{G}_1}|_{\mathcal{G}_2}$  is not Fredholm. Then Lemma 4.7 gives that  $T_{G_1^*G_2}$  is not Fredholm either. Now if  $G_1^*G_2$  is not invertible in  $\mathcal{A}$ , then we have  $d_v(p_1, p_2) = 1$  by definition. On the other hand, if  $G_1^*G_2 \in \text{inv } \mathcal{A}$  and  $W(G_1^*G_2) = (0, 0)$ , it follows from [5, Proposition 6.3, p. 27] that  $T_{G_1^*G_2}$  is invertible, a contradiction. Thus  $W(G_1^*G_2) = (0, 0)$ , and so  $d_v(p_1, p_2) = 1$  in this case as well.

Now that we have obtained  $d_v(p_1, p_2) = 1$ , it follows that there is no  $q \in \text{inv } \mathcal{A}_+$  such that  $\|G_1 - G_2q\|_\infty < 1$ . In other words, for each  $q \in \mathcal{A}_+$ ,  $\|G_1 - G_2q\|_\infty \geq 1$ . Also,  $\|G_1 - G_2q\|_\infty \leq \|G_1\|_\infty + \|G_2\|_\infty\|q\|_\infty \leq 1 + 1 \cdot \|q\|_\infty$ , and by taking  $q_1 = \frac{1}{n}I \in \text{inv } \mathcal{A}_+$ , we obtain

$$\inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty \leq \inf_n \left(1 + \frac{1}{n}\right) = 1.$$

Consequently,  $\inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty = 1 = d_g(p_1, p_2)$ .

3° Now suppose that  $d_g(p_1, p_2) = 1 = \vec{\delta}(p_1, p_2) = \vec{\delta}(p_2, p_1)$ . We have

$$\inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty \leq \inf_n \left\|G_1 - G_2 \frac{1}{n}I\right\|_\infty \leq \inf_n \left(1 + \frac{1}{n}\right) = 1.$$

Also,  $1 = \vec{\delta}(p_1, p_2) = \inf_{q \in \mathcal{A}_+} \|G_1 - G_2q\|_\infty \leq \inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty$ . Thus

$$\inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty = 1 = d_g(p_1, p_2).$$

This completes the proof.  $\square$

### 5. Equivalence of the $v$ -metric and the gap-metric

*Proof of Theorem 1.1.* We will show the following for  $p_1, p_2 \in \mathbb{S}(\mathcal{A}_+)$ :

$$d_g(p_1, p_2)\mu_{\text{opt}}(p_1) \leq d_v(p_1, p_2) \leq d_g(p_1, p_2), \tag{5.1}$$

where  $\mu_{\text{opt}}(p_1) := \sup_c \mu_{p_1, c}$ .

This will prove the fact that the topologies induced by metrics  $d_g$  and  $d_v$  on the set  $\mathbb{S}(\mathcal{A}_+)$  are identical.

The second inequality in (5.1) is an immediate consequence of the Propositions 3.6 and 4.9. Indeed, we have

$$\begin{aligned}
 d_v(p_1, p_2) &= \inf_{\substack{q \in \text{inv } \mathcal{A}, \\ W(q) = (0,0)}} \|G_1 - G_2q\|_\infty \\
 &\leq \inf_{\substack{q \in \mathcal{A}_+ \cap (\text{inv } \mathcal{A}), \\ W(q) = (0,0)}} \|G_1 - G_2q\|_\infty \\
 &= \inf_{q \in \text{inv } \mathcal{A}_+} \|G_1 - G_2q\|_\infty \quad (\text{using (I4)}) \\
 &= d_g(p_1, p_2).
 \end{aligned}$$

Now we will show the first inequality in (5.1). This inequality is trivially satisfied if  $d_v(p_1, p_2) \geq \mu_{\text{opt}}(p_1)$ , since  $d_g(p_1, p_2) \leq 1$ . So we will only consider the case when  $d_v(p_1, p_2) < \mu_{\text{opt}}(p_1)$ . Thus we can choose a  $c$  that stabilizes both  $p_1$  and  $p_2$ . (Since the above inequality shows that there exists a  $c_0$  stabilizing  $p_1$  such that  $d_v(p_1, p_2) < \mu_{p_1, c_0}$ . But by Theorem 3.5, it follows that  $c_0$  also stabilizes  $p_2$ .) If we now define  $q_0 := (\tilde{K}_0 G_1)^{-1} \tilde{K}_0 G_2$ , then we have  $G_2 - G_1 q_0 = G_2 - G_1 (\tilde{K}_0 G_1)^{-1} \tilde{K}_0 G_2 = (I - G_1 (\tilde{K}_0 G_1)^{-1} \tilde{K}_0) G_2$ . Also

$$I - \begin{bmatrix} p_1 \\ 1 \end{bmatrix} (1 - c_0 p_1)^{-1} \begin{bmatrix} -c_0 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ c_0 \end{bmatrix} (1 - p_1 c_0)^{-1} \begin{bmatrix} 1 & -p_1 \end{bmatrix}.$$

that is,  $I - G_1 (\tilde{K}_0 G_1)^{-1} \tilde{K}_0 = K_0 (\tilde{G}_1 K_0)^{-1} \tilde{G}_1$ . Thus

$$G_2 - G_1 q_0 = K_0 (\tilde{G}_1 K_0)^{-1} \tilde{G}_1 G_2.$$

Then we use  $\|K_0\| \leq 1$  (since  $K_0^* K_0 = 1$ ) to obtain

$$\begin{aligned}
 \|G_2 - G_1 q_0\|_\infty &= \|K_0 (\tilde{G}_1 K_0)^{-1} \tilde{G}_1 G_2\|_\infty \\
 &\leq \|K_0\|_\infty \|(\tilde{G}_1 K_0)^{-1} \tilde{G}_1 G_2\|_\infty \\
 &\leq 1 \cdot \|(\tilde{G}_1 K_0)^{-1} \tilde{G}_1 G_2\|_\infty \\
 &\leq \|(\tilde{G}_1 K_0)^{-1}\|_\infty \|\tilde{G}_1 G_2\|_\infty.
 \end{aligned}$$

As for each  $c$ ,  $\mu_{p_1, c} \leq 1$ , we have  $\mu_{\text{opt}}(p_1) \leq 1$ . So  $d_v(p_1, p_2) < \mu_{\text{opt}}(p_1) \leq 1$ , and we obtain  $d_v(p_1, p_2) = \|(\tilde{G}_1 G_2)\|_\infty$ .

From [1, Propositions 4.2,4.5],  $\|(\tilde{G}_1 K_0)^{-1}\|_\infty = 1/\mu_{c_0, p_1} = 1/\mu_{p_1, c_0}$ . So

$$\|G_2 - G_1 q_0\|_\infty \leq \|(\tilde{G}_1 K_0)^{-1}\|_\infty \|\tilde{G}_1 G_2\|_\infty \leq \frac{d_v(p_1, p_2)}{\mu_{p_1, c_0}}.$$

Thus

$$d_g(p_1, p_2) = \inf_{q \in \text{inv } \mathcal{A}} \|G_1 - G_2q\|_\infty \leq \|G_1 - G_2q_0\|_\infty \leq d_v(p_1, p_2)/\mu_{p_1, c_0}.$$

This inequality holds for any  $c_0$  that stabilizes  $p_1$  for which there holds  $d_v(p_1, p_2) < \mu_{p_1, c_0}$ . We can choose a sequence  $(c_{0,n})$  such  $\mu_{p_1, c_{0,n}} \rightarrow \mu_{\text{opt}}(p_1)$  as  $n \rightarrow \infty$ . Thus  $d_g(p_1, p_2) \leq d_v(p_1, p_2) / \mu_{\text{opt}}(p_1)$ . This completes the proof of the first inequality in (5.1).  $\square$

The question of whether our main result, Theorem 1.1 remains true for systems with multiple inputs and multiple outputs (as opposed to just *scalar* inputs and outputs) is open. A key technical difficulty is the validity of the analogue of Proposition 4.8 for matricial data. In this connection, we also point out [3, Remark 9.33, p. 413] to highlight the gravity of the problem.

## Acknowledgements

The author thanks Professor Ronald Douglas for providing clarifications in [4, Theorem 3], which are presented in Proposition 4.8 of this article. Part (3) of the Remark 3.1 and the proof of Lemma 4.3 are due to Kalle Mikkola, and the author thanks him for these. A useful discussion with Professor Joseph Ball is also gratefully acknowledged.

## REFERENCES

- [1] J. A. BALL AND A. J. SASANE, *Extension of the  $v$ -metric*, Complex Analysis and Operator Theory, to appear.
- [2] A. BRUDNYI AND A. J. SASANE, *Sufficient conditions for the projective freeness of Banach algebras*, Journal of Functional Analysis, in press.
- [3] A. BÖTTCHER AND B. SILBERMANN, *Analysis of Toeplitz operators*, Springer-Verlag, Berlin, 1990.
- [4] R. G. DOUGLAS, *On the  $C^*$ -algebra of a one-parameter semigroup of isometries*, Acta Mathematica **128**, 3–4 (1972), 143–151.
- [5] R. G. DOUGLAS, *Banach algebra techniques in the theory of Toeplitz operators*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 15. American Mathematical Society, Providence, R.I., 1973.
- [6] A. K. EL-SAKKARY, *The gap metric: robustness of stabilization of feedback systems*, IEEE Transactions on Automatic Control **30**, 3 (1985), 240–247.
- [7] T. T. GEORGIU, *On the computation of the gap metric*, Systems Control Letters **11**, 4 (1988), 253–257.
- [8] T. T. GEORGIU AND M. C. SMITH, *Optimal robustness in the gap metric*, IEEE Transactions on Automatic Control **35**, 6 (1990), 673–686.
- [9] B. JESSEN AND H. TORNEHAVE, *Mean motions and zeros of almost periodic functions*, Acta Mathematica **77** (1945), 137–279.
- [10] K. M. MIKKOLA, *Infinite-dimensional linear systems, optimal control and algebraic Riccati equations*, Doctoral dissertation, Technical Report A452, Institute of Mathematics, Helsinki University of Technology, 2002.
- [11] N. K. NIKOLSKI, *Treatise on the shift operator*, Spectral function theory. With an appendix by S.V. Khrushchëv and V. V. Peller. Grundlehren der Mathematischen Wissenschaften 273, Springer-Verlag, Berlin, 1986.
- [12] N. K. NIKOLSKI, *Operators, functions, and systems: an easy reading. Volume 1*. Mathematical Surveys and Monographs, 92. American Mathematical Society, Providence, RI, 2002.
- [13] J. R. PARTINGTON, *Linear operators and linear systems. An analytical approach to control theory*, London Mathematical Society Student Texts 60, Cambridge University Press, Cambridge, 2004.

- [14] M. VIDYASAGAR, *The graph metric for unstable plants and robustness estimates for feedback stability*, IEEE Transactions on Automatic Control **29**, 5 (1984), 403–418.
- [15] G. VINNICOMBE, *Frequency domain uncertainty and the graph topology*, IEEE Transactions on Automatic Control **38**, 9 (1993), 1371–1383.
- [16] G. ZAMES AND A. K. EL-SAKKARY, *Unstable systems and feedback: The gap metric*, In Proceedings of the Allerton Conference, 380–385, Oct. 1980.

(Received December 5, 2010)

*Amol Sasane*  
*Department of Mathematics*  
*London School of Economics*  
*Houghton Street, London WC2A 2AE*  
*United Kingdom*  
*e-mail: sasane@lse.ac.uk*