

## CLASS A OPERATORS AND THEIR EXTENSIONS

SUNGEUN JUNG AND EUNGIL KO

(Communicated by R. Curto)

*Abstract.* In this paper, we study various properties of analytic extensions of class  $A$  operators. In particular, we show that every analytic extension of a class  $A$  operator has a scalar extension. As a corollary, we get that such an operator with rich spectrum has a nontrivial invariant subspace.

### 1. Introduction

Let  $\mathcal{H}$  and  $\mathcal{K}$  be separable complex Hilbert spaces and let  $\mathcal{L}(\mathcal{H}, \mathcal{K})$  denote the space of all bounded linear operators from  $\mathcal{H}$  to  $\mathcal{K}$ . If  $\mathcal{H} = \mathcal{K}$ , we write  $\mathcal{L}(\mathcal{H})$  in place of  $\mathcal{L}(\mathcal{H}, \mathcal{H})$ . If  $T \in \mathcal{L}(\mathcal{H})$ , we write  $\sigma(T)$ ,  $\sigma_{ap}(T)$ , and  $\sigma_e(T)$  for the spectrum, the approximate point spectrum, and the essential spectrum of  $T$ , respectively.

An arbitrary operator  $T \in \mathcal{L}(\mathcal{H})$  has a unique polar decomposition  $T = U|T|$ , where  $|T| = (T^*T)^{\frac{1}{2}}$  and  $U$  is the appropriate partial isometry satisfying  $\ker(U) = \ker(|T|) = \ker(T)$  and  $\ker(U^*) = \ker(T^*)$ . Associated with  $T$  is a related operator  $|T|^{\frac{1}{2}}U|T|^{\frac{1}{2}}$ , called the *Aluthge transform* of  $T$ , and denoted throughout this paper by  $\widehat{T}$ . For an arbitrary operator  $T \in \mathcal{L}(\mathcal{H})$ , the sequence  $\{\widehat{T}^{(n)}\}$  of Aluthge iterates of  $T$  is defined by  $\widehat{T}^{(0)} = T$  and  $\widehat{T}^{(n+1)} = \widehat{\widehat{T}^{(n)}}$  for every positive integer  $n$ .

An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be  $p$ -hyponormal if  $(T^*T)^p \geq (TT^*)^p$ . If  $p = 1$ ,  $T$  is called *hyponormal* and if  $p = \frac{1}{2}$ ,  $T$  is called *semi-hyponormal*. An operator  $T$  is said to be  $w$ -hyponormal if  $|\widehat{T}| \geq |T| \geq |\widehat{T}^*|$ .  $w$ -Hyponormal operators were introduced by Aluthge and Wang (see [2] and [3]). An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be class  $A$  if  $|T^2| - |T|^2 \geq 0$ , and  $T$  is said to be  $F$ -quasiclass  $A$  if  $F(T)^*(|T^2| - |T|^2)F(T) \geq 0$  for some function  $F$  that is analytic and nonconstant on some neighborhood of  $\sigma(T)$ . We say that an operator  $T \in \mathcal{L}(\mathcal{H})$  is  $p$ -quasiclass  $A$  if there exists a nonconstant polynomial  $p$  such that  $p(T)^*(|T^2| - |T|^2)p(T) \geq 0$ . In particular, if  $p(z) = z^k$  for some positive integer  $k$  or  $p(z) = z$ , then  $T$  is said to be a  $k$ -quasiclass  $A$  operator or a *quasiclass  $A$  operator*, respectively. The class of these operators has been studied by many authors (see [10], [13], [14], [23], and [27], etc.). An operator

*Mathematics subject classification* (2010): Primary 47A11, Secondary 47A15, 47B20.

*Keywords and phrases:* Class  $A$  operators, subscalar operators, invariant subspaces, Weyl's theorem.

This work was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science, and Technology (2009-0087565).

$T \in \mathcal{L}(\mathcal{H})$  is called *normaloid* if  $\|T\| = r(T)$  where  $r(T) := \sup\{|\lambda| : \lambda \in \sigma(T)\}$  denotes the spectral radius of  $T$ . It is well known from [10] that

$$p\text{-hyponormal} \Rightarrow w\text{-hyponormal} \Rightarrow \text{class A} \Rightarrow \text{normaloid}.$$

We give the following example to indicate that there exists a  $k$ -quasiclass A operators which does not belong to class A.

EXAMPLE 1.1. Let  $T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \in \mathcal{L}(\mathbb{C}^3)$ . Then  $|T^2| - |T|^2 \not\geq 0$ , and so  $T$  is

not a class A operator. However,  $T^{k*}(|T^2| - |T|^2)T^k = 0$  for every positive integer  $k$ , which implies that  $T$  is a  $k$ -quasiclass A operator for every positive integer  $k$ .

From the above example, it is natural to ask whether  $k$ -quasiclass A operators are normaloid or not. Next we give a  $k$ -quasiclass A operator which is not normaloid.

EXAMPLE 1.2. Let  $W_\alpha$  be the unilateral weighted shift with weights  $\alpha := \{\alpha_n\}_{n \geq 0}$  of positive real numbers. Then it is easy to compute that  $W_\alpha$  belongs to  $k$ -quasiclass A if and only if

$$\alpha_k \leq \alpha_{k+1} \leq \alpha_{k+2} \leq \dots$$

Hence, if we take the weights  $\alpha$  such that  $\alpha_0 = 2$  and  $\alpha_n = \frac{1}{2}$  for all  $n \geq 1$ , then  $W_\alpha$  belongs to  $k$ -quasiclass A for all  $k \in \mathbb{N}$ , but it is not normaloid.

We also find an equivalent condition for some operator-valued bilateral weighted shifts to be  $k$ -quasiclass A operators.

EXAMPLE 1.3. Let  $\mathcal{H} = \bigoplus_{n=-\infty}^{\infty} \mathcal{H}_n$  where  $\mathcal{H}_n = \mathcal{H}$  for all integers  $n$ . Given two positive operators  $A$  and  $B$  in  $\mathcal{L}(\mathcal{H})$ , define an operator  $T \in \mathcal{L}(\mathcal{H})$  by  $Tx = y$  with the following relation; if  $x = \bigoplus_{n=-\infty}^{\infty} x_n \in \mathcal{H}$ , then  $y = \bigoplus_{n=-\infty}^{\infty} y_n \in \mathcal{H}$  is given by

$$y_n = \begin{cases} Ax_{n-1} & \text{if } n \leq 1 \\ Bx_{n-1} & \text{if } n > 1. \end{cases}$$

By straightforward computations, we get that  $T$  is a  $k$ -quasiclass A operator if and only if

$$A^k[(AB^2A)^{\frac{1}{2}} - A^2]A^k \geq 0.$$

For instance, we shall provide an example by using the Maple program. Let  $A = \begin{pmatrix} 3 & -2 \\ -2 & 3 \end{pmatrix}$  and  $B = \begin{pmatrix} 2 & 0 \\ 0 & 2\sqrt{23} \end{pmatrix}$  be operators on  $\mathcal{H} = \mathbb{R}^2$ , and let  $\mathcal{H}_n = \mathcal{H}$  for all positive integers  $n$ . Note that

$$(AB^2A)^{\frac{1}{2}} - A^2 = \begin{pmatrix} 0.17472\dots & -3.1798\dots \\ -3.1798\dots & 11.770\dots \end{pmatrix}$$

as computed in [10]. Then

$$A^3[(AB^2A)^{\frac{1}{2}} - A^2]A^3 = \begin{pmatrix} 70778. \dots & -71500. \dots \\ -71500. \dots & 72227. \dots \end{pmatrix}$$

and its eigenvalues are  $143010. \dots$  and  $-1.1705 \dots$ , and so

$$A^3[(AB^2A)^{\frac{1}{2}} - A^2]A^3 \not\geq 0.$$

Therefore if we define  $T$  on  $\oplus_{n=-\infty}^{\infty} \mathcal{H}_n$  as in the above, then  $T$  is not a 3-quasiclass  $A$  operator.

An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be *analytic* if there exists a nonconstant analytic function  $F$  on a neighborhood of  $\sigma(T)$  such that  $F(T) = 0$ . We say that an operator  $T \in \mathcal{L}(\mathcal{H})$  is *algebraic* if there is a nonconstant polynomial  $p$  such that  $p(T) = 0$ . In particular, if  $T^k = 0$  for some positive integer  $k$ , then  $T$  is called *nilpotent*. An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be *quasinilpotent* if  $\sigma(T) = \{0\}$ . If an operator  $T \in \mathcal{L}(\mathcal{H})$  is analytic, then  $F(T) = 0$  for some nonconstant analytic function  $F$  on a neighborhood  $D$  of  $\sigma(T)$ . Since  $F$  cannot have infinitely many zeros in  $D$ , we write  $F(z) = G(z)p(z)$  where  $G$  is a function that is analytic and does not vanish on  $D$  and  $p$  is a nonconstant polynomial with zeros in  $D$ . By Riesz-Dunford calculus,  $G(T)$  is invertible and then  $p(T) = 0$ , which means that  $T$  is algebraic (see [5]). *When  $p$  has degree  $k$ , we say that  $T$  is analytic with order  $k$  throughout this paper.*

An operator  $T \in \mathcal{L}(\mathcal{H})$  is called *scalar* of order  $m$  if it possesses a spectral distribution of order  $m$ , i.e., if there is a continuous unital homomorphism of topological algebras

$$\Phi : C_0^m(\mathbb{C}) \rightarrow \mathcal{L}(\mathcal{H})$$

such that  $\Phi(z) = T$ , where as usual  $z$  stands for the identical function on  $\mathbb{C}$ , and  $C_0^m(\mathbb{C})$  for the space of all continuously differentiable functions of order  $m$  which are compactly supported,  $0 \leq m \leq \infty$ . An operator is *subscalar* of order  $m$  if it is similar to the restriction of a scalar operator of order  $m$  to an invariant subspace.

In 1984, M. Putinar showed in [25] that every hyponormal operator is subscalar of order 2. In 1987, his theorem was used to show that hyponormal operators with thick spectra have a nontrivial invariant subspace, which was a result due to S. Brown (see [4]). In this paper, we study various properties of analytic extensions of class  $A$  operators. In particular, we show that every analytic extension of a class  $A$  operator has a scalar extension. As a corollary, we get that such an operator with rich spectrum has a nontrivial invariant subspace. In addition, we study some properties of analytic extensions of class  $A$  operators.

## 2. Preliminaries

An operator  $T \in \mathcal{L}(\mathcal{H})$  is called *left semi-Fredholm* if  $T$  has closed range and  $\dim(\ker(T)) < \infty$ , and  $T$  is called *right semi-Fredholm* if  $T$  has closed range and  $\dim(\mathcal{H}/\text{ran}(T)) < \infty$ . When  $T$  is either left semi-Fredholm or right semi-Fredholm,  $T$

is called *semi-Fredholm*. In this case, the Fredholm index of  $T$  is defined by  $\text{ind}(T) := \dim(\ker(T)) - \dim(\mathcal{H}/\text{ran}(T))$ . Note that  $\text{ind}(T)$  is an integer or  $\pm\infty$ . We say that  $T$  is *Fredholm* if it is both left and right semi-Fredholm. Especially, an operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be *Weyl* if it is Fredholm of index zero. The Weyl spectrum is given by  $\sigma_w(T) = \{\lambda \in \mathbb{C} : T - \lambda \text{ is not Weyl}\}$  and we write  $\pi_{00}(T) := \{\lambda \in \text{iso}\sigma(T) : 0 < \dim(\ker(T - \lambda)) < \infty\}$ . We say that Weyl's theorem holds for  $T$  if  $\sigma(T) \setminus \sigma_w(T) = \pi_{00}(T)$ . A hole in  $\sigma_e(T)$  is a nonempty bounded component of  $\mathbb{C} \setminus \sigma_e(T)$ , and a pseudohole in  $\sigma_e(T)$  is a nonempty component of  $\sigma_e(T) \setminus \sigma_{le}(T)$  or of  $\sigma_e(T) \setminus \sigma_{re}(T)$ , where  $\sigma_{le}(T)$  and  $\sigma_{re}(T)$  denotes the left essential spectrum and the right essential spectrum of  $T$ , respectively. The *spectral picture* of  $T$  is the structure consisting of  $\sigma_e(T)$ , the collection of holes and pseudoholes in  $\sigma_e(T)$ , and it is denoted by  $SP(T)$  (see [24] for more details).

An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to have the *single-valued extension property* (or SVEP) if for every open subset  $G$  of  $\mathbb{C}$  and any analytic function  $f : G \rightarrow \mathcal{H}$  such that  $(T - z)f(z) \equiv 0$  on  $G$ , we have  $f(z) \equiv 0$  on  $G$ . For  $T \in \mathcal{L}(\mathcal{H})$  and  $x \in \mathcal{H}$ , the set  $\rho_T(x)$  is defined to consist of elements  $z_0$  in  $\mathbb{C}$  such that there exists an analytic function  $f(z)$  defined in a neighborhood of  $z_0$ , with values in  $\mathcal{H}$ , which verifies  $(T - z)f(z) \equiv x$ , and it is called *the local resolvent set of  $T$  at  $x$* . We denote the complement of  $\rho_T(x)$  by  $\sigma_T(x)$ , called *the local spectrum of  $T$  at  $x$* , and define *the local spectral subspace* of  $T$ ,  $H_T(F) = \{x \in \mathcal{H} : \sigma_T(x) \subseteq F\}$  for each subset  $F$  of  $\mathbb{C}$ . An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to have *property  $(\beta)$*  if for every open subset  $G$  of  $\mathbb{C}$  and every sequence  $f_n : G \rightarrow \mathcal{H}$  of  $\mathcal{H}$ -valued analytic functions such that  $(T - z)f_n(z)$  converges uniformly to 0 in norm on compact subsets of  $G$ , then  $f_n(z)$  converges uniformly to 0 in norm on compact subsets of  $G$ . An operator  $T \in \mathcal{L}(\mathcal{H})$  is said to have *Dunford's property (C)* if  $H_T(F)$  is closed for each closed subset  $F$  of  $\mathbb{C}$ . It is well known from [18] that

$$\text{Property } (\beta) \Rightarrow \text{Dunford's property (C)} \Rightarrow \text{SVEP.}$$

Let  $z$  be the coordinate function in the complex plane  $\mathbb{C}$  and  $d\mu(z)$  the planar Lebesgue measure. Consider a bounded (connected) open subset  $U$  of  $\mathbb{C}$ . We shall denote by  $L^2(U, \mathcal{H})$  the Hilbert space of measurable functions  $f : U \rightarrow \mathcal{H}$ , such that

$$\|f\|_{2,U} = \left( \int_U \|f(z)\|^2 d\mu(z) \right)^{\frac{1}{2}} < \infty.$$

The space of functions  $f \in L^2(U, \mathcal{H})$  that are analytic in  $U$  is denoted by

$$A^2(U, \mathcal{H}) = L^2(U, \mathcal{H}) \cap \mathcal{O}(U, \mathcal{H})$$

where  $\mathcal{O}(U, \mathcal{H})$  denotes the Fréchet space of  $\mathcal{H}$ -valued analytic functions on  $U$  with respect to uniform topology.  $A^2(U, \mathcal{H})$  is called the Bergman space for  $U$ . Note that  $A^2(U, \mathcal{H})$  is a Hilbert space.

Now, let us define a special Sobolev type space. For a fixed non-negative integer  $m$ , the vector-valued Sobolev space  $W^m(U, \mathcal{H})$  with respect to  $\bar{\partial}$  and of order  $m$  will be the space of those functions  $f \in L^2(U, \mathcal{H})$  whose derivatives  $\bar{\partial}f, \dots, \bar{\partial}^m f$  in the

sense of distributions still belong to  $L^2(U, \mathcal{H})$ . Endowed with the norm

$$\|f\|_{W^m}^2 = \sum_{i=0}^m \|\bar{\partial}^i f\|_{2,U}^2,$$

$W^m(U, \mathcal{H})$  becomes a Hilbert space contained continuously in  $L^2(U, \mathcal{H})$ .

We can easily show that the linear operator  $M$  of multiplication by  $z$  on  $W^m(U, \mathcal{H})$  is continuous and it has a spectral distribution  $\Phi$  of order  $m$  defined by the following relation; for  $\varphi \in C_0^m(\mathbb{C})$  and  $f \in W^m(U, \mathcal{H})$ ,  $\Phi(\varphi)f = \varphi f$ . Hence  $M$  is a scalar operator of order  $m$ .

### 3. Main results

In this section, we will show that every analytic extension of a class  $A$  operator has a scalar extension. For this, we begin with the following lemmas.

LEMMA 3.1. ([25]) For a bounded open disk  $D$  in the complex plane  $\mathbb{C}$  there is a constant  $C_D$  such that for any operator  $T \in \mathcal{L}(\mathcal{H})$  and  $f \in W^m(D, \mathcal{H})$  ( $m \geq 2$ ) we have

$$\|(I - P)\bar{\partial}^i f\|_{2,D} \leq C_D (\|(T - z)^* \bar{\partial}^{i+1} f\|_{2,D} + \|(T - z)^* \bar{\partial}^{i+2} f\|_{2,D})$$

for  $i = 0, 1, \dots, m - 2$ , where  $P$  denotes the orthogonal projection of  $L^2(D, \mathcal{H})$  onto the Bergman space  $A^2(D, \mathcal{H})$ .

LEMMA 3.2. ([25]) Let  $T \in \mathcal{L}(\mathcal{H})$  be a hyponormal operator and let  $D$  be a bounded disk in  $\mathbb{C}$ . If  $\{f_n\}$  is a sequence in  $W^m(D, \mathcal{H})$  ( $m > 2$ ) such that

$$\lim_{n \rightarrow \infty} \|(T - z)\bar{\partial}^i f_n\|_{2,D} = 0$$

for  $i = 1, 2, \dots, m$ , then  $\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n\|_{2,D_0} = 0$  for  $i = 1, 2, \dots, m - 2$  where  $D_0$  is a disk strictly contained in  $D$ .

LEMMA 3.3. Let  $D$  be a bounded disk in  $\mathbb{C}$  and let  $m$  be a positive integer with  $m > 12$ . If  $T \in \mathcal{L}(\mathcal{H})$  is a class  $A$  operator and  $f_n$  is a sequence in  $W^m(D, \mathcal{H})$  such that

$$\lim_{n \rightarrow \infty} \|(T - z)\bar{\partial}^i f_n\|_{2,D} = 0$$

for  $i = 1, 2, \dots, m$ , then it holds that

$$\lim_{n \rightarrow \infty} \|(I - P)\bar{\partial}^i f_n\|_{2,D_1} = 0$$

for  $i = 0, 1, 2, \dots, m - 12$ , where  $P$  denotes the orthogonal projection of  $L^2(D, \mathcal{H})$  onto  $A^2(D, \mathcal{H})$  and  $D_1$  is any disk relatively compact in  $D$ . Furthermore, we have

$$\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n\|_{2,D_2} = 0$$

for  $i = 1, 2, \dots, m - 12$ , where  $D_2$  is any disk relatively compact in  $D_1$ .

*Proof.* As in [14, Lemma 3.1], we can show that

$$\lim_{n \rightarrow \infty} \|(I - P)\bar{\partial}^i f_n\|_{2,D_1} = 0$$

for  $i = 0, 1, 2, \dots, m - 12$ , where  $P$  denotes the orthogonal projection of  $L^2(D, \mathcal{H})$  onto  $A^2(D, \mathcal{H})$  and  $D_1$  is any disk relatively compact in  $D$ . Then it follows that

$$\lim_{n \rightarrow \infty} \|(T - z)P\bar{\partial}^i f_n\|_{2,D_1} = 0$$

for  $i = 1, 2, \dots, m - 12$ . Since  $T$  has property  $(\beta)$  from [14], we get that

$$\lim_{n \rightarrow \infty} \|P\bar{\partial}^i f_n\|_{2,D_2} = 0$$

for  $i = 1, 2, \dots, m - 12$ , where  $D_2$  is any disk relatively compact in  $D_1$ . Hence we complete our proof.  $\square$

The next lemma is the key step to prove the subscalarity for analytic extensions of class  $A$  operators.

LEMMA 3.4. Let  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  be an analytic extension of a class  $A$  operator, i.e.,  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix}$  where  $T_1$  is a class  $A$  operator and  $T_3$  is analytic with order  $k$  and let  $D$  be a bounded disk in  $\mathbb{C}$  containing  $\sigma(T)$ . Define the map  $V : \mathcal{H} \oplus \mathcal{K} \rightarrow H(D)$  by

$$Vh = \widetilde{1} \otimes h \left( \equiv 1 \otimes h + \overline{(T - z)W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})} \right)$$

where

$$H(D) := W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K}) / \overline{(T - z)W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})}$$

and  $\widetilde{1} \otimes h$  denotes the constant function sending any  $z \in D$  to  $h$ . Then  $V$  is one-to-one and has closed range.

*Proof.* Let  $f_n = f_n^1 \oplus f_n^2 \in W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})$  and  $h_n = h_n^1 \oplus h_n^2 \in \mathcal{H} \oplus \mathcal{K}$  be sequences such that

$$\lim_{n \rightarrow \infty} \|(T - z)f_n + 1 \otimes h_n\|_{W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})} = 0. \tag{1}$$

Then from (1) we have the following equations:

$$\begin{cases} \lim_{n \rightarrow \infty} \|(T_1 - z)f_n^1 + T_2 f_n^2 + 1 \otimes h_n^1\|_{W^{2k+12}} = 0 \\ \lim_{n \rightarrow \infty} \|(T_3 - z)f_n^2 + 1 \otimes h_n^2\|_{W^{2k+12}} = 0. \end{cases} \tag{2}$$

By the definition of the norm for the Sobolev space, (2) implies that

$$\begin{cases} \lim_{n \rightarrow \infty} \|(T_1 - z)\bar{\partial}^i f_n^1 + T_2 \bar{\partial}^i f_n^2\|_{2,D} = 0 \\ \lim_{n \rightarrow \infty} \|(T_3 - z)\bar{\partial}^i f_n^2\|_{2,D} = 0 \end{cases} \tag{3}$$

for  $i = 1, 2, \dots, 2k + 12$ . Since  $T_3$  is analytic with order  $k$ , there exists a nonconstant analytic function  $F$  on a neighborhood of  $\sigma(T_3)$  such that  $F(T_3) = 0$ . As remarked in section one, write  $F(z) = G(z)p(z)$  where  $G$  is analytic and does not vanish on a neighborhood of  $\sigma(T_3)$  and  $p(z) = (z - z_1)(z - z_2) \cdots (z - z_k)$  is a polynomial of degree  $k$ . Set  $q_j(z) = (z - z_{j+1}) \cdots (z - z_k)$  for  $j = 0, 1, 2, \dots, k - 1$  and  $q_k(z) = 1$ .

*Claim.* It holds for every  $j = 0, 1, 2, \dots, k$  that

$$\lim_{n \rightarrow \infty} \|q_j(T_3) \bar{\partial}^i f_n^2\|_{2, D_j} = 0$$

for  $i = 1, 2, \dots, 2k - 2j + 12$ , where  $\sigma(T) \subsetneq D_k \subsetneq \cdots \subsetneq D_2 \subsetneq D_1 \subsetneq D$ .

To prove the claim, we will use the induction on  $j$ . Since  $0 = F(T_3) = G(T_3)p(T_3)$  and  $G(T_3)$  is invertible, it follows that  $q_0(T_3) = p(T_3) = 0$ , and so the claim holds when  $j = 0$ . Suppose that the claim is true for some  $j = r$  where  $0 \leq r < k$ . That is,

$$\lim_{n \rightarrow \infty} \|q_r(T_3) \bar{\partial}^i f_n^2\|_{2, D_r} = 0 \tag{4}$$

for  $i = 1, 2, \dots, 2k - 2r + 12$ , where  $\sigma(T) \subsetneq D_r \subsetneq \cdots \subsetneq D_1 \subsetneq D$ . By the second equation of (3) and (4), we get that

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \|q_{r+1}(T_3)(T_3 - z) \bar{\partial}^i f_n^2\|_{2, D_r} \\ &= \lim_{n \rightarrow \infty} \|q_{r+1}(T_3)(T_3 - z_{r+1} + z_{r+1} - z) \bar{\partial}^i f_n^2\|_{2, D_r} \\ &= \lim_{n \rightarrow \infty} \|(z_{r+1} - z)q_{r+1}(T_3) \bar{\partial}^i f_n^2\|_{2, D_r} \end{aligned} \tag{5}$$

for  $i = 1, 2, \dots, 2k - 2r + 12$ . Since  $z_{r+1}I$  is hyponormal, by applying Lemma 3.2 we obtain that

$$\lim_{n \rightarrow \infty} \|q_{r+1}(T_3) \bar{\partial}^i f_n^2\|_{2, D_{r+1}} = 0 \tag{6}$$

for  $i = 1, 2, \dots, 2k - 2r + 10$ , where  $\sigma(T) \subsetneq D_{r+1} \subsetneq D_r$ . Hence we complete the proof of our claim.

From the claim with  $j = k$ , we have

$$\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n^2\|_{2, D_k} = 0 \tag{7}$$

for  $i = 1, 2, \dots, 12$ , which implies by Lemma 3.1 that

$$\lim_{n \rightarrow \infty} \|(I - P_2) f_n^2\|_{2, D_k} = 0 \tag{8}$$

where  $P_2$  denotes the orthogonal projection of  $L^2(D_k, \mathcal{H})$  onto  $A^2(D_k, \mathcal{H})$ . By combining (7) with the first equation of (3), we obtain that

$$\lim_{n \rightarrow \infty} \|(T_1 - z) \bar{\partial}^i f_n^1\|_{2, D_k} = 0 \tag{9}$$

for  $i = 1, 2, \dots, 12$ . From Lemma 3.3, it follows that

$$\lim_{n \rightarrow \infty} \|(I - P_1)f_n^1\|_{2, D_{k,1}} = 0. \tag{10}$$

Set  $Pf_n := \begin{pmatrix} P_1 f_n^1 \\ P_2 f_n^2 \end{pmatrix}$ . Combining (8) and (10) with (2), we have

$$\lim_{n \rightarrow \infty} \|(T - z)Pf_n + 1 \otimes h_n\|_{2, D_{k,1}} = 0.$$

Let  $\Gamma$  be a curve in  $D_{k,1}$  surrounding  $\sigma(T)$ . Then

$$\lim_{n \rightarrow \infty} \|Pf_n(z) + (T - z)^{-1}(1 \otimes h_n)(z)\| = 0$$

uniformly for all  $z \in \Gamma$ . Applying Riesz-Dunford functional calculus, we obtain that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{2\pi i} \int_{\Gamma} Pf_n(z) dz + h_n \right\| = 0.$$

But by Cauchy’s theorem,  $\frac{1}{2\pi i} \int_{\Gamma} Pf_n(z) dz = 0$ . Hence  $\lim_{n \rightarrow \infty} \|h_n\| = 0$ , and so  $V$  is one-to-one and has closed range.  $\square$

Now we are ready to prove that every analytic extension of a class  $A$  operator has a scalar extension.

**THEOREM 3.5.** Every analytic extension of a class  $A$  operator is subscalar.

*Proof.* Let  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix}$  be an operator matrix defined on  $\mathcal{H} \oplus \mathcal{K}$ , where  $T_1$  is a class  $A$  operator and  $T_3$  is analytic with order  $k$ . Let  $D$  be an arbitrary bounded open disk in  $\mathbb{C}$  that contains  $\sigma(T)$ . As in Lemma 3.4, if we define an operator  $V : \mathcal{H} \oplus \mathcal{K} \rightarrow H(D)$  by  $Vh = \widetilde{1 \otimes h}$ , then  $V$  is one-to-one and has closed range. The class of a vector  $f$  or an operator  $S$  on  $H(D)$  will be denoted by  $\tilde{f}$ , respectively  $\tilde{S}$ . Let  $M$  be the operator of multiplication by  $z$  on  $W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})$ . Then  $M$  is a scalar operator of order  $2k + 12$  and has a spectral distribution  $\Phi$ . Since the range of  $T - z$  is invariant under  $M$ ,  $\tilde{M}$  can be well-defined. Moreover, consider the spectral distribution  $\Phi : C_0^{2k+12}(\mathbb{C}) \rightarrow \mathcal{L}(W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K}))$  defined by the following relation; for  $\varphi \in C_0^{2k+12}(\mathbb{C})$  and  $f \in W^{2k+12}(D, \mathcal{H}) \oplus W^{2k+12}(D, \mathcal{K})$ ,  $\Phi(\varphi)f = \varphi f$ . Then the spectral distribution  $\Phi$  of  $M$  commutes with  $T - z$ , and so  $\tilde{M}$  is still a scalar operator of order  $2k + 12$  with  $\tilde{\Phi}$  as a spectral distribution. Since

$$VTh = \widetilde{1 \otimes Th} = z \widetilde{\otimes h} = \tilde{M}(\widetilde{1 \otimes h}) = \tilde{M}Vh$$

for all  $h \in \mathcal{H} \oplus \mathcal{K}$ ,  $VT = \tilde{M}V$ . In particular,  $\text{ran}(V)$  is invariant under  $\tilde{M}$ , where  $\text{ran}(V)$  is the range of  $V$ . Since  $\text{ran}(V)$  is closed, it is a closed invariant subspace of the scalar operator  $\tilde{M}$ . Since  $T$  is similar to the restriction  $\tilde{M}|_{\text{ran}(V)}$  and  $\tilde{M}$  is a scalar operator of order  $2k + 12$ ,  $T$  is subscalar of order  $2k + 12$ .  $\square$

As an application of our main theorem, we prove that every  $F$ -quasiclass  $A$  operator is subscalar with the following lemma.

LEMMA 3.6. Let  $T \in \mathcal{L}(\mathcal{H})$  be  $F$ -quasiclass  $A$  and let  $\mathcal{M}$  be an invariant subspace for  $T$ . Then the restriction  $T|_{\mathcal{M}}$  is a  $p$ -quasiclass  $A$  operator.

*Proof.* Since  $T$  is an  $F$ -quasiclass  $A$  operator,  $F(T)^*(|T^2| - |T|^2)F(T) \geq 0$  for some function  $F$  analytic and nonconstant on a neighborhood of  $\sigma(T)$ . Set  $F(z) = G(z)p(z)$  where  $G$  is a nonvanishing analytic function on a neighborhood of  $\sigma(T)$  and  $p$  is a nonconstant polynomial. Since  $\mathcal{M}$  is a  $T$ -invariant subspace, we can write  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix}$  on the decomposition  $\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^\perp$ , where  $T_1 = T|_{\mathcal{M}}$ ,  $T_3 = (I - P)T(I - P)|_{\mathcal{M}^\perp}$ , and  $P$  denotes the orthogonal projection of  $\mathcal{H}$  onto  $\mathcal{M}$ . Since  $((T^2)^*T^2)^{\frac{1}{2}} \geq 0$ , from [9] we can set

$$|T^2| = ((T^2)^*T^2)^{\frac{1}{2}} = \begin{pmatrix} B & C \\ C^* & D \end{pmatrix},$$

where  $B \geq 0$ ,  $D \geq 0$ , and  $C = B^{\frac{1}{2}}SD^{\frac{1}{2}}$  for some contraction  $S : \mathcal{M}^\perp \rightarrow \mathcal{M}$ . Then a simple calculation gives that

$$(T^2)^*T^2 = |T^2|^2 = \begin{pmatrix} B & C \\ C^* & D \end{pmatrix}^2 = \begin{pmatrix} B^2 + CC^* & BC + CD \\ C^*B + DC^* & C^*C + D^2 \end{pmatrix}.$$

Since

$$(T^2)^*T^2 = \begin{pmatrix} (T_1^2)^*T_1^2 & * \\ * & * \end{pmatrix},$$

we get that  $B^2 + CC^* = (T_1^2)^*T_1^2$ . Hence

$$|T_1^2| = ((T_1^2)^*T_1^2)^{\frac{1}{2}} = (B^2 + CC^*)^{\frac{1}{2}} \geq B.$$

Also, since

$$|T|^2 = T^*T = \begin{pmatrix} T_1^*T_1 & * \\ * & * \end{pmatrix} = \begin{pmatrix} |T_1|^2 & * \\ * & * \end{pmatrix},$$

we have

$$\begin{aligned} 0 &\leq F(T)^*(|T^2| - |T|^2)F(T) \\ &= F(T)^* \begin{pmatrix} B - |T_1|^2 & * \\ * & * \end{pmatrix} F(T) = G(T)^* \begin{pmatrix} p(T_1)^*(B - |T_1|^2)p(T_1) & * \\ * & * \end{pmatrix} G(T) \end{aligned}$$

by Riesz-Dunford's functional calculus. Since  $G(T)$  is invertible, we obtain from [9] that  $p(T_1)^*(B - |T_1|^2)p(T_1) \geq 0$ , which completes our proof.  $\square$

THEOREM 3.7. Every  $F$ -quasiclass  $A$  operator is subscalar. In particular, every  $k$ -quasiclass  $A$  operator is subscalar of order  $2k + 12$ .

*Proof.* Suppose that  $T \in \mathcal{L}(\mathcal{H})$  satisfies that  $F(T)^*(|T^2| - |T|^2)F(T) \geq 0$  for some analytic function  $F$  on a neighborhood of  $\sigma(T)$ . If the range of  $F(T)$  is norm dense in  $\mathcal{H}$ , then  $T$  is a class  $A$  operator. Hence  $T$  is subscalar of order 12 by

Theorem 3.5. So it suffices to assume that the range of  $F(T)$  is not norm dense in  $\mathcal{H}$ . Since  $F(T)$  commutes with  $T$ ,  $\overline{\text{ran}(F(T))}$  is a  $T$ -invariant subspace, and so we can express  $T$  as  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix}$  on  $\mathcal{H} = \overline{\text{ran}(F(T))} \oplus \ker(F(T)^*)$  where  $T_1 = T|_{\overline{\text{ran}(F(T))}}$ ,  $T_3 = (I - P)T(I - P)|_{\ker(F(T)^*)}$ , and  $P$  denotes the projection of  $\mathcal{H}$  onto  $\overline{\text{ran}(F(T))}$ . Note that  $F(z) = G(z)p(z)$  where  $G$  is a nonvanishing analytic function on a neighborhood of  $\sigma(T)$  and  $p$  is a nonconstant polynomial. Then  $G(T)$  is invertible and thus we obtain that  $\ker(F(T)^*) = \ker(p(T)^*)$ . Since  $p(T_3) = (I - P)p(T)(I - P)|_{\ker(F(T)^*)}$ , it holds for any  $x \in \ker(F(T)^*)$  that

$$\langle p(T_3)x, x \rangle = \langle p(T)x, x \rangle = \langle x, p(T)^*x \rangle = 0.$$

Hence  $p(T_3) = 0$  and so  $T_3$  is analytic. In addition, since  $P(|T|^2 - |T|^2)P \geq 0$ , we have

$$|T_1^2| - |T_1|^2 \geq B - |T_1|^2 \geq 0$$

from the proof of Lemma 3.6 and [9]. This means that  $T_1$  is a class  $A$  operator. Therefore if  $T_3$  is analytic with order  $k$ , then  $T$  is subscalar of order  $2k + 12$  by Theorem 3.5.  $\square$

In the next corollary, we obtain a partial solution to the invariant subspace problem for analytic extensions of class  $A$  operators, which is a generalization of S. Brown’s result mentioned in section one.

COROLLARY 3.8. Let  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  be an analytic extension of a class  $A$  operator. If  $\sigma(T)$  has nonempty interior in  $\mathbb{C}$ , then  $T$  has a nontrivial invariant subspace.

*Proof.* The proof follows from Theorem 3.5 and [8].  $\square$

For the following corollary, note that an operator  $T \in \mathcal{L}(\mathcal{H})$  is said to be *power regular* if  $\{\|T^n x\|^{\frac{1}{n}}\}_{n=0}^\infty$  converges for each  $x \in \mathcal{H}$  and  $r_T(x)$  denotes the *local spectral radius* of  $T$  at  $x$  given by  $r_T(x) := \limsup_{n \rightarrow \infty} \|T^n x\|^{\frac{1}{n}}$ . Moreover, we recall that for an operator  $T \in \mathcal{L}(\mathcal{H})$ , a *spectral maximal space* of  $T$  is defined to be a closed  $T$ -invariant subspace  $\mathcal{M}$  of  $\mathcal{H}$  with the property that  $\mathcal{M}$  contains any closed  $T$ -invariant subspace  $\mathcal{N}$  of  $\mathcal{H}$  such that  $\sigma(T|_{\mathcal{N}}) \subset \sigma(T|_{\mathcal{M}})$ . Furthermore, recall that an operator  $X \in \mathcal{L}(\mathcal{H}, \mathcal{K})$  is called a *quasiaffinity* if it has trivial kernel and dense range. An operator  $S \in \mathcal{L}(\mathcal{H})$  is said to be a *quasiaffine transform* of an operator  $T \in \mathcal{L}(\mathcal{H})$  if there is a quasiaffinity  $X \in \mathcal{L}(\mathcal{H}, \mathcal{K})$  such that  $XS = TX$ . Also, operators  $S \in \mathcal{L}(\mathcal{H})$  and  $T \in \mathcal{L}(\mathcal{H})$  are *quasisimilar* if there are quasiaffinities  $X \in \mathcal{L}(\mathcal{H}, \mathcal{K})$  and  $Y \in \mathcal{L}(\mathcal{K}, \mathcal{H})$  such that  $XS = TX$  and  $SY = YT$ .

COROLLARY 3.9. If  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic extension of a class  $A$  operator, then the following statements hold.

- (i)  $T$  has property  $(\beta)$ , Dunford’s property  $(C)$ , and the single-valued extension property.

(ii)  $T$  is power regular.

(iii)  $r_T(x) = \lim_{n \rightarrow \infty} \|T^n x\|^{\frac{1}{n}}$  for all  $x \in \mathcal{H}$ .

(iv)  $H_T(E)$  is a spectral maximal space of  $T$  and  $\sigma(T|_{H_T(E)}) \subset \sigma(T) \cap E$  for any closed subset  $E$  in  $\mathbb{C}$ .

(v) If  $S$  is a quasilinear transform of  $T$  such that  $XS = TX$  where  $X$  is a quasilinearity, then  $S$  has the single-valued extension property and  $XH_S(E) \subseteq H_T(E)$  for any subset  $E$  in  $\mathbb{C}$ .

*Proof.* (i) From section two, it suffices to prove that  $T$  has property  $(\beta)$ . Since property  $(\beta)$  is transmitted from an operator to its restrictions to closed invariant subspaces, we are reduced by Theorem 3.5 to the case of a scalar operator. Since every scalar operator has property  $(\beta)$  (see [25]),  $T$  has property  $(\beta)$ .

(ii) From Theorem 3.5,  $T$  is similar to the restriction of a scalar operator to one of its invariant subspaces. Since a scalar operator is power regular and the restrictions of power regular operators to their invariant subspaces are still power regular,  $T$  is also power regular.

(iii) The proof follows from (i) and [18].

(iv) Since  $T$  has property (C) from (i),  $H_T(E)$  is closed for any closed subset  $E$  in  $\mathbb{C}$ . Hence the proof follows from [6] or [18].

(v) Let  $f : G \rightarrow \mathcal{H} \oplus \mathcal{H}$  be an analytic function on an open set  $G$  in  $\mathbb{C}$  such that  $(S - z)f(z) \equiv 0$ . Then  $(T - z)Xf(z) = X(S - z)f(z) \equiv 0$  on  $G$ . Since  $T$  has the single-valued extension property,  $Xf(z) \equiv 0$  on  $G$ . Since  $X$  is a quasilinearity,  $f(z) \equiv 0$  on  $G$ . Hence  $S$  has the single-valued extension property. To prove the last conclusion, it suffices to show that  $\sigma_T(Xx) \subseteq \sigma_S(x)$  for any  $x \in \mathcal{H} \oplus \mathcal{H}$ ; in fact, if it holds, then  $x \in H_S(E)$  implies  $\sigma_T(Xx) \subset E$ , which means that  $Xx \in H_T(E)$ . If  $z_0 \in \rho_S(x)$ , then we can choose an  $\mathcal{H} \oplus \mathcal{H}$ -valued analytic function  $f$  on some neighborhood of  $z_0$  for which  $(S - z)f(z) \equiv x$ . Since  $XS = TX$ , we have  $(T - z)Xf(z) = X(S - z)f(z) \equiv Xx$ , and so  $z_0 \in \rho_T(Xx)$ .  $\square$

**COROLLARY 3.10.** Let  $C$  and  $D$  be operator matrices in  $\mathcal{L}(\mathcal{H} \oplus \mathcal{H})$  which are analytic extensions of class A operators. If  $C$  and  $D$  are quasisimilar, then  $\sigma(C) = \sigma(D)$  and  $\sigma_e(C) = \sigma_e(D)$ .

*Proof.* Since  $C$  and  $D$  satisfy property  $(\beta)$  from Corollary 3.9, the proof follows from [26].  $\square$

**COROLLARY 3.11.** Let  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{H})$  be an analytic extension of a class A operator. If there exists a nonzero vector  $x \in \mathcal{H} \oplus \mathcal{H}$  such that  $\sigma_T(x) \subsetneq \sigma(T)$ , then  $T$  has a nontrivial hyperinvariant subspace.

*Proof.* Set  $\mathcal{M} := H_T(\sigma_T(x))$ , i.e.,  $\mathcal{M} = \{y \in \mathcal{H} \oplus \mathcal{H} : \sigma_T(y) \subseteq \sigma_T(x)\}$ . Since  $T$  has Dunford's property (C) by Corollary 3.9,  $\mathcal{M}$  is a  $T$ -hyperinvariant subspace from [6] or [18]. Since  $x \in \mathcal{M}$ , we get  $\mathcal{M} \neq \{0\}$ . Suppose  $\mathcal{M} = \mathcal{H} \oplus \mathcal{H}$ . Since  $T$  has the single-valued extension property by Corollary 3.9, it follows from [18] that

$$\sigma(T) = \bigcup \{ \sigma_T(y) : y \in \mathcal{H} \oplus \mathcal{H} \} \subseteq \sigma_T(x) \subsetneq \sigma(T),$$

which is a contradiction. Hence  $\mathcal{M}$  is a nontrivial  $T$ -hyperinvariant subspace.  $\square$

Next we show that every analytic extension  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  of a class  $A$  operator is isoloid (i.e.,  $\text{iso } \sigma(T) \subseteq \sigma_p(T)$  where  $\text{iso } \sigma(T)$  denotes the set of all isolated points of  $\sigma(T)$ ). If  $T \in \mathcal{L}(\mathcal{H})$  is analytic, then there exists a nonconstant polynomial  $p(z)$  such that  $p(T) = 0$ . If  $q(z)$  is a minimal polynomial satisfying  $q(T) = 0$ , it is obvious that  $q(z)$  is a factor of  $p(z)$ .

LEMMA 3.12. Suppose that  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic extension of a class  $A$  operator, i.e.,  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix}$  is an operator matrix on  $\mathcal{H} \oplus \mathcal{K}$  where  $T_1$  is a class  $A$  operator and  $F(T_3) = 0$  for a nonconstant analytic function  $F$  on a neighborhood  $D$  of  $\sigma(T_3)$ . Then the spectrum  $\sigma(T) = \sigma(T_1) \cup \sigma(T_3)$  and  $\sigma(T_3)$  is a subset of  $\{z \in \mathbb{C} : p(z) = 0\}$  where  $F(z) = G(z)p(z)$ ,  $G$  is analytic and does not vanish on  $D$ , and  $p$  is a polynomial.

*Proof.* Since  $p(T_3) = 0$ , choose a minimal polynomial  $q$  such that  $q(T_3) = 0$  and  $q(z)$  is a factor of  $p(z)$  as remarked in the above. Then  $\{z \in \mathbb{C} : q(z) = 0\}$  is nonempty and is contained in  $\{z \in \mathbb{C} : p(z) = 0\}$ . First we will show that  $\sigma(T_3) = \sigma_p(T_3) = \{z \in \mathbb{C} : q(z) = 0\}$ . Since  $q(T_3) = 0$ , we have  $q(\sigma(T_3)) = \sigma(q(T_3)) = \{0\}$  by the spectral mapping theorem. This means that  $\sigma(T_3) \subseteq \{z \in \mathbb{C} : q(z) = 0\}$ . Moreover if we assume that  $z_1, \dots, z_k$  are all the roots of  $q(z) = 0$ , not necessarily distinct, then  $(T_3 - z_1)(T_3 - z_2) \cdots (T_3 - z_k)x = 0$  for all  $x \in \mathcal{H}$ . By the minimality of the degree of  $q$ , we can select a vector  $x_0 \in \mathcal{H}$  such that  $(T_3 - z_2) \cdots (T_3 - z_k)x_0 \neq 0$ , and so  $z_1 \in \sigma_p(T_3)$ . Similarly,  $z_i \in \sigma_p(T_3)$  for all  $i = 1, 2, \dots, k$ . Hence  $\sigma(T_3) = \sigma_p(T_3) = \{z \in \mathbb{C} : q(z) = 0\}$ . Since  $\{z \in \mathbb{C} : q(z) = 0\}$  is a finite set,  $\sigma(T_1) \cap \sigma(T_3)$  is also finite, which implies that  $\sigma(T_1) \cap \sigma(T_3)$  has no interior point. By using [11], we get  $\sigma(T) = \sigma(T_1) \cup \sigma(T_3)$ , which completes the proof.  $\square$

THEOREM 3.13. Every analytic extension of a class  $A$  operator is isoloid.

*Proof.* Suppose that  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic extension of a class  $A$  operator. Then we get by Lemma 3.12 that  $\sigma(T) = \sigma(T_1) \cup \sigma(T_3)$  and  $\sigma(T_3)$  is a finite set. Let  $\lambda \in \mathbb{C}$  be an isolated point of  $\sigma(T)$ . Then either  $\lambda$  is an isolated point of  $\sigma(T_1)$  or  $\lambda \in \sigma(T_3)$ . If  $\lambda$  is an isolated point of  $\sigma(T_1)$ , then  $\lambda \in \sigma_p(T_1) \subseteq \sigma_p(T)$  because every class  $A$  operator is isoloid by [13]. Thus we may assume that  $\lambda \in \sigma_p(T_3)$  and  $\lambda \notin \sigma(T_1)$ . Since  $\lambda \in \sigma_p(T_3)$ , we get  $\ker(T_3 - \lambda) \neq \{0\}$ . In addition it holds for any  $x \in \ker(T_3 - \lambda)$  that  $(T - \lambda)(- (T_1 - \lambda)^{-1} T_2 x \oplus x) = 0$ . Hence  $\lambda \in \sigma_p(T)$ .  $\square$

COROLLARY 3.14. Let  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  be an analytic extension of a class  $A$  operator. If  $T$  is quasinilpotent, then it is nilpotent.

*Proof.* Since  $\sigma(T) = \{0\}$ , Lemma 3.12 implies that  $\sigma(T_1) = \{0\}$  and  $T_3$  is nilpotent. Since  $T_1$  is a class  $A$  operator, it is normaloid by [10]. Hence we get  $\|T_1\| = r(T_1) = 0$ . Therefore,  $T$  is nilpotent.  $\square$

PROPOSITION 3.15. Suppose that  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix} \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic

extension of a class  $A$  operator, i.e.,  $T_1$  is a class  $A$  operator and  $F(T_3) = 0$  for some nonconstant analytic function  $F$  on a neighborhood  $D$  of  $\sigma(T_3)$  with the representation  $F(z) = G(z)p(z)$  where  $G$  is analytic and does not vanish on  $D$  and  $p(z) = (z - z_1)(z - z_2) \cdots (z - z_k)$  is a polynomial. Then

- (i)  $H_T(E) \supset H_{T_1}(E) \oplus \{0\}$  for every subset  $E$  of  $\mathbb{C}$ , and
- (ii) if  $E$  is a closed subset of  $\mathbb{C}$  with  $z_i \notin E$  for some  $i = 1, 2, \dots, k$  and  $\{T_j\}_{j=1}^3$  are mutually commuting, then

$$H_T(E) \subseteq \{x_1 \oplus x_2 \in \mathcal{H} \oplus \mathcal{K} : p_i(T_3)x_1 \in H_{T_1}(E) \text{ and } x_2 \in \ker(p_i(T_3))\}$$

where  $p_i(z) = (z - z_1) \cdots (z - z_{i-1})(z - z_{i+1}) \cdots (z - z_k)$ .

*Proof.* (i) Let  $E$  be any subset of  $\mathbb{C}$  and let  $x_1 \in H_{T_1}(E)$  be given. Since  $T$  has the single-valued extension property by Corollary 3.9, there exists an  $\mathcal{H}$ -valued analytic function  $f_1$  on  $\mathbb{C} \setminus E$  for which  $(T_1 - z)f_1(z) \equiv x_1$  on  $\mathbb{C} \setminus E$ . Hence  $(T - z)(f_1(z) \oplus 0) \equiv x_1 \oplus 0$  on  $\mathbb{C} \setminus E$ , and so  $x_1 \oplus 0 \in H_T(E)$ .

(ii) We may assume that  $E$  is any closed subset of  $\mathbb{C}$  with  $z_1 \notin E$ , and let  $x_1 \oplus x_2 \in H_T(E)$  be given. Since  $T$  has the single-valued extension property by Corollary 3.9, we can choose an  $\mathcal{H} \oplus \mathcal{K}$ -valued analytic function  $f(z) = f_1(z) \oplus f_2(z)$  defined on  $\mathbb{C} \setminus E$  such that  $(T - z)f(z) = x_1 \oplus x_2$  for all  $z \in \mathbb{C} \setminus E$ . Then we have

$$\begin{cases} (T_1 - z)f_1(z) + T_2f_2(z) = x_1 \\ (T_3 - z)f_2(z) = x_2 \end{cases} \tag{11}$$

for all  $z \in \mathbb{C} \setminus E$ . Since  $p(T_3) = (T_3 - z_1)p_1(T_3) = 0$ , it follows from (11) that

$$(z - z_1)p_1(T_3)f_2(z) + p_1(T_3)x_2 \equiv 0 \text{ on } \mathbb{C} \setminus E. \tag{12}$$

By taking  $z = z_1$  in (12), we obtain that  $p_1(T_3)x_2 = 0$ , which means  $x_2 \in \ker(p_1(T_3))$ . Moreover,  $(T_1 - z)p_1(T_3)f_1(z) \equiv p_1(T_3)x_1$  on  $\mathbb{C} \setminus E$  from (11), which implies  $p_1(T_3)x_1 \in H_{T_1}(E)$ .  $\square$

In the following proposition, we will consider the Putnam’s type inequality corresponding to the analytic extension of a class  $A$  operator. Note that the Putnam’s inequality holds for class  $A$  operators;

$$|||T^2| - |T|^2| \leq \frac{1}{\pi} \mu(\sigma(T))$$

where  $\mu$  denotes the planar Lebesgue measure (see [23]).

PROPOSITION 3.16. Suppose that  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix} \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic

extension of a class  $A$  operator, i.e.,  $T_1$  is a class  $A$  operator and  $F(T_3) = 0$  for some nonconstant analytic function  $F$  on a neighborhood  $D$  of  $\sigma(T)$  with the representation

$F(z) = G(z)p(z)$  where  $G$  is analytic and does not vanish on  $D$  and  $p(z)$  is a polynomial.

(i) If  $T$  is compact, then both  $p(T)$  and  $F(T)$  are expressed as the sum of a normal operator and a nilpotent operator of order 2.

(ii) The following inequality holds;

$$||P(|T^2| - |T|^2)P|| \leq \frac{1}{\pi} \mu(\sigma(T))$$

where  $P$  is the orthogonal projection of  $\mathcal{H} \oplus \mathcal{H}$  onto  $\mathcal{H} \oplus \{0\}$ . Moreover, if  $\sigma(T)$  is a Lebesgue null set, then  $T_1$  is normal.

*Proof.* (i) We have  $F(T) = \begin{pmatrix} F(T_1) & S \\ 0 & 0 \end{pmatrix}$  for some operator  $S : \mathcal{H} \rightarrow \mathcal{H}$ . Since  $T$  is compact and  $T_1$  is the restriction of  $T$  to the invariant subspace  $\mathcal{H} \oplus \{0\}$ ,  $T_1$  is also compact. Thus  $T_1$  is normal by [14], and so is  $F(T_1)$ . Since  $F(T) - F(T_1) \oplus 0$  is a nilpotent operator of order 2, we complete the proof for  $F(T)$ , and the proof for  $p(T)$  is analogous.

(ii) Since  $PTP = TP$ , we get that  $|T_1^2| = (P|T^2|^2P)^{\frac{1}{2}} \geq P|T^2|P$  by Hansen’s inequality (see [10]). Since  $|T_1|^2 = (TP)^*(TP) = P|T|^2P$ , we have  $|T_1^2| - |T_1|^2 \geq P(|T^2| - |T|^2)P$ . Since  $\sigma(T) = \sigma(T_1) \cup \sigma(T_3)$  and  $\sigma(T_3)$  is a finite set by Lemma 3.12, it follows from [23] that

$$||P(|T^2| - |T|^2)P|| \leq |||T_1^2| - |T_1|^2|| \leq \frac{1}{\pi} \mu(\sigma(T_1)) = \frac{1}{\pi} \mu(\sigma(T)).$$

Moreover, if  $\mu(\sigma(T)) = 0$ , then  $\mu(\sigma(T_1)) = 0$ , and hence  $T_1$  is normal from [28].  $\square$

**COROLLARY 3.17.** Under the same hypotheses as in Proposition 3.16, let  $\sigma(T)$  be a Lebesgue null set. If  $T_1$  has dense range, then  $T$  is the direct sum of a normal operator and an analytic operator.

*Proof.* Since  $T_1$  is normal by Proposition 3.16, it suffices to show that  $T_2 = 0$ . Since  $\sigma(T)$  is a Lebesgue null set, we know that  $P(|T^2| - |T|^2)P = 0$  and  $|T_1^2| = |T_1|^2$  from Proposition 3.16. From easy computations, we get that

$$|T^2|^2 = \begin{pmatrix} |T_1^2|^2 & * \\ * & * \end{pmatrix} \quad \text{and} \quad |T|^4 = \begin{pmatrix} |T_1|^4 + T_1^* T_2 T_2^* T_1 & * \\ * & * \end{pmatrix}.$$

Hence  $|T_1^2|^2 = |T_1|^4 + T_1^* T_2 T_2^* T_1$ . Since  $|T_1^2| = |T_1|^2$ ,  $T_1^* T_2 T_2^* T_1 = 0$ . Since  $T_1$  has dense range,  $T_2 = 0$ . Thus  $T = T_1 \oplus T_3$ .  $\square$

Next we show that the spectral mapping theorem for the Weyl spectrum and Weyl’s theorem hold for an analytic extension  $T$  of a class  $A$  operator, more generally for  $f(T)$  where  $f$  is any analytic function on some neighborhood of  $\sigma(T)$ .

**THEOREM 3.18.** If  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{H})$  is an analytic extension of a class  $A$  operator, then

- (i) it satisfies Weyl’s theorem, and
- (ii)  $f(\sigma_w(T)) = \sigma_w(f(T))$  for any analytic function  $f$  on some neighborhood of  $\sigma(T)$ .

*Proof.* Suppose that  $T = \begin{pmatrix} T_1 & T_2 \\ 0 & T_3 \end{pmatrix} \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  is an analytic extension of a class A operator, i.e.,  $T_1$  is a class A operator and  $F(T_3) = 0$  for some nonconstant analytic function  $F$  on a neighborhood  $D$  of  $\sigma(T_3)$ .

(i) Note that every class A operator is isoloid and satisfies Weyl’s theorem by [5]. Furthermore, since every analytic operator is algebraic as noted in section one or [5],  $T_3$  is isoloid and it satisfies Weyl’s theorem by [22]. Since  $\sigma_w(T_1) \cap \sigma_w(T_3)$  has no interior points by Lemma 3.12, Weyl’s theorem holds for  $T_1 \oplus T_3$  from [20]. If  $\lambda_0 \notin \sigma_{le}(T_3) \cap \sigma_{re}(T_3)$  and  $\lambda_0 \in \sigma_e(T_3)$ , then  $T_3 - \lambda_0$  is semi-Fredholm and  $\lambda_0 \in \sigma(T_3)$ . Since  $T_3$  is algebraic,  $\lambda_0$  is an isolated point of  $\sigma(T_3)$ . By [7],  $T_3 - \lambda_0$  is Fredholm and  $\text{ind}(T_3 - \lambda_0) = 0$ , which is a contradiction. Thus we have  $\sigma_e(T_3) = \sigma_{le}(T_3) \cap \sigma_{re}(T_3)$ , which induces  $\sigma_e(T_3) = \sigma_{le}(T_3) = \sigma_{re}(T_3)$ . Therefore  $SP(T_3)$  has no pseudoholes, and so we finally get that Weyl’s theorem holds for  $T$  by [19].

(ii) If  $f$  is analytic on some neighborhood of  $\sigma(T)$ , then  $\sigma_w(f(T_1)) = f(\sigma_w(T_1))$  by [5]. Moreover since  $T_3$  is algebraic, we know that  $\sigma_w(f(T_3)) = f(\sigma_w(T_3))$  and  $\sigma_w(T_1) \cap \sigma_w(T_3)$  is finite and so has no interior points. Since  $\sigma_w(T_1) \cap \sigma_w(T_3)$  is finite,  $\sigma_w(f(T_1)) \cap \sigma_w(f(T_3)) = f(\sigma_w(T_1)) \cap f(\sigma_w(T_3))$  also has no interior points. Hence, we obtain from [20] that

$$\begin{aligned} \sigma_w(f(T)) &= \sigma_w(f(T_1)) \cup \sigma_w(f(T_3)) = f(\sigma_w(T_1)) \cup f(\sigma_w(T_3)) \\ &= f(\sigma_w(T_1) \cup \sigma_w(T_3)) = f(\sigma_w(T)). \end{aligned}$$

Thus we complete our proof.  $\square$

**COROLLARY 3.19.** Let  $T \in \mathcal{L}(\mathcal{H} \oplus \mathcal{K})$  be an analytic extension of a class A operator. Then Weyl’s theorem holds for  $f(T)$  where  $f$  is any analytic function on some neighborhood of  $\sigma(T)$ .

*Proof.* If  $T$  is an analytic extension of a class A operator, then  $T$  is isoloid by Theorem 3.13. Let  $f$  be an analytic function on some neighborhood of  $\sigma(T)$ . Then it follows from [21] that

$$\sigma(f(T)) \setminus \pi_{00}(f(T)) = f(\sigma(T) \setminus \pi_{00}(T)).$$

Since Weyl’s theorem holds for  $T$  and  $f(\sigma_w(T)) = \sigma_w(f(T))$  by Theorem 3.18,

$$\sigma(f(T)) \setminus \pi_{00}(f(T)) = f(\sigma(T) \setminus \pi_{00}(T)) = f(\sigma_w(T)) = \sigma_w(f(T)).$$

Accordingly, Weyl’s theorem holds for  $f(T)$ .  $\square$

REFERENCES

[1] A. ALUTHGE, *On p-hyponormal operators for  $0 < p < 1$* , Int. Eq. Op. Th. **13** (1990), 307–315.  
 [2] A. ALUTHGE AND D. WANG, *w-Hyponormal operators*, Int. Eq. Op. Th. **36** (2000), 1–10.  
 [3] A. ALUTHGE AND D. WANG, *w-Hyponormal operators II*, Int. Eq. Op. Th. **37** (2000), 324–331.  
 [4] S. BROWN, *Hyponormal operators with thick spectrum have invariant subspaces*, Ann. of Math. **125** (1987), 93–103.

- [5] X. CAO, *Analytically class A operators and Weyl's theorem*, J. Math. Anal. Appl. **320** (2006), 795–803.
- [6] I. COLOJOARĂ AND C. FOIAȘ, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
- [7] J. B. CONWAY, *A course in functional analysis*, Springer-Verlag New York, 1990.
- [8] J. ESCHMEIER, *Invariant subspaces for subscalar operators*, Arch. Math. **52** (1989), 562–570.
- [9] C. FOIAȘ AND A. E. FRAZHO, *The commutant lifting approach to interpolation problem*, Operator Theory Adv. Appl., vol. 44, Birkhäuser, Boston, 1990.
- [10] T. FURUTA, *Invitation to linear operators*, Taylor and Francis, 2001.
- [11] J. K. HAN, H. Y. LEE, AND W. Y. LEE, *Invertible completions of  $2 \times 2$  upper triangular operator matrices*, Proc. Amer. Math. Soc. **128** (1999), 119–123.
- [12] M. ITO AND T. YAMAZAKI, *Relations between two inequalities  $(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{p}{p+r}} \geq B^r$  and  $A^p \geq (A^{\frac{p}{2}}B^rA^{\frac{p}{2}})^{\frac{p}{p+r}}$  and their applications*, Int. Eq. Op. Th. **44** (2002), 442–450.
- [13] I. H. JEON AND B. P. DUGGAL, *On operators with an absolute value conditions*, J. Korean Math. Soc. **41** (2004), 617–627.
- [14] S. JUNG, E. KO, AND M. LEE, *On class A operators*, Studia Math. **198** (2010), 249–260.
- [15] I. B. JUNG, E. KO, AND C. PEARCY, *Aluthge transforms of operators*, Int. Eq. Op. Th. **37** (2000), 449–456.
- [16] I. H. KIM, *On  $(p, k)$ -quasihyponormal operators*, Math. Ineq. Appl. **7** (2004), 629–638.
- [17] E. KO,  *$K$ th roots of  $p$ -hyponormal operators are subscalar operators of order  $4k$* , Int. Eq. Op. Th. **59** (2007), 173–187.
- [18] K. LAURSEN AND M. NEUMANN, *An introduction to local spectral theory*, Clarendon Press, Oxford, 2000.
- [19] W. Y. LEE, *Weyl's theorem for operator matrices*, Int. Eq. Op. Th. **32** (1998), 319–331.
- [20] W. Y. LEE, *Weyl spectra of operator matrices*, Proc. Amer. Math. Soc. **129** (2000), 131–138.
- [21] W. Y. LEE AND S. H. LEE, *A spectral mapping theorem for the Weyl spectrum*, Glasgow Math. J. **38** (1996), 61–64.
- [22] M. OUDGHIRI, *Weyl's theorem and perturbations*, Int. Eq. Op. Th. **53** (2005), 535–545.
- [23] S. M. PATEL, M. CHO, K. TANAHASHI, AND A. UCHIYAMA, *Putnam's inequality for class A operators and an operator transform by Cho and Yamazaki*, Scientiae Mathematicae Japonicae **67** (2008), 393–402.
- [24] C. M. PEARCY, *Some recent developments in operator theory*, Conference board of the mathematical sciences regional conference series in mathematics, **36** (1975).
- [25] M. PUTINAR, *Hyponormal operators are subscalar operators*, J. Op. Th. **12** (1984), 385–395.
- [26] M. PUTINAR, *Quasimilarity of tuples with Bishop's property  $(\beta)$* , Int. Eq. Op. Th. **15** (1992), 1047–1052.
- [27] K. TANAHASHI, A. UCHIYAMA, AND M. CHO, *Isolated points of spectrum of  $(p, k)$ -quasihyponormal operators*, Linear Alge. Appl. **382** (2004), 221–229.
- [28] D. WANG AND J. I. LEE, *Spectral properties of class A operators*, Trends in Math, Information Center for Math. Science, **6** (2003), 93–98.

(Received May 12, 2011)

Sungeun Jung  
 Institute of Mathematical Sciences and Department of Mathematics  
 Ewha Womans University  
 Seoul, 120-750, Korea  
 e-mail: ssung105@ewhain.net

Eungil Ko  
 Department of Mathematics Ewha Womans University  
 Seoul, 120-750, Korea  
 e-mail: eiko@ewha.ac.kr