

SUBMAXIMAL OPERATOR SPACE STRUCTURES ON BANACH SPACES

VINOD KUMAR. P AND M. S. BALASUBRAMANI

(Communicated by Z.-J. Ruan)

Abstract. Subspaces of maximal operator spaces are called *submaximal* spaces and in general, they need not be maximal. We call those maximal operator spaces with the property that all submaximal spaces turn out to be maximal as *hereditarily maximal* spaces. Any two Banach isomorphic subspaces of a hereditarily maximal space will be completely isomorphic as operator spaces. We derive a characterization of these spaces. We introduce a notion of distance of an operator space to the class of submaximal spaces and discuss some related results.

1. Introduction

Just as one can treat C^* -algebras either concretely as closed $*$ -subalgebras of $\mathcal{B}(\mathcal{H})$ for some Hilbert space \mathcal{H} , or abstractly as a Banach algebra satisfying certain properties, operator spaces can also be considered in two different ways. A (*concrete*) *operator space* X is a closed linear subspace of $\mathcal{B}(\mathcal{H})$. Here, in each *matrix level* $M_n(X)$ (the space of all $n \times n$ matrices with entries from X), we have a norm $\|\cdot\|_n$, induced by the inclusion $M_n(X) \subset M_n(\mathcal{B}(\mathcal{H}))$, where the norm in $M_n(\mathcal{B}(\mathcal{H}))$ is given by the natural identification $M_n(\mathcal{B}(\mathcal{H})) \approx \mathcal{B}(\mathcal{H}^n)$. If X and Y are linear spaces and $\varphi : X \rightarrow Y$ is a linear map, $\varphi^{(n)} : M_n(X) \rightarrow M_n(Y)$, given by $[x_{ij}] \rightarrow [\varphi(x_{ij})]$, with $[x_{ij}] \in M_n(X)$ and $n \in \mathbb{N}$, determines a linear map from $M_n(X)$ to $M_n(Y)$. Suppose that each of the spaces $M_n(X)$ has a given norm $\|\cdot\|_n$, then the *complete bound norm* (in short *cb-norm*) of φ is defined as $\|\varphi\|_{cb} = \sup \left\{ \left\| \varphi^{(n)} \right\|; n \in \mathbb{N} \right\}$. φ is *completely bounded* if $\|\varphi\|_{cb} < \infty$. φ is a *complete isometry* if each map $\varphi^{(n)} : M_n(X) \rightarrow M_n(Y)$ is an isometry. If φ is a complete isometry, then $\|\varphi\|_{cb} = 1$. If $\|\varphi\|_{cb} \leq 1$, φ is said to be a complete contraction. If $\varphi : X \rightarrow Y$ is a completely bounded linear bijection and if its inverse is also completely bounded, then φ is said to be a *complete isomorphism*.

An *abstract operator space*, is a pair $(X, \{\|\cdot\|_n\}_{n \in \mathbb{N}})$ consisting of a linear space X and a complete norm $\|\cdot\|_n$ on $M_n(X)$ for every $n \in \mathbb{N}$, such that there exists a linear complete isometry $\varphi : X \rightarrow \mathcal{B}(\mathcal{H})$ for some Hilbert space \mathcal{H} . Two operator spaces are considered to be the same if there is a complete isometric isomorphism from X to Y . i.e., if there is a linear isomorphism $\varphi : X \rightarrow Y$ such that $\|\varphi\|_{cb} = \|\varphi^{-1}\|_{cb} = 1$.

Mathematics subject classification (2010): 46L07, 47L25.

Keywords and phrases: Operator spaces, maximal operator spaces, submaximal spaces, hereditarily maximal spaces.

This research work was supported by University Grants Commission of India, under Faculty Development Programme.

In that case, we write $X \approx Y$ *completely isometrically*. The sequence of matrix norms $\{\|\cdot\|_n\}_{n \in \mathbb{N}}$ is called an *operator space structure* on the linear space X . In 1988, Z.-J. Ruan [11], characterized the sequence of matrix norms on a linear space X that defines an operator space structure on X in terms of two properties of matrix norms. This allows us to view an operator space in an abstract way free of any concrete representation on a Hilbert space and so we no longer distinguish between concrete and abstract operator spaces. More information about operator spaces and completely bounded mappings may be found in [3], [8] or [9].

If X is a Banach space, any linear embedding of X into $B(\mathcal{H})$ for some Hilbert space \mathcal{H} , defines an operator space structure on X , by the natural identification $M_n(\mathcal{B}(\mathcal{H})) \approx \mathcal{B}(\mathcal{H}^n)$. If such an embedding preserves the original norm on X , (i.e., if the embedding is isometric), the corresponding operator space structure on X is said to be *admissible*. Blecher and Paulsen [2] observed that the set of all operator space structures admissible on a given Banach space X admits a minimal and maximal element. These structures were further investigated in [6] and [7]. Subspace structure of various maximal operator spaces were studied in [5].

Let X be a Banach space and $K = \text{Ball}(X^*)$ the closed unit ball of the dual space X^* of X , with its weak* topology. Then the canonical embedding $J : X \rightarrow C(K)$, defined by $J(x)(f) = f(x), x \in X$ and $f \in K$ is a linear isometry. Since by Gelfand-Naimark theorem [4], subspaces of C*-algebras are operator spaces, this identification of X induces matrix norms on $M_n(X)$ that makes X an operator space. The matrix norms on X are given by

$$\|[x_{ij}]\|_n = \sup\{\|[f(x_{ij})]\| : f \in K\}$$

for all $[x_{ij}] \in M_n(X)$ and for all $n \in \mathbb{N}$. The above defined operator space structure on X is called the *minimal operator space structure* on X , and we denote this operator space as $\text{Min}(X)$. For $[x_{ij}] \in M_n(X)$, we write $\|[x_{ij}]\|_{\text{Min}(X)}$ to denote its norm as an element of $M_n(\text{Min}(X))$. This minimal quantization of a normed space is characterized by the property that for any arbitrary operator space Y any bounded linear map $\varphi : Y \rightarrow \text{Min}(X)$ is completely bounded and satisfies

$$\|\varphi : Y \rightarrow \text{Min}(X)\|_{cb} = \|\varphi : Y \rightarrow X\|.$$

Thus, if X and Y are Banach spaces and $\varphi \in B(X, Y)$, then φ is completely bounded and $\|\varphi\|_{cb} = \|\varphi\|$, when considered as a map from $X \rightarrow \text{Min}(Y)$. It may be noted that $\text{Min}(X)$ is the smallest operator space structure on X . An operator space X is minimal if $\text{Min}(X) = X$. Also, an operator space is minimal if and only if it is completely isometric to a subspace of a commutative C*-algebra.

If X is a Banach space, there is a maximal way to consider it as an operator space. The matrix norms given by

$$\|[x_{ij}]\|_n = \sup\{\|[\varphi(x_{ij})]\|; \varphi \in \text{Ball}(B(X, Y))\}$$

where the supremum is taken over all operator spaces Y and all linear maps $\varphi \in \text{Ball}(B(X, Y))$, makes X an operator space. We denote this operator space as $\text{Max}(X)$

and is called the *maximal operator space structure* on X . For $[x_{ij}] \in M_n(X)$, we write $\|[x_{ij}]\|_{Max(X)}$ to denote its norm as an element of $M_n(Max(X))$. We say that an operator space X is maximal if $Max(X) = X$. By Ruan’s theorem [11], we also have

$$\|[x_{ij}]\|_{Max(X)} = \sup\{\|[\varphi(x_{ij})]\|; \varphi \in Ball(B(X, B(\mathcal{H})))\}$$

where the supremum is taken over all Hilbert spaces \mathcal{H} and all linear maps $\varphi \in Ball(B(X, B(\mathcal{H})))$. By the definition of $Max(X)$, any operator space structure that we can put on X , will be smaller than $Max(X)$. This maximal quantization of a Banach space is characterized by the property that for any arbitrary operator space Y any bounded linear map $\varphi : Max(X) \rightarrow Y$ is completely bounded and satisfies

$$\|\varphi : Max(X) \rightarrow Y\|_{cb} = \|\varphi : X \rightarrow Y\|.$$

Thus, if X and Y are Banach spaces and $\varphi \in B(X, Y)$, then φ is completely bounded and $\|\varphi\|_{cb} = \|\varphi\|$, when considered as a map from $Max(X) \rightarrow Y$. If X is any operator space, then the identity map on X defines completely contractive maps $Max(X) \rightarrow X \rightarrow Min(X)$. From the above discussions, we have the following observation.

PROPOSITION 1.1. *Let X and Y be operator spaces and $\varphi : X \rightarrow Y$ a bounded linear map, then φ is completely bounded and $\|\varphi\|_{cb} = \|\varphi\|$ if X is given the maximal operator space structure or Y is given the minimal operator space structure.*

Any subspace of a minimal operator space is again minimal, but quotient of a minimal space need not be minimal. Quotients of minimal operator spaces are called Q -spaces. Also, the class of Q -spaces is stable under taking quotients and subspaces. The subspace of a maximal space need not be maximal and such spaces are called *submaximal* spaces [5]. But quotient spaces inherits maximality.

THEOREM 1.2. ([9]) *If X is a maximal operator space and Y a closed subspace then $Max(X/Y) \approx Max(X)/Y$ completely isometrically.*

An operator space Z is *injective* [3] if for any operator spaces X and Y where Y contains X as a closed subspace, and for any completely bounded linear map $\varphi : X \rightarrow Z$, there exists a completely bounded extension $\tilde{\varphi} : Y \rightarrow Z$ such that $\tilde{\varphi}|_X = \varphi$ and $\|\tilde{\varphi}\|_{cb} = \|\varphi\|_{cb}$. An important class of operator spaces are those $X \subset B(H)$ which are isomorphic (as a Banach space) to a Hilbert space. We call such spaces as *Hilbertian* operator spaces. The spaces $Min(\ell^2)$ and $Max(\ell^2)$ are Hilbertian operator spaces. Blecher and Paulsen [2] observed that any separable infinite dimensional Hilbertian operator space lies between $Min(\ell^2)$ and $Max(\ell^2)$.

We discuss submaximal spaces and some of their properties in the next section. In section 3, we introduce *hereditarily maximal* spaces and give a characterization of these spaces. In section 4, we define the distance $d_{sm}(X)$ of a space to the class of submaximal spaces and make use of this to obtain a necessary and sufficient condition for a space to be submaximal. Also, we compute $d_{sm}(X)$ for some spaces and explore its relation with $d_Q(X)$, the distance of a space X to the class of Q -spaces.

2. Submaximal spaces

We have noted that subspace of a maximal space need not be maximal, i.e., if Y is a subspace of X and if $x_{ij} \in Y$ for $i, j = 1, 2, \dots, n$, then the norm of $[x_{ij}]$ in $M_n(\text{Max}(Y))$ can be larger than the norm of $[x_{ij}]$ as an element of $M_n(\text{Max}(X))$. However, we have the following result.

THEOREM 2.1. ([7]) *Let $x_{ij} \in X$, $i, j = 1, 2, \dots, n$, then $\|[x_{ij}]\|_{\text{Max}(X)} = \inf\{\|[x_{ij}]\|_{\text{Max}(Y)}; x_{ij} \in Y \text{ and } Y \subset X, \text{ finite dimensional}\}$.*

Also, if every subspace of $X = \text{Max}(X)$ is maximal, then by proposition 1.1, any two Banach isomorphic subspaces of X will be completely isomorphic as operator spaces.

DEFINITION 2.2. ([9]) An operator space X is said to be submaximal if it embeds completely isometrically into a maximal operator space Y .

Paulsen ([6]) observed that every separable submaximal space embeds completely isometrically into a separable maximal space. We now observe that just like every operator space embeds completely isometrically into $B(\mathcal{H})$ for some Hilbert space \mathcal{H} , every submaximal space embeds completely isometrically into $\text{Max}(B(\mathcal{H}))$ for some Hilbert space \mathcal{H} .

PROPOSITION 2.3. *Let X be a submaximal space of a maximal operator space $Y = \text{Max}(Y)$. If $\iota : X \rightarrow B(\mathcal{H})$ is a completely isometric inclusion, then ι defines a completely isometric inclusion $\tilde{\iota} : X \rightarrow \text{Max}(B(\mathcal{H}))$ and there is a completely contractive extension $\varphi : Y \rightarrow \text{Max}(B(\mathcal{H}))$.*

Proof. Let $\iota : X \rightarrow B(\mathcal{H})$ be a complete isometric inclusion. Since $X \hookrightarrow Y = \text{Max}(Y)$ and since $B(\mathcal{H})$ is injective, $\iota : X \rightarrow B(\mathcal{H})$ extends to a complete contraction $\varphi : Y \rightarrow B(\mathcal{H})$. The maximal operator space structure of Y implies that

$$\|\varphi : Y \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} = \|\varphi : Y \rightarrow \text{Max}(B(\mathcal{H}))\| \leq \|\varphi : Y \rightarrow B(\mathcal{H})\|_{cb} \leq 1$$

Thus, $\|\tilde{\iota} = \varphi|_X : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \leq 1$. If $\tilde{X} = \tilde{\iota}(X)$, then $\|\tilde{\iota}^{-1} : \tilde{X} \rightarrow X\|_{cb} \leq 1$, so that $\tilde{\iota}$ is a complete isometric inclusion of X into $\text{Max}(B(\mathcal{H}))$. \square

Using the above result, we now derive a characterization of submaximal spaces.

THEOREM 2.4. *An operator space X is a submaximal space if and only if any complete contraction $u : X \rightarrow B(\mathcal{H})$ defines a complete contraction $\tilde{u} = u : X \rightarrow \text{Max}(B(\mathcal{H}))$.*

Proof. Assume that X is a submaximal space and let $X \subset \text{Max}(Y)$. Let $u : X \rightarrow B(\mathcal{H})$ be a complete contraction. As in proposition 2.3, there exists a complete contractive extension $\phi : \text{Max}(Y) \rightarrow \text{Max}(B(\mathcal{H}))$ such that $\|\tilde{u} = \phi|_X : X \rightarrow$

$Max(B(\mathcal{H}))\|_{cb} \leq \|\phi : Max(Y) \rightarrow Max(B(\mathcal{H}))\|_{cb} \leq 1$. Conversely assume that $\iota : X \rightarrow B(\mathcal{H})$ be a complete isometric inclusion of X in $B(\mathcal{H})$, then by assumption, $\tilde{\iota} = \iota : X \rightarrow Max(B(\mathcal{H}))$ is a complete contraction. Also, $\|\tilde{\iota}^{-1} : \tilde{\iota}(X) \rightarrow X\|_{cb} \leq 1$, so that $\tilde{\iota}$ defines a complete isometric inclusion of X in $Max(B(\mathcal{H}))$, so that X is completely isometrically isomorphic to the submaximal space $\tilde{\iota}(X) \subset Max(B(\mathcal{H}))$. \square

Now we identify the completely bounded maps between two submaximal spaces.

THEOREM 2.5. *Let $X \subset Max(B(\mathcal{H}_1))$ and $Y \subset Max(B(\mathcal{H}_2))$ be submaximal spaces. Then any completely bounded map $\varphi : X \rightarrow Y$ extends to a bounded linear map from $B(\mathcal{H}_1)$ to $B(\mathcal{H}_2)$. Also, any bounded linear map $\varphi : B(\mathcal{H}_1) \rightarrow B(\mathcal{H}_2)$ with $\varphi(X) \subset Y$ induces a completely bounded map from X to Y .*

Proof. Let $\varphi : X \rightarrow Y$ be completely bounded. It is known from proposition 2.3 that $\tilde{\varphi} = \varphi : X \rightarrow Max(B(\mathcal{H}_2))$ is also completely bounded and $\|\tilde{\varphi}\|_{cb} \leq \|\varphi\|_{cb}$. Since $B(\mathcal{H}_2)$ is injective, we have an extension $\phi : Max(B(\mathcal{H}_1)) \rightarrow Max(B(\mathcal{H}_2))$ such that $\phi|_X = \tilde{\varphi}$ and $\|\phi\|_{cb} = \|\tilde{\varphi}\|_{cb}$. Thus, $\|\phi\| \leq \|\phi\|_{cb} = \|\tilde{\varphi}\|_{cb} \leq \|\varphi\|_{cb} < \infty$. This implies that ϕ is a bounded linear map from $B(\mathcal{H}_1)$ to $B(\mathcal{H}_2)$. Now let $\varphi : B(\mathcal{H}_1) \rightarrow B(\mathcal{H}_2)$ be a bounded linear map such that $\varphi(X) \subset Y$. Regarding φ as a map $\tilde{\varphi} = \varphi : Max(B(\mathcal{H}_1)) \rightarrow Max(B(\mathcal{H}_2))$, we see that $\tilde{\varphi}$ is completely bounded and $\|\tilde{\varphi}\|_{cb} = \|\varphi : B(\mathcal{H}_1) \rightarrow B(\mathcal{H}_2)\|$, so that $\|\tilde{\varphi}|_X : X \rightarrow Y\|_{cb} \leq \|\tilde{\varphi}\|_{cb} = \|\varphi\| < \infty$. \square

REMARK 2.6. The class of submaximal spaces is closed under taking closed subspaces and quotients. To see this, if X is a submaximal space, it embeds completely isometrically into a maximal operator space Y ; say $\varphi : X \hookrightarrow Max(Y)$. If Z is a closed subspace of X , then the restriction of φ to Z will be a complete isometric embedding of Z in $Max(Y)$. Also, in this case, $X/Z \subset Max(Y)/Z$ and from theorem 1.2, we have $Max(Y/Z) \approx Max(Y)/Z$ completely isometrically. This shows that the quotient space X/Z is a submaximal space.

Now we show that submaximal spaces are stable under ℓ^1 -sums.

THEOREM 2.7. *Let $\{X_i\}_{i \in I}$ be a family of submaximal operator spaces. Then $\ell^1(\{X_i; i \in I\})$ is submaximal.*

Proof. First, we note that ℓ^1 -sum of maximal spaces is again maximal. Let X_i 's be maximal, so that $Max(X_i) = X_i$ for all $i \in I$. Let $u : \ell^1(\{X_i; i \in I\}) \rightarrow B(H)$ be a bounded linear map. Corresponding to this u , there exists bounded linear maps $u_i : X_i \rightarrow B(H)$ for every $i \in I$. Then $\|u\| = \sup\{\|u_i\|; i \in I\}$. But by the definition of operator space structure on $\ell^1(\{X_i; i \in I\})$, and using the fact that X_i has maximal operator space structure, we see that $\|u\|_{cb} = \sup\{\|u_i\|_{cb}; i \in I\} = \sup\{\|u_i\|; i \in I\} = \|u\|$. This shows that $\ell^1(\{X_i; i \in I\})$ is maximal.

If X_i 's are submaximal, we have $X_i \subset Max(Y_i)$, for every $i \in I$, so that $\ell^1(\{X_i; i \in I\}) \subset \ell^1(\{Max(Y_i); i \in I\})$. Hence $\ell^1(\{X_i; i \in I\})$ is a submaximal space. \square

Now we show that submaximality of a space will be reflected in its bidual also.

THEOREM 2.8. *An operator space X is submaximal if and only if its bidual X^{**} is submaximal.*

Proof. Let $X \subset \text{Max}(Y)$. Then $X^{**} \subset (\text{Max}(Y))^{**}$. But from the duality relations [1], $(\text{Max}(Y))^* = \text{Min}(Y^*)$, so that $(\text{Max}(Y))^{**} = (\text{Min}(Y^*))^* = \text{Max}(Y^{**})$. Thus X^{**} is submaximal. The converse part follows from the fact that $X \subset X^{**}$ and from the remark 2.6. \square

3. Hereditarily maximal spaces

It is known that submaximal operator space structure on a Banach space X need not equals with the maximal operator space structure on X . But we show that in some spaces this will happen. We call such spaces as *hereditarily maximal*. (This name was suggested by Prof. Gilles Pisier in a private communication with us.)

DEFINITION 3.1. A maximal operator space X is said to be hereditarily maximal if any closed subspace of X is again a maximal operator space. i.e., for any closed subspace $Y \subset X = \text{Max}(X)$, we have $\text{Max}(Y) = Y$.

Note that any two closed subspaces of a hereditarily maximal space which are isomorphic as Banach spaces, are in fact completely isomorphic as operator spaces.

Now we derive a characterization of these spaces.

THEOREM 3.2. *A maximal operator space X is hereditarily maximal if and only if the space X has the following extension property: For any closed subspace $Y \subset X$, and for any bounded linear map $\varphi : Y \rightarrow B(\mathcal{H})$, there exists a bounded extension $\tilde{\varphi} : X \rightarrow B(\mathcal{H})$ such that $\|\tilde{\varphi}\| = \|\varphi\|$.*

Proof. Assume that X is hereditarily maximal. Let $Y \subset X$ be a closed subspace. Then $\text{Max}(Y) = Y$. Let $\varphi : Y \rightarrow B(\mathcal{H})$ be a bounded linear map. Since $Y = \text{Max}(Y)$, we see that φ is completely bounded and $\|\varphi\|_{cb} = \|\varphi\|$. Since $B(\mathcal{H})$ is injective, there exists a mapping $\tilde{\varphi} : X \rightarrow B(\mathcal{H})$ such that $\tilde{\varphi}|_Y = \varphi$ and $\|\tilde{\varphi}\|_{cb} = \|\varphi\|_{cb}$. Since X has maximal structure, $\|\tilde{\varphi}\|_{cb} = \|\tilde{\varphi}\|$. Thus $\|\tilde{\varphi}\| = \|\tilde{\varphi}\|_{cb} = \|\varphi\|_{cb} = \|\varphi\|$.

Conversely, assume that X has the above described extension property. We have to show that X is hereditarily maximal. Let $Y \subset X = \text{Max}(X)$. Let $u : Y \rightarrow \text{Max}(Y) \subset B(\mathcal{H})$ be the isometric inclusion mapping. Then u is bounded. By assumption it has an extension $\tilde{u} : X \rightarrow B(\mathcal{H})$ such that $\tilde{u}|_Y = u$ and $\|\tilde{u}\| = \|u\|$. Then $\|u\|_{cb} = \|\tilde{u}|_Y\|_{cb} \leq \|\tilde{u}\|_{cb} = \|\tilde{u}\| = \|u\|$. The second last equality follows from the fact that $X = \text{Max}(X)$. This shows that $\|u\|_{cb} = \|u\|$. Thus the formal isometric inclusion mapping of Y to $\text{Max}(Y)$ is a complete isometry, which implies that $\text{Max}(Y) = Y$. \square

The following theorem will serve us some examples for these type of spaces.

THEOREM 3.3. *If X is a Hilbertian operator space and Y be a closed subspace of $\text{Max}(X)$, then $Y = \text{Max}(Y)$.*

Proof. Let $p : X \rightarrow Y$ be the orthogonal projection. Then $p : \text{Max}(X) \rightarrow \text{Max}(Y)$ is a complete contraction. Note that the inclusion map $j : Y \hookrightarrow \text{Max}(X)$ is completely bounded. Therefore, $pj : Y \rightarrow \text{Max}(Y)$ is completely contractive. Thus $\text{Max}(Y) = Y$. \square

REMARK 3.4. The above theorem shows that every maximal Hilbertian operator space is hereditarily maximal. This can also be obtained by noting that a Hilbertian operator space X has the bounded extension property described in theorem 3.2. Let Y be a closed subspace of X and let u be any bounded linear map $u : Y \rightarrow B(\mathcal{H})$. Then Y admits an orthogonal projection $p : X \rightarrow Y$. Now the composition map $\tilde{u} = u \circ p : X \rightarrow B(\mathcal{H})$ is bounded and $\tilde{u}|_Y = u$. Also, $\|\tilde{u}\| = \|u \circ p\| \leq \|u\|$. Since \tilde{u} is an extension of u , this shows that $\|u\| = \|\tilde{u}\|$.

4. Distance to the class of submaximal spaces

We now introduce a notion of distance of an operator space to the class of submaximal spaces, and by using it we derive a characterization of submaximal spaces. Note that completely bounded Banach-Mazur distance between two operator spaces X and Y is defined as $d_{cb}(X, Y) = \inf\{\|\varphi\|_{cb} \|\varphi^{-1}\|_{cb} : \varphi : X \rightarrow Y \text{ a complete isomorphism}\}$.

DEFINITION 4.1. For a given operator space X , we define $d_{sm}(X) = \inf\{d_{cb}(X, Y); Y \text{ a submaximal space}\}$

We make use of this concept to characterize a submaximal space up to complete isometric isomorphism.

THEOREM 4.2. *An operator space X is a submaximal space, up to complete isometric isomorphism, if and only if $d_{sm}(X) = 1$.*

Proof. If X is a submaximal space, then clearly $d_{sm}(X) = 1$. For proving the other direction we use the fact that submaximal spaces are stable under ultra products. To see this, note if $X_i \subset \text{Max}(Y_i)$, and if \mathcal{U} is a nontrivial ultra filter on the indexing set I , then $\prod X_i / \mathcal{U} \subset \prod \text{Max}(Y_i) / \mathcal{U}$, which is completely isometrically isomorphic to $\text{Max}(\prod Y_i / \mathcal{U})$ [5]. If $d_{sm}(X) = 1$, then for any $n \in \mathbb{N}$, there exists a submaximal space X_n such that $d_{cb}(X, X_n) < 1 + 1/n$, and a complete isomorphism $u_n : X \rightarrow X_n$ such that $\|u_n\|_{cb} < 1 + 1/n$ and $\|u_n^{-1}\|_{cb} = 1$. Choose a nontrivial ultra filter \mathcal{U} on \mathbb{N} such that $\lim_{\mathcal{U}} \|u_n\|_{cb} = 1$. Then the ultra product $\prod X_n / \mathcal{U}$ is a submaximal space and $u : X \rightarrow \prod X_n / \mathcal{U}$ defined by $u = (u_n)_{n \in \mathbb{N}}$ is a complete isometric isomorphism. This shows that X is a submaximal space. \square

Operator spaces X and Y are said to be *C-completely isomorphic* if there exists a linear isomorphism $\varphi : X \rightarrow Y$ such that $\|\varphi\|_{cb} \|\varphi^{-1}\|_{cb} \leq C$. The following result gives another characterization of submaximal spaces upto *C*-complete isomorphism. This result is implicitly contained in [5].

THEOREM 4.3. *Let X be an operator space and $C > 0$. Then X is C -completely isomorphic to a submaximal space Y if and only if for every complete contraction $u : X \rightarrow B(\mathcal{H})$, we have $\|u : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \leq C$*

Proof. Assume that X is C -completely isomorphic to a submaximal space Y . Let $\varphi : X \rightarrow Y$ be such that $\|\varphi\|_{cb} \|\varphi^{-1}\|_{cb} \leq C$. Since Y is submaximal, $Y \subset \text{Max}(Z)$, for some operator space Z . Let $u : X \rightarrow B(\mathcal{H})$ be a complete contraction. Then the map $v = u\varphi^{-1} : Y \rightarrow B(\mathcal{H})$ is completely bounded. As in proposition 2.3, there exists a completely bounded extension $\tilde{v} : \text{Max}(Z) \rightarrow \text{Max}(B(\mathcal{H}))$ such that $\tilde{v}|_Y = u\varphi^{-1} : Y \rightarrow \text{Max}(B(\mathcal{H}))$ is completely bounded. Therefore,

$$\begin{aligned} \|u : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} &= \|u\varphi^{-1}\varphi : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \\ &\leq \|\varphi\|_{cb} \|u\varphi^{-1} : Y \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \\ &\leq \|\varphi\|_{cb} \|\varphi^{-1}\|_{cb} \leq C \end{aligned}$$

Conversely, let $\iota : X \rightarrow B(\mathcal{H})$ be a complete isometric inclusion, then by assumption, $\|\iota : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \leq C$. Then $\iota(X) \subset \text{Max}(B(\mathcal{H}))$ is a submaximal space and since $\iota^{-1} : \iota(X) \rightarrow X$ is a complete contraction, ι defines a C -complete isomorphism from X onto the submaximal space $\iota(X) \subset \text{Max}(B(\mathcal{H}))$. \square

COROLLARY 4.4. *For an operator space X , $d_{sm}(X)$ is the smallest $C > 0$ such that for any complete contraction $u : X \rightarrow B(\mathcal{H})$, $\|u : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} \leq C$*

As an illustration, we compute $d_{sm}(X)$ for some finite dimensional operator spaces.

PROPOSITION 4.5. *Let R_n and C_n denotes the n -dimensional row and column Hilbert spaces and M_n denotes the space of $n \times n$ scalar matrices. Then, we have $d_{sm}(R_n) = d_{sm}(C_n) = \sqrt{n}$ and $d_{sm}(M_n) = n$.*

Proof. Consider the formal identity map $id : R_n \rightarrow \ell_2^n$. From the factorization of the identity map from R_n to C_n , along $\text{Max}(\ell_2^n)$, $R_n \rightarrow \text{Max}(\ell_2^n) \rightarrow C_n$, we have

$$\|id : R_n \rightarrow \text{Max}(\ell_2^n)\|_{cb} \geq \|id : R_n \rightarrow C_n\|_{cb} = \|id : R_n \rightarrow C_n\|_{HS} = \sqrt{n}.$$

Therefore by the above corollary 4.4, $d_{sm}(R_n) \geq \sqrt{n}$. Since $\text{Max}(\ell_2^n)$ itself is a submaximal space, and since $\|id : R_n \rightarrow \text{Max}(\ell_2^n)\|_{cb} = \sqrt{n}$ and $\|id^{-1} : \text{Max}(\ell_2^n) \rightarrow R_n\|_{cb} = 1$, from the definition of $d_{sm}(X)$, we have $d_{sm}(R_n) \leq \sqrt{n}$. Thus $d_{sm}(R_n) = \sqrt{n}$. Similarly $d_{sm}(C_n) = \sqrt{n}$. Note that $id : M_n \rightarrow \text{Max}(M_n)$ has the cb -norm n , so that $d_{sm}(M_n) \geq n$. But $\text{Max}(M_n)$ itself is a submaximal space, hence we have $d_{sm}(M_n) = n$. \square

THEOREM 4.6. *Let X be an operator space. Then X is C -complete isomorphic to a submaximal space, for some $C > 0$, if and only if $d_{sm}(Y) \leq C$ for any finite dimensional subspace Y of X .*

Proof. If X is C -complete isomorphic to a submaximal space, for some $C > 0$, then there exists a submaximal space Z such that $\varphi : X \rightarrow Z$ satisfies $\|\varphi\|_{cb} \|\varphi^{-1}\|_{cb} \leq C$. Then for any subspace Y of X , $\varphi|_Y : Y \rightarrow \varphi(Y) \subset Z$, is such that $\|\varphi|_Y\|_{cb} \|\varphi|_Y^{-1}\|_{cb} \leq C$. This shows that $d_{sm}(Y) \leq C$.

Conversely assume that $d_{sm}(Y) \leq C$ for any finite dimensional subspace Y of X for some $C > 0$. If X is not C -completely isomorphic to any submaximal space, then by theorem 4.3, there exists a Hilbert space \mathcal{H} and a complete contraction $\varphi : X \rightarrow \text{Max}(B(\mathcal{H}))$, with $\|\varphi : X \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} > C$. Therefore, by definition of cb-norm, there exists a finite set of elements $\{x_{i,j}; 1 \leq i, j \leq n\}$ for some $n \in \mathbb{N}$, in X such that $\|x_{i,j}\| \leq 1$ and $\|\varphi(x_{i,j})\| > C$. Choose Y as the linear span of $\{x_{i,j}; 1 \leq i, j \leq n\}$. Then Y is a finite dimensional subspace of X , and $\|\varphi|_Y : Y \rightarrow \text{Max}(B(\mathcal{H}))\|_{cb} > C$. So by theorem 4.3, Y is not C -complete isomorphic to a submaximal space. This implies that $d_{sm}(Y) > C$, which is a contradiction. \square

A Q -space ([10]) is an (operator) quotient of a minimal space. In a similar way of defining $d_{sm}(X)$, we can have $d_Q(X) = \inf\{d_{cb}(X, Y); Y \text{ is a } Q\text{-space}\}$ which measures the distance of a given operator space X to the class of Q -spaces. Also, X is a Q -space if and only if $d_Q(X) = 1$. Note that if X is a Q -space, then X^* is a submaximal space. More precisely, if X is C -completely isomorphic to a Q -space, then X^* is C -completely isomorphic to a submaximal space. Conversely, the dual of a submaximal space is a Q -space. Thus we have the following observation.

THEOREM 4.7. *For any operator space X , we have $d_{sm}(X) = d_Q(X^*)$.*

COROLLARY 4.8. *For $n \in \mathbb{N}$, we have: $d_Q(R_n) = d_Q(C_n) = \sqrt{n}$ and $d_Q(S_1^n) = n$, where $S_1^n = M_n^*$, the space of trace class operators.*

Acknowledgement. The authors extend thanks to referee for his valuable suggestions. The first author is grateful to Prof. Gilles Pisier for his encouragement, kind advice, and valuable discussions on this subject, and to Prof. Eric Ricard for having many discussions on these ideas. Also, the first author would like to thank Prof. V.S. Sunder, the organizer of the Instructional Workshop on the Functional Analysis of Quantum Information Theory in the Institute of Mathematical Sciences, Chennai, India, in January 2012, which gave an opportunity to interact with Prof. Gilles Pisier.

REFERENCES

- [1] D. BLECHER, *The standard dual of an operator space*, Pacific J. Math. **153** (1992), 15–30.
- [2] D. BLECHER AND V. PAULSEN, *Tensor products of operator spaces*, J. Funct. Anal. **99** (1991), 262–292.
- [3] E. EFFROS AND Z.-J. RUAN, *Operator Spaces*, London Math. Soc. Monographs. New Series, Oxford University Press, 2000.
- [4] I.M. GELFAND AND M.A. NAIMARK, *On the imbedding of normed rings into the rings of operators on a Hilbert space*, Math. Sbornik. **12** (1943).
- [5] T. OIKHBERG, *Subspaces of Maximal Operator Spaces*, Integr. equ. oper. theory **48** (2004), 81–102.
- [6] V. PAULSEN, *Representation of function algebras, Abstract operator spaces and Banach space geometry*, J. Funct. Anal. **109** (1992), 113–129.

- [7] V. PAULSEN, *The maximal operator space of a normed space*, Proc. Edinburgh. Math. Soc. **39** (1996), 309–323.
- [8] V. PAULSEN, *Completely Bounded maps and Operator Algebras*, Cambridge University Press, 2002.
- [9] G. PISIER, *Introduction to operator space theory*, Cambridge University Press, 2003.
- [10] E. RICARD, *A tensor norm for \mathcal{Q} -spaces*, J. Operator theory **48** (2002), 431–445.
- [11] Z.-J. RUAN, *Subspaces of C^* -algebras*, J. Funct. Anal. **76** (1988), 217–230.

(Received May 15, 2012)

Vinod Kumar. P
Department of Mathematics
Thunchan Memorial Govt. College
Tirur, Kerala
India
e-mail: vinodunical@gmail.com

M. S. Balasubramani
Department of Mathematics
University of Calicut
Calicut University
P. O. Kerala, India
e-mail: msbalaa@rediffmail.com