

WEAK MAJORIZATION INEQUALITIES FOR SINGULAR VALUES

LIMIN ZOU AND CHUANJIANG HE

(Communicated by R. Bhatia)

Abstract. In this paper, we refine an inequality due to Bhatia and Kittaneh [Linear Algebra Appl. 308 (2000) 203–211], and generalize another inequality by Bhatia and Kittaneh [Lett. Math. Phys. 43 (1998) 225–231].

1. Introduction

Let M_n be the space of $n \times n$ complex matrices. Let $\|\cdot\|$ denote any unitarily invariant norm on M_n . We shall always denote the singular values of A by $s_1(A) \geq \dots \geq s_n(A) \geq 0$. Let M_n^+ be the set of positive semidefinite matrix on M_n .

Let $x = (x_1, \dots, x_n)$ be an element of R^n . Let x^\downarrow be the vector obtained by rearranging the coordinates of x in the decreasing order. For $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ belonging to R^n , if

$$\sum_{i=1}^k x_i^\downarrow \leq \sum_{i=1}^k y_i^\downarrow, \quad k = 1, \dots, n,$$

then we say that x is weakly majorized by y , denoted $x \prec_w y$. If the components of x and y are nonnegative and

$$\prod_{i=1}^k x_i^\downarrow \leq \prod_{i=1}^k y_i^\downarrow, \quad k = 1, \dots, n,$$

then we say that x is weakly log-majorized by y , denoted $x \prec_{w \log} y$.

It is well known that $x \prec_{w \log} y$ implies $x \prec_w y$. For more information on majorization and matrix inequalities the reader is referred to [1–3].

Let A and B be positive semidefinite. Bhatia and Kittaneh [4, Theorem 1] (see also [2, p. 77]) obtained the following inequality:

$$s(AB) \prec_w s \left(\left(\frac{A+B}{2} \right)^2 \right). \quad (1.1)$$

Mathematics subject classification (2010): 15A18, 15A42, 47A63.

Keywords and phrases: Weak majorization, singular values, positive semidefinite matrices.

Zhan [5, Theorem 2.2] proved that for any complex number z ,

$$s(A - |z|B) \prec_{w \log} s(A + zB) \prec_{w \log} s(A + |z|B).$$

This is a strengthening of the following inequality:

$$s(A - |z|B) \prec_w s(A + zB) \prec_w s(A + |z|B),$$

which is due to Bhatia and Kittaneh [6, Theorem 2.1]. These authors also proved [6, Theorem 2.2] that for any positive integer m ,

$$s(A^m + B^m) \prec_w s((A + B)^m). \quad (1.2)$$

In Section 2, we shall refine (1.1) and generalize (1.2). Section 3 contains some remarks.

2. Main results

In this section, we first refine (1.1).

THEOREM 2.1. *If $A, B \in M_n$ are positive semidefinite, then*

$$s(AB) \prec_w s\left(\int_0^1 A^{1/2+t} B^{3/2-t} dt\right) \prec_w s\left(\left(\frac{A+B}{2}\right)^2\right). \quad (2.1)$$

Proof. The well-known arithmetic-geometric mean inequality for singular values due to Bhatia and Kittaneh [7] (see also [1, p. 262]) says that

$$2s_j(PQ^*) \leq s_j(P^*P + Q^*Q), \quad j = 1, 2, \dots, n \quad (2.2)$$

for any $P, Q \in M_n$. Let

$$P = A^{1/2}(A+B)^{1/2}, \quad Q = B^{1/2}(A+B)^{1/2}.$$

By (2.2), we have

$$2s_j\left(A^{1/2}(A+B)B^{1/2}\right) \leq s_j\left((A+B)^2\right), \quad j = 1, 2, \dots, n. \quad (2.3)$$

Hiai and Kosaki [8, Corollary 2.3] proved that for all unitarily invariant norms

$$\left\|A^{1/2}XB^{1/2}\right\| \leq \left\|\int_0^1 A^tXB^{1-t} dt\right\| \leq \left\|\frac{AX+XB}{2}\right\|.$$

Putting

$$X = A^{1/2}B^{1/2}$$

in this last inequality, gives

$$\|AB\| \leq \left\|\int_0^1 A^{1/2+t}B^{3/2-t} dt\right\| \leq \left\|\frac{A^{1/2}(A+B)B^{1/2}}{2}\right\|.$$

By Fan's dominance principle, this is equivalent to

$$s(AB) \prec_w s\left(\int_0^1 A^{1/2+t} B^{3/2-t} dt\right) \prec_w s\left(\frac{A^{1/2}(A+B)B^{1/2}}{2}\right). \quad (2.4)$$

It follows from (2.3) and (2.4) that

$$s(AB) \prec_w s\left(\int_0^1 A^{1/2+t} B^{3/2-t} dt\right) \prec_w s\left(\left(\frac{A+B}{2}\right)^2\right).$$

This completes the proof. \square

Next, we shall generalize (1.2). To do this, we need the following result [9, Theorem 2.1].

LEMMA 2.1. *Let $A, B \in M_n$ be normal and let $f: [0, \infty) \rightarrow [0, \infty)$ be concave. Then, for all unitarily invariant norms,*

$$\|f(|A+B|)\| \leq \|f(|A|) + f(|B|)\|. \quad (2.5)$$

THEOREM 2.2. *Let $g(t) = \sum_{k=1}^m a_k t^k$ be a polynomial vanishing at 0 and with non-negative coefficients $a_k, k = 1, \dots, m$. Then for all normal matrices $A, B \in M_n$,*

$$s(g(A) + g(B)) \prec_w s(g(|A+B|)). \quad (2.6)$$

In particular,

$$s(A^m + B^m) \prec_w s(|A+B|^m).$$

Proof. Let X, Y be any pair of normal matrices in M_n and let $f(t) = g^{-1}(t)$ be the reciprocal function of $g(t)$ for $t \in [0, \infty)$. By (2.5), since f is concave, we have

$$s(f(|X+Y|)) \prec_w s(f(|X|) + f(|Y|)).$$

Since g is convex and increasing on $[0, \infty)$, it preserves weak majorization on M_n^+ , hence the above majorization yields

$$s(|X+Y|) \prec_w s(g(f(|X|) + f(|Y|))).$$

Now, set $X = g(A)$ and $Y = g(B)$. We then have

$$s(|g(A) + g(B)|) \prec_w s(g(f(|g(A)|) + f(|g(B)|))).$$

Since $|g(A)| = g(|A|)$, $|g(B)| = g(|B|)$, and $f(g(t)) = t$ on $[0, \infty)$, the last majorization completes the proof. \square

3. Remarks

REMARK 3.1. The inequality (2.3) has been obtained by Bhatia and Kittaneh [4, p. 206]. Here, we give a simple proof.

REMARK 3.2. Let $A, B \in M_n$ be positive semidefinite. Then

$$s_j(AB) \leq s_j \left(\left(\frac{A+B}{2} \right)^2 \right), \quad 1 \leq j \leq n. \quad (3.1)$$

This was a question posed by Bhatia and Kittaneh [4](see also [10-11]), and settled in the affirmative by Drury in [12]. In view of (1.1), (2.1) and (3.1), we ask the following: Is it true that

$$s_j(AB) \leq s_j \left(\int_0^1 A^{1/2+t} B^{3/2-t} dt \right) \leq s_j \left(\left(\frac{A+B}{2} \right)^2 \right), \quad 1 \leq j \leq n?$$

This would be a strengthening of (2.1).

REMARK 3.3. Let $A, B \in M_n$ be positive semidefinite. Tao [13, Theorem 3] proved that the following inequality

$$2s_j \left(A^{1/2} (A+B)^r B^{1/2} \right) \leq s_j \left((A+B)^{r+1} \right), \quad j = 1, \dots, n \quad (3.2)$$

holds for any positive integer r . It is a generalization of (2.3). Bhatia and Kittaneh [10, p. 2186] proved that the inequality (3.2) holds for any positive real number r . Now, we give a simple proof of (3.2). In fact, for any $r > 0$, let

$$P = A^{1/2} (A+B)^{r/2}, \quad Q = B^{1/2} (A+B)^{r/2},$$

we obtain the inequality (2.4) from the inequality (2.1).

Moreover, for any $r, r_1, r_2 > 0$, let

$$P = A^{r_1} (A+B)^{r/2}, \quad Q = B^{r_2} (A+B)^{r/2}.$$

Then, for $j = 1, \dots, n$, we have

$$2s_j \left(A^{r_1} (A+B)^r B^{r_2} \right) \leq s_j \left((A+B)^{r/2} (A^{2r_1} + B^{2r_2}) (A+B)^{r/2} \right).$$

This is a generalization of (3.2).

Acknowledgement. The authors wish to express their heartfelt thanks to the referees and Professor Rajendra Bhatia for their detailed and helpful suggestions for revising the manuscript. This research was supported by Natural Science Foundation Project of Chongqing Municipal Education Commission (KJ121103).

REFERENCES

- [1] R. BHATIA, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [2] X. ZHAN, *Matrix Inequalities*, Lecture Notes in Mathematics, vol. **1790**, Springer-Verlag, Berlin, 2002.
- [3] F. ZHANG, *Matrix Theory-Basic Results and Techniques*, Springer-Verlag, New York, 2011.
- [4] R. BHATIA, F. KITTANEH, *Notes on matrix arithmetic-geometric mean inequalities*, Linear Algebra Appl. **308** (2000), 203–211.
- [5] X. ZHAN, *Singular values of differences of positive semidefinite matrices*, SIAM J. Matrix Anal. Appl. **22** (2000), 819–823.
- [6] R. BHATIA, F. KITTANEH, *Norm inequalities for positive operators*, Lett.Math.Phys. **43** (1998), 225–231.
- [7] R. BHATIA, F. KITTANEH, *On the singular values of a product of operators*, SIAM J. Matrix Anal. Appl. **11** (1990), 272–277.
- [8] F. HIAI, H. KOSAKI, *Means for matrices and comparison of their norms*, Indiana Univ.Math.J. **48** (1999), 899–936.
- [9] J.-C. BOURIN, *A matrix subadditivity inequality for symmetric norms*, Proc.Amer.Math.Soc. **138** (2010), 495–504.
- [10] R. BHATIA, F. KITTANEH, *The matrix arithmetic-geometric mean inequality revisited*, Linear Algebra Appl. **428** (2008), 2177–2191.
- [11] R. BHATIA, *Positive Definite Matrices*, Princeton University Press, Princeton, 2007.
- [12] S. W. DRURY, *On a question of Bhatia and Kittaneh*, Linear Algebra Appl. **437** (2012), 1955–1960.
- [13] Y. TAO, *More results on singular value inequalities of matrices*, Linear Algebra Appl. **416** (2006), 724–729.

(Received April 14, 2012)

Limin Zou
 College of Mathematics and Statistics
 Chongqing University
 Chongqing, 401331, P. R. China
 and
 School of Mathematics and Statistics
 Chongqing Three Gorges University
 Chongqing, 404100, P. R. China
 e-mail: limin-zou@163.com

Chuanjiang He
 Mathematical Sciences Research Institute in Chongqing
 Chongqing University
 Chongqing, 401331, P. R. China
 and
 College of Mathematics and Statistics
 Chongqing University
 Chongqing, 401331, P. R. China