

DIAGRAM VECTORS AND TIGHT FRAME SCALING IN FINITE DIMENSIONS

MARTIN S. COPENHAVER, YEON HYANG KIM, CORTNEY LOGAN, KYANNE
MAYFIELD, SIVARAM K. NARAYAN, MATTHEW J. PETRO AND JONATHAN
SHEPERD

Abstract. We consider frames in a finite-dimensional Hilbert space \mathcal{H}_n where frames are exactly the spanning sets of the vector space. The diagram vector of a vector in \mathbb{R}^2 was previously defined using polar coordinates and was used to characterize tight frames in \mathbb{R}^2 in a geometric fashion. Reformulating the definition of a diagram vector in \mathbb{R}^2 we provide a natural extension of this notion to \mathbb{R}^n and \mathbb{C}^n . Using the diagram vectors we give a characterization of tight frames in \mathbb{R}^n or \mathbb{C}^n . Further we provide a characterization of when a unit-norm frame in \mathbb{R}^n or \mathbb{C}^n can be scaled to a tight frame. This classification allows us to determine all scaling coefficients that make a unit-norm frame into a tight frame.

Mathematics subject classification (2010): 42C15, 05B20, 15A03.

Keywords and phrases: Frames, tight frames, tight frame scaling, diagram vectors, Gramian operator.

REFERENCES

- [1] J. J. BENEDETTO AND M. FICKUS, *Finite Normalized Tight Frames*, Adv. Comput. Math., (2003), **18**, p. 357–385.
- [2] P. G. CASAZZA AND N. LEONHARD, *Classes of finite equal norm Parseval frames*, Frames and operator theory in analysis and signal processing, Contemp. Math., Vol. 451, p. 11–31, Amer. Math. Soc., Providence, RI (2008).
- [3] P. G. CASAZZA, M. FICKUS, J. KOVAČEVIĆ, M. T. LEON AND J. C. TREMAIN, *A Physical Interpretation of Finite Frames*, Appl. Numer. Harmon. Anal., (2006), **2–3**, p. 51–76.
- [4] P. G. CASAZZA AND J. KOVAČEVIĆ, *Equal-Norm Tight Frames with Erasures*, Advances in Computational Mathematics, **392**, p. 387–430, (2006).
- [5] E. W. CHENEY, *Introduction to approximation theory*, American Mathematical Society, (2000), Providence, Rhode Island.
- [6] I. DAUBECHIES, *Ten Lectures on Wavelets*, SIAM, (1992), Philadelphia, PA.
- [7] R. J. DUFFIN AND A. C. SHAEFFER, *A Class of Nonharmonic Fourier Series*, Transactions of the American Mathematical Society, **72**, (1952), p. 341–366.
- [8] D. HAN, K. KORNELSON, D. LARSON AND E. WEBER, *Frames for undergraduates*, Student Mathematical Library, Vol. 40, American Mathematical Society, Providence, RI, (2007), p. xiv+295.
- [9] R. A. HORN AND C. R. JOHNSON, *Matrix Analysis*, Cambridge University Press, (1985).
- [10] K. V. MITAL, *Optimization Methods in Operations Research and System Analysis*, New Age International, (2007).
- [11] A. RON AND Z. SHEN, *Frames and stable bases for shift-invariant subspaces of $L_2(\mathbb{R}^d)$* , Canadian Journal of Mathematics (Journal Canadien de Mathématiques), Vol. 47, (1995), **5**, p. 1051–1094.
- [12] F. ROSENBLATT, *The perceptron: A theory of statistical separability in cognitive systems*, Cornell Aeronautical Laboratory, Inc., Rep. No. VG-1196-G-1. U.S. Department of Commerce, Office of Technical Services, PB 151247, (1958), p. xii+262.
- [13] N. SOHEILI AND J. PENA, *A smooth perceptron algorithm*, (2011).