

## BISHOP'S PROPERTY ( $\beta$ ), HYPERCYCLICITY AND HYPERINVARIANT SUBSPACES

SALAH MECHERI

**Abstract.** The question whether every operator on  $H$  has an hyperinvariant subspace is one of the most difficult problems in operator theory. The purpose of this paper is to make a beginning on the hyperinvariant subspace problems for another class of operators closely related to the normal operators namely, the class of  $k$ -quasi-class  $A$  operators. A necessary and sufficient condition for the hypercyclicity of the adjoint of a quasi-class  $A$  operator is also presented.

*Mathematics subject classification (2010):* 47B20, 47A15.

*Keywords and phrases:* Hyperinvariant subspace, hypercyclicity, quasi-class  $A$  operators.

### REFERENCES

- [1] P. AIENA, S. JOSE EDUARDO, *On left and right poles of the resolvent*, Acta Sci. Math. (Szeged) **74** (2008), 669–687.
- [2] A. ALUTHGE, *On  $p$ -hyponormal operators for  $0 < p < 1$* , Integr. Equat. Oper. Th. **13** (1990), 307–315.
- [3] A. ALUTHGE AND D. WANG,  $\omega$ -hyponormal operators, Integral Equation operator theory, **36** (2000), 1–10.
- [4] T. ANDO, *Operators with a norm conditions*, Acta. Sci. Math (Szeged), **33** (1972), 169–178.
- [5] P. BOURDON, *Orbits of hyponormal operators*, Michigan Math. J. **44** (1997), 444–447.
- [6] R. E. CURTO AND Y. M. HAN, *Weyl's theorem for algebraically paranormal operators*, Integr. Equa. Oper. Theory. **47** (2003), 307–314.
- [7] I. COLOJOARA, C. FOIAS, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
- [8] J. ESCHMEIER, *Invariant subspaces for subscalar operators*, Arch. Math. **52** (1989), 562–570.
- [9] T. FURUTA, M. ITO, T. YAMAZAKI, *A subclass of paranormal operators including class of log-hyponormal and several related classes*, Sci. Math. **1** (1998), 389–403.
- [10] T. FURUTA, *Invitation to linear operatorss-From Matrices to bounded Linear Operators in Hilbert space*, Taylor and Francis, London, 2001.
- [11] T. FURUTA, *On the class of paranormal operators*, Proc. Japan Acad. **43** (1967), 594–598.
- [12] C. FOIAS, I. B. JUNG, E. KO, C. PEARCY, *Hyperinvariant subspaces for some subnormal operators*, Trans. Amer. Math. Soc. **359** (2007), 2899–2913.
- [13] N. S. FIELDMAN, V. G. MILLER, AND T. L. MILLER, *Hypercyclic and supercyclic cohyponormal operators*, Acta. Sci. Math (Szeged). **68** (2002), 965–990.
- [14] F. GAO AND X. FANG, *On  $k$ -quasi-class  $A$  operators*, Jour. Ineq. Appl., Article ID 921634, 10 page (2009).
- [15] F. GAO AND X. FANG, *Weyl's theorem for algebraically  $k$ -quasi-class  $A$  operators*, Opuscula Mathematica, **32**(2012), 125–136.
- [16] F. GAO AND X. FANG, *The Fuglede-Putnam's Theorem and Putnam's inequality for quasi-class  $(A,k)$  operators*, Ann. Funct. Anal. **2**, 1 (2011), 105–113.
- [17] B. HOOVER, *Hyperinvariant subspaces for  $n$ -normal operators*, Acta. Sci. Math. (Szeged). **32** (1972), 121–126.
- [18] IN HO JEAN AND IN HYOUN KIM, *On operators satisfying  $T^*|T^2|T \geq T^*|T|^2T$* , Linear Algebra. Appl., **418** (2006), 854–862.

- [19] S. JUNG, E. KO, AND M. LEE, *On class A operators*, Studia Math. **198** (2010), 249–260.
- [20] H. W. KIM, C. H. PEARCY, *Subnormal operators and hyperinvariant subspaces*, Illinois J. Math. **23** (1979), 459–463.
- [21] R. LANGE AND S. WANG, *New approaches in spectral decomposition*, Contemp. Math., 128, Amer. Math. Soc., 1992.
- [22] K. B. LAURSEN AND M. M. NEUMANN, *An introduction to local spectral theory*, London Math. Soc. Monogr. (N.S) 20, Clarendon Press, Oxford, 2000.
- [23] S. MECHERI, *Bishop's property ( $\beta$ ) and Riesz idempotent of  $k$ -quasi-paranormal operators*, Banach J. Math. Anal. **6** (2012), 147–154.
- [24] S. MECHERI, *Bishop's property ( $\beta$ ), SVEP and Dunford Property (C)*, Electronic Jour. Lin. Alg. **23** (2012), 523–529.
- [25] V. G. MILLER, *Remarks on finitely hypercyclic and finitely hypercyclic operators*, Integral Equations Operator Theory. **29** (1997), 110–115.
- [26] S. Z. NAGY, C. FOIAS, *Harmonic Analysis of operators in Hilbert spaces*, North Holland, Amsterdam, 1970.
- [27] M. PUTINAR, *Hyponormal operators are subscalar*, J. Operator Theory. **12** (1984), 385–395.
- [28] M. PUTINAR, *Hyponormal operators and eigendistribution*, Oper. Theory Adv. Appl. **17** (1986), 249–273.
- [29] H. RADJAVI, P. ROSENTHAL, *Invarinat subspaces*, Springer, 1973.
- [30] K. TANAHASHI, *Putnam's Inequality for log-hyponormal operators*, Integr. equ. oper. theory, **43** (2004), 364–372.
- [31] D. XIA, *Spectral theory of hyponormal operators*, Birkhauser Verlag, Boston, 1983.
- [32] T. YAMAZAKI, *Relations between two equalities  $(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{r}{r+p}} \geq B^r$  and  $A^p \geq (A^{\frac{p}{2}}B^rA^{\frac{p}{2}})^{\frac{p}{r+p}}$  and their applications*, Integral Equation Operator Theory. **44** (2002), 442–450.
- [33] T. YAMAZAKI AND M. YANAGIDA, *A further generalization of paranormal operators*, Sci. Math. **3** (2000), 23–31.