

TIGHT PROJECTIONS OF FRAMES ON INFINITE DIMENSIONAL HILBERT SPACES

JOHN JASPER

Abstract. We characterize the frames on an infinite dimensional separable Hilbert space that can be projected to a tight frame for an infinite dimensional subspace. A result of Casazza and Leon states that an arbitrary frame for a $2N$ – or $(2N - 1)$ -dimensional Hilbert space can be projected to a tight frame for an N -dimensional subspace. Surprisingly, we demonstrate a large class of frames for infinite dimensional Hilbert spaces which cannot be projected to a tight frame for any infinite dimensional subspace.

Mathematics subject classification (2010): Primary: 46C05; Secondary: 47A20.

Keywords and phrases: Frames, projections of frames, compressions.

REFERENCES

- [1] P. G. CASAZZA, M. LEON, *Projections of frames*, Proc. SPIE 5914, 591402 (2005), 277–289.
- [2] O. CHRISTENSEN, *An introduction to frames and Riesz bases*, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Inc., Boston, MA, 2003.
- [3] J. B. CONWAY, *A course in functional analysis*, Second edition, Graduate Texts in Mathematics, Springer-Verlag, New York, 1990.
- [4] M. S. COPENHAVER, Y. H. KIM, C. LOGAN, K. MAYFIELD, S. K. NARAYAN, M. J. PETRO, J. SHEPERD, *Diagram vectors and tight frame scaling in finite dimensions*, preprint, 2012.
- [5] M. FICKUS, D. G. MIXON AND M. J. POTEET, *Frame completions for optimally robust reconstruction*, preprint arXiv:1107.1912, 2011.
- [6] G. KUTYNIOK, K. A. OKOUDJOU, F. PHILIPP, E. K. TULEY, *Scalable frames*, preprint arXiv:1204.1880v3, 2012.
- [7] P. MASSEY, M. RUIZ, D. STOJANOFF, *Optimal completions of a frame*, preprint arXiv:1206.3588v1, 2012.