

A BOUND FOR THE HILBERT–SCHMIDT NORM OF GENERALIZED COMMUTATORS OF NONSELF–ADJOINT OPERATORS

MICHAEL GIL’

(Communicated by F. Kittaneh)

Abstract. Let A, \tilde{A} and B be bounded linear operators in a Hilbert space, and $f(z)$ be a function regular on the convex hull of the union of the spectra of A and \tilde{A} . Let SN_2 be the ideal of Hilbert-Schmidt operators. In the paper, a sharp estimate for the Hilbert-Schmidt norm of the commutator $f(A)B - Bf(\tilde{A})$ is established, provided $AB - B\tilde{A} \in SN_2$, $A - A^* \in SN_2$ and $\tilde{A} - \tilde{A}^* \in SN_2$. Here the star means the adjointness. Our results are new even in the finite dimensional case.

1. Introduction and statement of the main result

Let \mathcal{H} be a separable complex Hilbert space with a scalar product (\cdot, \cdot) , the norm $\|\cdot\| = \sqrt{(\cdot, \cdot)}$ and unit operator I . $\mathcal{B}(\mathcal{H})$ means the algebra of all bounded linear operators in \mathcal{H} . For an $A \in \mathcal{B}(\mathcal{H})$, $\sigma(A)$ denotes the spectrum, $R_z(A) = (A - zI)^{-1}$ ($z \notin \sigma(A)$) is the resolvent, $r_s(A)$ is the spectral radius, A^* is the adjoint operator. SN_2 denotes the ideal of Hilbert-Schmidt operators C with the finite norm $N_2(C) := (\text{Trace}(CC^*))^{1/2}$.

Let $\Omega \supset \sigma(A)$ be an open set whose boundary L consists of a finite number of rectifiable Jordan curves, oriented in the positive sense customary in the theory of complex variables. Suppose that $\Omega \cup L$ is contained in the domain of analyticity of a scalar-valued function f . Then $f(A)$ is defined by

$$f(A) = -\frac{1}{2\pi i} \int_L f(z)R_z(A)dz. \tag{1.1}$$

Furthermore, for $A, B, \tilde{A} \in \mathcal{B}(\mathcal{H})$, $[A, B] := AB - BA$ is the commutator, $[A, B, \tilde{A}] := AB - B\tilde{A}$ is the generalized commutator; $[f(A), B] := f(A)B - Bf(A)$ and $[f(A), B, f(\tilde{A})] := f(A)B - Bf(\tilde{A})$ will be called the function commutator and the generalized function commutator, respectively.

As it is well-known, the generalized function commutator plays an essential role in the perturbation theory of operators, cf. [2] and references therein. Various estimates for the generalized function commutator and its particular cases have been derived by many mathematicians. For details and further references we recommend the

Mathematics subject classification (2010): 47B47, 47A30, 47B10, 47A56, 47A55.

Keywords and phrases: Generalized commutator, inequality, nonself-adjoint operators, operator function, Hilbert-Schmidt norm.

papers [1, 3, 4, 11]. Besides, mainly selfadjoint and normal operators have been considered. Recently, for a class of analytic functions, Kittaneh [10] considered the operator $f(A)B - Bf(\tilde{A})$ in symmetric ideals of \mathcal{H} . Besides A satisfies the inequality

$$\|R_z(A)\| \leq \frac{1}{\text{dist}(z, \sigma(A))}.$$

The similar condition is satisfied by \tilde{A} . In that paper, bounds are established for an arbitrary unitarily invariant norm of the commutator.

In the present paper we consider the generalized function commutator with non-normal operators satisfying the conditions

$$A_I := (A - A^*)/2i \in SN_2, \quad \tilde{A}_I := (\tilde{A} - \tilde{A}^*)/2i \in SN_2, \tag{1.2}$$

and

$$K := AB - B\tilde{A} \in SN_2. \tag{1.3}$$

To formulate the result put

$$g_I(A) := \left[2N_2^2(A_I) - 2 \sum_{k=1}^{\infty} |Im \lambda_k(A)|^2 \right]^{1/2},$$

where $\lambda_k(A)$ ($k = 1, 2, \dots$) are the nonreal eigenvalues of A taken with their multiplicities. If A is normal, then $g_I(A) = 0$, cf. [5, Lemma 7.7.2]. Obviously, $g_I(A) \leq \sqrt{2}N_2(A_I)$.

Denote by $co(A, \tilde{A})$ the closed convex hull of $\sigma(A) \cup \sigma(\tilde{A})$. Now we are in a position to formulate the main result of the paper.

THEOREM 1.1. *Let conditions (1.2) and (1.3) hold. Let $f(\lambda)$ be holomorphic on a neighborhood of $co(A, \tilde{A})$. Then with the notations*

$$\psi_{j,k} := \sup_{z \in co(A, \tilde{A})} \frac{|f^{(k+j+1)}(z)|}{\sqrt{k!j!(k+j+1)!}} \quad (j, k = 0, 1, 2, \dots),$$

we have the inequality

$$N_2(f(A)B - Bf(\tilde{A})) \leq N_2(K) \sum_{j,k=0}^{\infty} \psi_{j,k} g_I^j(A) g_I^k(\tilde{A}).$$

The proof of this theorem is presented in the next two sections. Theorem 1.1 particularly generalizes the main results from [6, 7].

If A and \tilde{A} are normal operators, then Theorem 1.1 implies the inequality.

$$N_2(f(A)B - Bf(\tilde{A})) \leq N_2(K) \sup_{z \in co(A, \tilde{A})} |f'(z)|.$$

If $A \in SN_2$, then due to [5, Lemma 6.5.2] we have $g_I(A) = g(A)$, where

$$g(A) := \left[N_2^2(A) - \sum_{k=1}^{\infty} |\hat{\lambda}_k(A)|^2 \right]^{1/2}.$$

Here $\hat{\lambda}_k(A)$ ($k = 1, 2, \dots$) are all the eigenvalues of A taken with their multiplicities. In addition,

$$g^2(A) \leq N_2^2(A) - |\text{Trace } A^2|,$$

since

$$\sum_{k=1}^{\infty} |\hat{\lambda}_k(A)|^2 \geq |\text{Trace } A^2|.$$

EXAMPLE 1.2. Let $f(A) = e^{At}, t \geq 0$. Then

$$\sup_{z \in co(A, \tilde{A})} \left| \frac{d^{k+j+1} e^{zI}}{dz^{k+j+1}} \right| = e^{\alpha t} t^{k+j+1} \quad (j, k = 0, 1, 2, \dots; t \geq 0),$$

where

$$\alpha := \max\{\sup \text{Re} \sigma(A), \sup \text{Re} \sigma(\tilde{A})\}.$$

Thus,

$$\psi_{j,k} = \frac{e^{\alpha t} t^{k+j+1}}{\sqrt{k!j!(k+j+1)!}} \quad (j, k = 0, 1, 2, \dots).$$

Due to Theorem 1.1

$$N_2(e^{At}B - Be^{\tilde{A}t}) \leq e^{\alpha t} N_2(K) \sum_{j,k=0}^{\infty} \frac{t^{k+j+1} g_I^k(A) g_I^j(\tilde{A})}{\sqrt{k!j!(k+j+1)!}} \quad (t \geq 0).$$

2. Auxiliary results

In this section our reasonings are valid if \mathcal{H} is considered as a Banach space.

Let $A, \tilde{A}, B \in \mathcal{B}(\mathcal{H})$. Then for any $z \notin \sigma(A) \cup \sigma(\tilde{A})$, we have

$$(zI - A)^{-1}B - B(zI - \tilde{A})^{-1} = (Iz - A)^{-1}K(Iz - \tilde{A})^{-1}. \tag{2.1}$$

This well known identity can be checked by multiplying the both sides of (2.1) by $zI - A$ from the left and by $zI - \tilde{A}$ from the right.

Formulas (2.1) and (1.1) imply

$$\begin{aligned} f(A)B - Bf(\tilde{A}) &= -\frac{1}{2\pi i} \int_L f(z)(R_z(A)B - BR_z(\tilde{A}))dz \\ &= \frac{1}{2\pi i} \int_L f(z)R_z(A)KR_z(\tilde{A})dz \end{aligned} \tag{2.2}$$

for any $f(z)$ regular on a neighborhood of $\sigma(A) \cup \sigma(\tilde{A})$.

Put $\Omega(r) = \{|z| \leq r\}$ and $\partial\Omega(r) = \{|z| = r\}$. In the rest of this section it is assumed that $f(z)$ is regular on $\Omega(r)$ with $r > r_s(A, \tilde{A}) := \max\{r_s(A), r_s(\tilde{A})\}$. Take into account that

$$R_\lambda(A) = - \sum_{k=0}^{\infty} \frac{A^k}{\lambda^{k+1}} \quad (|\lambda| > r_s(A)).$$

Then by the previous lemma

$$f(A)B - Bf(\tilde{A}) = \frac{1}{2\pi i} \int_{\partial\Omega(r)} f(z)R_z(A)KR_z(\tilde{A})dz = \sum_{j,k=0}^{\infty} \frac{1}{2\pi i} \int_{\partial\Omega(r)} \frac{f(z)dz}{z^{k+j+2}} A^j K \tilde{A}^k.$$

Or

$$f(A)B - Bf(\tilde{A}) = \sum_{j,k=0}^{\infty} f_{j+k+1} A^j K \tilde{A}^k, \tag{2.3}$$

where f_j are the Taylor coefficients of f at zero.

If, in particular, $f(z) = z^m$, for an integer $m \geq 1$, then $f_{jk} = 0$ for $j+k+1 \neq m$ and $f_{jk} = 1$ for $j+k+1 = m$. Thus we arrive at the identity

$$A^m B - B \tilde{A}^m = \sum_{j=0}^{m-1} A^j K \tilde{A}^{m-j-1}. \tag{2.4}$$

3. Proof of Theorem 1.1

Let A, B and \tilde{A} have n -dimensional ranges ($n < \infty$). Then by the triangular (Schur) representation

$$A = D + V \quad (\sigma(A) = \sigma(D)), \tag{3.1}$$

where D is a normal and V is a nilpotent operators having the joint invariant subspaces. Similarly,

$$\tilde{A} = \tilde{D} + \tilde{V} \quad (\sigma(\tilde{A}) = \sigma(\tilde{D})), \tag{3.2}$$

where \tilde{D} is a normal and \tilde{V} is a nilpotent operators having the same invariant subspaces. Let us prove that

$$N_2(f(A)B - Bf(\tilde{A})) \leq N_2(K) \sum_{j,k=0}^{n-1} \psi_{j,k} N_2^j(V) N_2^k(\tilde{V}). \tag{3.3}$$

Indeed, by (3.1)

$$R_\lambda(A) = (D + V - I\lambda)^{-1} = (I + R_\lambda(D)V)R_\lambda(D).$$

Note that $R_\lambda(D)V$ is a nilpotent matrix and therefore $(R_\lambda(D)V)^n = 0$. Consequently,

$$R_\lambda(A) = \sum_{k=0}^{n-1} (-1)^k (R_\lambda(D)V)^k R_\lambda(D).$$

Similarly,

$$R_\lambda(\tilde{A}) = \sum_{k=0}^{n-1} (-1)^k (R_\lambda(\tilde{D})\tilde{V})^k R_\lambda(\tilde{D}).$$

So by (2.2) we have

$$f(A)B - Bf(\tilde{A}) = \sum_{m,k=0}^{n-1} C_{mk} \tag{3.4}$$

where

$$C_{mk} = (-1)^{k+m} \frac{1}{2\pi i} \int_L f(\lambda) (R_\lambda(D)V)^m R_\lambda(D)K(R_\lambda(\tilde{D})\tilde{V})^k R_\lambda(\tilde{D})d\lambda.$$

Since D is a diagonal matrix in the orthonormal basis of the triangular representations of A (the Schur basis) $\{e_k\}$ of A , and \tilde{D} is a diagonal matrix in the Schur basis $\{\tilde{e}_k\}$ of \tilde{A} , we can write out

$$R_\lambda(D) = \sum_{j=1}^n \frac{Q_j}{\lambda_j - \lambda}, \quad R_\lambda(\tilde{D}) = \sum_{j=1}^n \frac{\tilde{Q}_j}{\tilde{\lambda}_j - \lambda},$$

where $\lambda_j = \lambda_j(A), \tilde{\lambda}_j = \lambda_j(\tilde{A}), Q_k = (\cdot, e_k)e_k, \tilde{Q}_k = (\cdot, \tilde{e}_k)\tilde{e}_k$. Consequently,

$$C_{mk} = \sum_{i_1=1}^n Q_{i_1} V \sum_{i_2=1}^n Q_{i_2} V \dots V \sum_{i_{m+1}=1}^n Q_{i_{m+1}} K \sum_{j_1=1}^n \tilde{Q}_{j_1} \tilde{V} \sum_{j_2=1}^n \tilde{Q}_{j_2} \tilde{V} \dots \tag{3.5}$$

$$\tilde{V} \sum_{j_{k+1}=1}^n \tilde{Q}_{j_{k+1}} J_{i_1, i_2, \dots, i_{m+1}, j_1 j_2 \dots j_{k+1}}.$$

Here

$$J_{i_1, i_2, \dots, i_{m+1}, j_1 j_2 \dots j_{k+1}} = \frac{(-1)^{k+m}}{2\pi i} \int_L \frac{f(\lambda)d\lambda}{(\lambda_{i_1} - \lambda) \dots (\lambda_{i_{m+1}} - \lambda)(\tilde{\lambda}_{j_1} - \lambda) \dots (\tilde{\lambda}_{j_{k+1}} - \lambda)}.$$

Below the symbol $|V|$ means the operator whose entries are absolute values of the entries of V in the basis $\{e_k\}$ and $|\tilde{V}|$ means the operator whose entries are absolute values of the entries of \tilde{V} in the basis $\{\tilde{e}_k\}$. That is, if

$$Ve_k = \sum_{j=1}^{k-1} a_{jk}e_j, \quad \tilde{V}e_k = \sum_{j=1}^{k-1} \tilde{a}_{jk}\tilde{e}_j,$$

then $|V|$ and $|\tilde{V}|$ are defined by

$$|V|e_k = \sum_{j=1}^{k-1} |a_{jk}|e_j \quad \text{and} \quad |\tilde{V}|e_k = \sum_{j=1}^{k-1} |\tilde{a}_{jk}|\tilde{e}_j,$$

respectively.

Furthermore, denote $K_{kj} = (K\tilde{e}_j, e_k)$ and $c_{kj}^{(ml)} = (C_{ml}\tilde{e}_j, e_k)$. Then

$$K = \sum_{j,k=1}^n K_{kj}(\cdot, \tilde{e}_j)e_k \quad \text{and} \quad C_{ml} = \sum_{j,k=1}^n c_{kj}^{(ml)}(\cdot, \tilde{e}_j)e_k.$$

Introduce the operators

$$|K| = \sum_{j,k=1}^n |K_{kj}|(\cdot, \tilde{e}_j)e_k \quad \text{and} \quad |C_{ml}| = \sum_{j,k=1}^n |c_{kj}^{(ml)}|(\cdot, \tilde{e}_j)e_k.$$

So $|K|$ and $|C_{ml}|$ are defined as the operators whose entries are absolute values of the entries of $|K|$ and $|C_{ml}|$ in the corresponding base.

By [5, Lemma 1.5.1],

$$|J_{i_1, i_2, \dots, i_{m+1}, j_1, j_2, \dots, j_{k+1}}| \leq \tilde{\Psi}_{m,k} := \sup_{z \in \text{co}(A, \tilde{A})} \frac{|f^{(k+m+1)}(z)|}{(m+k+1)!}.$$

Now (3.5) implies

$$|C_{mk}| \leq \tilde{\Psi}_{m,k} \sum_{i_1=1}^n Q_{i_1} |V| \sum_{i_2=1}^n Q_{i_2} |V| \dots |V| \sum_{i_{m+1}=1}^n Q_{i_{m+1}} |K| \sum_{j_1=1}^n \tilde{Q}_{j_2} |\tilde{V}| \dots |\tilde{V}| \sum_{j_{k+1}=1}^n \tilde{Q}_{j_{k+1}}.$$

But

$$\sum_{i=1}^n Q_i = I.$$

Thus,

$$|C_{mk}| \leq \tilde{\Psi}_{m,k} |V|^m |K| |\tilde{V}|^k. \tag{3.6}$$

Note that

$$N_2^2(|K|) = \sum_{k=1}^n \| |K| \tilde{e}_k \|^2 = \sum_{k=1}^n \sum_{j=1}^n |K_{jk}|^2 = N_2^2(K).$$

Hence (3.6) yields the inequality

$$N_2(C_{mk}) \leq \tilde{\Psi}_{m,k} \| |V|^m \| N_2(K) \| |\tilde{V}|^k \|.$$

By [5, Corollary 2.5.2] we have

$$\| |V|^m \| \leq \frac{N_2^m(|V|)}{\sqrt{m!}}.$$

But obviously,

$$N_2^2(|V|) = \sum_{k=2}^n \sum_{j=1}^{k-1} |a_{jk}|^2 = N_2^2(V).$$

Recall that a_{jk} are the entries of V in the Schur basis. Thus,

$$\| |V|^m \| \leq \frac{N_2^m(V)}{\sqrt{m!}}$$

So

$$N_2(C_{mk}) \leq \tilde{\Psi}_{m,k} N_2(K) \frac{N_2^m(V) N_2^k(\tilde{V})}{\sqrt{m!} \sqrt{k!}}.$$

Now (3.4) implies the required inequality (3.3).

Furthermore, taking into account the equality $N_2(V) = g_I(A)$, cf. [5, Lemma 7.7.2], from (3.3) we have

$$N_2(f(A)B - Bf(\tilde{A})) \leq N_2(K) \sum_{j,k=0}^{n-1} \psi_{j,k} g_I^j(A) g_I^k(\tilde{A}), \tag{3.7}$$

provided A and \tilde{A} are n -dimensional.

Now let us recall that there is a sequence of n -dimensional operators A_n , strongly converging to A , such that $\sigma(A_n) \subseteq \sigma(A)$, cf. [5, Section 7.11]. In addition, $g_I(A_n) \rightarrow g_I(A)$ as $n \rightarrow \infty$. Indeed, according to [8, Theorem I.5.2], the nonreal spectrum of A under conditions (1.2) consists of no more countable number of points which are normal eigenvalues (that is, isolated and having finite multiplicities). Denote by \mathcal{E} the linear closed convex hull of all the root vectors of A corresponding to non-real eigenvalues. Choice in each root subspace a Jordan basis. Then we obtain vectors ϕ_k for each of which either $A\phi_k = \lambda_k(A)\phi_k$, or $A\phi_k = \lambda_k(A)\phi_k + \phi_{k+1}$. Orthogonalizing the system $\{\phi_k\}$, we obtain the (orthonormal) Schur basis $\{e_k\}$ of the triangular representation:

$$Ae_k = a_{1k}e_1 + a_{2k}e_2 + \dots + a_{kk}e_k \quad (k = 1, 2, \dots) \tag{3.8}$$

with $a_{kk} = \lambda_k(A)$ (see [8, Section II.6]). Besides, \mathcal{E} is an invariant subspace of A . Let P be the orthogonal projection of \mathcal{H} onto \mathcal{E} and $C = AP = PAP$. So $\sigma(C)$ consists of the nonreal spectrum of A . Denote $P_1 = I - P$, $M = P_1AP_1$ and $W = PAP_1$. We have

$$A = (P + P_1)A(P + P_1) = C + M + W,$$

since $P_1AP = P_1PAP = 0$. Take into account that

$$(C + M - \lambda)^{-1} = (C - P\lambda)^{-1} + (M - P_1\lambda)^{-1},$$

where the inverse $(C - P\lambda)^{-1}$ is understood in $P\mathcal{H}$ and the inverse $(M - P_1\lambda)^{-1}$ is understood in $P_1\mathcal{H}$. Hence,

$$(C + M - \lambda)^{-1}W = (C + M - \lambda)^{-1}PAP_1 = P(C - P\lambda)^{-1}PAP_1$$

and therefore, $((C + M - \lambda)^{-1}W)^2 = 0$. Consequently, for sufficiently large λ ,

$$(I + (C + M - \lambda)^{-1}W)^{-1} = \sum_{k=0}^{\infty} (-1)^k (C + M - \lambda)^{-1}W^k = I - (C + M - \lambda)^{-1}W.$$

So for all regular λ ,

$$\begin{aligned} (A - \lambda)^{-1} &= (C + M + W - \lambda)^{-1} = (I + (C + M - \lambda)^{-1}W)^{-1}(C + M - \lambda)^{-1} \\ &= (C + M - \lambda)^{-1} + (C + M - \lambda)^{-1}W(C + M - \lambda)^{-1}. \end{aligned}$$

Consequently,

$$\sigma(A) = \sigma(C) \cup \sigma(M). \tag{3.9}$$

Moreover,

$$\begin{aligned} (AP_1 - \lambda)^{-1} &= (W + M - \lambda)^{-1} = (I + (M - \lambda)^{-1}W)^{-1}(M - \lambda)^{-1} \\ &= (M - \lambda)^{-1} + (M - \lambda)^{-1}W(M - \lambda)^{-1}. \end{aligned}$$

Thus $\sigma(AP_1) = \sigma(M)$ and it is real. Let C_n ($n = 1, 2, \dots$) be the sequence of n -dimensional operators, defined by

$$C_n e_k = a_{1k} e_1 + a_{2k} e_2 + \dots + a_{kk} e_k \quad (k = 1, \dots, n).$$

Then according to (3.8) $\lambda_k(C_n) = \lambda_k(A)$ ($k = 1, \dots, n$) and $C_n \rightarrow C$ at least strongly. Let Q_n be a sequence of n -dimensional orthogonal projections strongly converging to I . Then $Q_n AP_1 \rightarrow AP_1$ strongly. Since $C_n = C_n P$, according to (3.9),

$$\sigma(Q_n AP_1 + C_n) = \sigma(Q_n AP_1) \cup \sigma(C_n).$$

Besides, $A_n := Q_n AP_1 + C_n \rightarrow A$ strongly. Due to the upper semicontinuity of the spectrum [9, p. 56, Problem 103], $\lim_{n \rightarrow \infty} \sigma(A_n) \subseteq \sigma(A)$. Hence, since $\sigma(AP_1)$ is real, we can write

$$\lim_{n \rightarrow \infty} \Im \sigma(Q_n AP_1) = 0$$

and therefore, under condition (1.2),

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n |\Im \lambda_k(A_n)|^2 = \lim_{n \rightarrow \infty} \sum_{k=1}^n |\Im \lambda_k(C_n)|^2 = \sum_{k=1}^{\infty} |\Im \lambda_k(C)|^2 = \sum_{k=1}^{\infty} |\Im \lambda_k(A)|^2. \tag{3.10}$$

Since, $A_n \rightarrow A$ strongly, we have $A_n - A_n^* \rightarrow A - A^*$ strongly, but $A - A^* \in SN_2$, and therefore, $N_2(A_n - A_n^*) \rightarrow N_2(A - A^*)$. This and (3.10) implies that really $g_I(A_n) \rightarrow g_I(A)$ as $n \rightarrow \infty$. Moreover, due to the relation $\sigma(A_n) \subseteq \sigma(A)$ we have $\lim_n co(A_n) \subseteq co(A)$. So replacing in (3.7) A and \tilde{A} by A_n and \tilde{A}_n , respectively, we get the required result by passing to the limit $n \rightarrow \infty$ in that inequality. \square

Acknowledgement. I am very grateful to the referee of this paper for his (her) really helpful remarks.

REFERENCES

- [1] R. BHATIA AND F. KITTANEH, *Some inequalities for norms of commutators*, SIAM J. Matrix Anal. Appl. **18** (1997), 258–263.
- [2] M. BIRMAN, AND M. SOLOMYAK, *Double operator integrals in a Hilbert space*, Integral Equations Operator Theory **47**, no. 2, (2003), 131–168.
- [3] K. N. BOYADZHIEV, *Some inequalities for generalized commutators*, Publ. RIMS, Kyoto University, **26**, no. 3 (1990), 521–527.
- [4] K. N. BOYADZHIEV, *Norm estimates for commutators of operators*, J. London Math. Soc. **57** (2) (1998), 739–745.

- [5] M. I. GIL', *Operator Functions and Localization of Spectra*, Lecture Notes In Mathematics, vol. 1830, Springer-Verlag, Berlin, 2003.
- [6] M. I. GIL', *Perturbations of functions of operators in a Banach space*, Math. Phys. Anal. Geom. **13**, no. 1, (2010) 69–82.
- [7] M. I. GIL', *Perturbations of operator functions in a Hilbert Space*, Communications in Mathematical Analysis **13**, no. 2, (2012) 108–115.
- [8] I. C. GOHBERG, AND M. G. KREIN, *Introduction to the Theory of Linear Nonselfadjoint Operators*, Trans. Mathem. Monographs, v. 18, Amer. Math. Soc., Providence, R. I, 1969.
- [9] P. HALMOS, *Hilbert Space Problem Book*, Second edition, Springer Verlag, New York, 1982.
- [10] F. KITTANEH, *Norm Inequalities for commutators of G_1 operators*, Complex Anal. Oper. Theory **10**, (2016) 109–114.
- [11] V. PELLER, *Hankel operators in perturbation theory of unbounded self-adjoint operators*, in the book *Analysis and Partial Differential Equations*, Lecture Notes in Pure and Appl. Math. **122**, Marcel Dekker, New York, 1990, pp. 529–544.

(Received April 1, 2016)

Michael Gil'
Department of Mathematics
Ben Gurion University of the Negev
P. O. Box 653, Beer-Sheva 84105, Israel
e-mail: gilmi@bezeqint.net