

SPECTRA OF LAPLACIANS ON FORMS ON AN INFINITE GRAPH

HÈLA AYADI

(Communicated by M. Embree)

Abstract. In the context of infinite weighted graphs, we consider the discrete Laplacians on 0-forms and 1-forms. Using Weyl's criterion, we prove the relation between the nonzero spectrum of Δ_0 and that of Δ_1 . Moreover, we give an extension of the work of John Lott to characterize the 0-spectrum of these two Laplacians.

1. Introduction

In recent years, much attention has been paid to the analysis of discrete Laplacians and elliptic differential operators acting on graphs [13], [5], [6] and [19]. More precisely, authors have intensively studied the spectrum of the discrete Laplacian on an infinite graph in various areas, for example, harmonic analysis on graphs (see [16], [20]), probability theory especially Markov chains (see [8], [12]), potential theory such as electric networks (see [17], [12]), and so on. In this paper, we define two Laplacians, mentioned in [1] and [3], one as an operator acting on functions on vertices denoted by Δ_0 and the other one acting on functions on edges denoted by Δ_1 . So, it is a natural question to characterize the relation between their spectrum in terms of a certain geometric property of the graph and properties of the operators. Especially we show that the nonzero spectrum of Δ_0 and Δ_1 are the same, by using Weyl's criterion. Moreover, with suitable weight conditions we prove that 0 is in the spectrum of Δ_1 , if the operator Δ_0 is invertible. This result is inspired from J. Lott's work [11] (Proposition 9, p. 12) which proves in the case of a simple graph that 0 is either in the spectrum of the Laplacian on 0-forms, or in the spectrum of the Laplacian on 1-forms. In fact, the major interest of J. Lott concerns the zero-in-the-spectrum question for the Laplace-de Rham operator acting on L^2 differential forms of any degree on a complete connected oriented Riemannian manifold. The article [11] is rather expository and gives some positive answers, in relation with topology, for small dimensions. We finish the paper with examples of constructions of Δ_1 -harmonic nonzero square-integrable functions.

Mathematics subject classification (2010): 39A12, 05C63, 05C05, 47A53.

Keywords and phrases: Infinite weighted graph, discrete Laplacian, Weyl's criterion, spectrum.

2. Preliminaries

2.1. Definition and notation

- A graph G is a couple $(\mathcal{V}, \mathcal{E})$ where \mathcal{V} is a set at most countable whose elements are called vertices and \mathcal{E} is a set of oriented edges, considered as a subset of $\mathcal{V} \times \mathcal{V}$.
- If the graph G has a finite set of vertices, it is called a finite graph. Otherwise, G is called an infinite graph.
- We assume that \mathcal{E} has no self-loops and is symmetric:

$$v \in \mathcal{V} \Rightarrow (v, v) \notin \mathcal{E}, \quad (v_1, v_2) \in \mathcal{E} \Rightarrow (v_2, v_1) \in \mathcal{E}.$$

- Choosing an orientation of G consists of defining a partition of \mathcal{E} : $\mathcal{E}^+ \sqcup \mathcal{E}^- = \mathcal{E}$

$$(v_1, v_2) \in \mathcal{E}^+ \Leftrightarrow (v_2, v_1) \in \mathcal{E}^-.$$

- For $e = (v_1, v_2)$, we denote

$$e^- = v_1, \quad e^+ = v_2 \quad \text{and} \quad -e = (v_2, v_1).$$

- We write $v_1 \sim v_2$ for $e = (v_1, v_2) \in \mathcal{E}$.
- The graph G is connected if any two vertices x, y in \mathcal{V} can be joined by a path of edges γ_{xy} , that means $\gamma_{xy} = \{e_k\}_{k=1, \dots, n}$ such that

$$e_1^- = x, \quad e_n^+ = y \quad \text{and, if } n \geq 2, \quad \forall j; \quad 1 \leq j \leq (n-1) \Rightarrow e_j^+ = e_{j+1}^-.$$

- The degree (or valence) of a vertex x is the number of edges emanating from x . We denote

$$\deg(x) := \#\{e \in \mathcal{E}; e^- = x\}.$$

- If $\deg(x) < \infty, \forall x \in \mathcal{V}$, we say that G is a locally finite graph.

2.2. Weighted graphs

DEFINITION 2.1. A weighted graph (G, c) is given by a graph $G = (\mathcal{V}, \mathcal{E})$ and weights on the edges $c : \mathcal{E} \rightarrow [0, \infty[$ such that

- $c(x, x) = 0, \forall x \in \mathcal{V}$.
- $c(x, y) > 0, \forall (x, y) \in \mathcal{E}$.
- $c(x, y) = c(y, x), \forall (x, y) \in \mathcal{E}$.

If $\sum_{y \sim x} c(x, y) < \infty$ for each $x \in \mathcal{V}$, we can define a weight on \mathcal{V} by

$$\tilde{c}(x) = \sum_{y \sim x} c(x, y), \quad x \in \mathcal{V}.$$

REMARK 2.1. If the graph G is locally finite, the weight \tilde{c} on any vertex is well defined.

EXAMPLES.

An infinite electrical network is a weighted graph (G, c) where the weight c on the edges are called conductances and their reciprocals are called resistances. This is the convention used in the study of random walks on weighted graphs, see [12] and [16]. Then, $\tilde{c}(x) = \sum_{y \in \mathcal{V}} c(x, y)$ is the weight associated to the vertex x .

A graph G is called a simple graph if the edge weights are equal to 1. In this case,

$$\tilde{c}(x) = \deg(x), \quad \forall x \in \mathcal{V}.$$

All the graphs we shall consider in the sequel will be connected, locally finite and weights c given in Definition 2.1.

2.3. Functional spaces

We denote the set of real functions on \mathcal{V} by:

$$\mathcal{C}(\mathcal{V}) = \{f : \mathcal{V} \rightarrow \mathbb{R}\}$$

and the set of functions of finite support by $\mathcal{C}_0(\mathcal{V})$.

Moreover, we denote the set of real skew-symmetric functions on \mathcal{E} by:

$$\mathcal{C}^a(\mathcal{E}) = \{\varphi : \mathcal{E} \rightarrow \mathbb{R} ; \varphi(-e) = -\varphi(e)\}$$

and the set of functions of finite support by $\mathcal{C}_0^a(\mathcal{E})$.

We define on the weighted graph (G, c) the following function spaces endowed with the scalar products.

a)

$$l^2(\mathcal{V}) := \left\{ f \in \mathcal{C}(\mathcal{V}) ; \sum_{x \in \mathcal{V}} \tilde{c}(x) f^2(x) < \infty \right\},$$

with the inner product

$$\langle f, g \rangle_{\mathcal{V}} = \sum_{x \in \mathcal{V}} \tilde{c}(x) f(x) g(x)$$

and the norm

$$\|f\|_{\mathcal{V}} = \sqrt{\langle f, f \rangle_{\mathcal{V}}}.$$

b)

$$l^2(\mathcal{E}) := \left\{ \varphi \in \mathcal{C}^a(\mathcal{E}); \frac{1}{2} \sum_{e \in \mathcal{E}} c(e) \varphi^2(e) < \infty \right\},$$

with the inner product

$$\langle \varphi, \psi \rangle_{\mathcal{E}} = \frac{1}{2} \sum_{e \in \mathcal{E}} c(e) \varphi(e) \psi(e)$$

and the norm

$$\|\varphi\|_{\mathcal{E}} = \sqrt{\langle \varphi, \varphi \rangle_{\mathcal{E}}}.$$

Then, $l^2(\mathcal{V})$ and $l^2(\mathcal{E})$ are separable Hilbert spaces (since \mathcal{V} is countable).

2.4. Operators and properties

The difference operator

$$d : l^2(\mathcal{V}) \longrightarrow l^2(\mathcal{E}),$$

is given by

$$d(f)(e) = f(e^+) - f(e^-).$$

The coboundary operator is δ , the formal adjoint of d . Thus it satisfies

$$\langle df, \varphi \rangle_{\mathcal{E}} = \langle f, \delta \varphi \rangle_{\mathcal{V}} \quad (2.1)$$

for all $f \in l^2(\mathcal{V})$ and for all $\varphi \in l^2(\mathcal{E})$.

As consequence, we have the following formula characterizing δ :

LEMMA 2.1. *The coboundary operator δ is characterized by the formula*

$$\delta \varphi(x) = \frac{1}{\tilde{c}(x)} \sum_{e, e^+ = x} c(e) \varphi(e),$$

for all $\varphi \in l^2(\mathcal{E})$.

Proof. For $f \in l^2(\mathcal{V})$ and $\varphi \in l^2(\mathcal{E})$, using (2.1), we get

$$\begin{aligned} \langle df, \varphi \rangle_{\mathcal{E}} &= \frac{1}{2} \sum_{e \in \mathcal{E}} c(e) df(e) \varphi(e) \\ &= \frac{1}{2} \sum_{e \in \mathcal{E}} c(e) (f(e^+) - f(e^-)) \varphi(e) \\ &= \frac{1}{2} \sum_{x \in \mathcal{V}} f(x) \left(\sum_{e, e^+ = x} c(e) \varphi(e) - \sum_{e, e^- = x} c(e) \varphi(e) \right). \end{aligned}$$

But $c(-e) = c(e)$ and φ is skew-symmetric, so we have

$$\sum_{e, e^+ = x} c(e)\varphi(e) = - \sum_{e, e^- = x} c(e)\varphi(e).$$

Then,

$$\begin{aligned} \langle df, \varphi \rangle_{\mathcal{E}} &= \sum_{x \in \mathcal{V}} \tilde{c}(x)f(x) \left(\frac{1}{\tilde{c}(x)} \sum_{e, e^+ = x} c(e)\varphi(e) \right) \\ &= \langle f, \delta\varphi \rangle_{\mathcal{V}} \end{aligned}$$

and the formula for $\delta\varphi$ follows. \square

DEFINITION 2.2. The Laplacian on 0-forms Δ_0 defined by δd on $l^2(\mathcal{V})$ is given by

$$\Delta_0 f(x) = \frac{1}{\tilde{c}(x)} \sum_{y \sim x} c(x, y) (f(x) - f(y)).$$

In fact, we have

$$\begin{aligned} \Delta_0 f(x) &= \delta(df)(x) \\ &= \frac{1}{\tilde{c}(x)} \sum_{e, e^+ = x} c(e)df(e) \\ &= \frac{1}{\tilde{c}(x)} \sum_{e, e^+ = x} c(e) (f(e^+) - f(e^-)) \\ &= \frac{1}{\tilde{c}(x)} \sum_{y \sim x} c(x, y) (f(x) - f(y)). \end{aligned}$$

DEFINITION 2.3. The Laplacian on 1-forms Δ_1 defined by $d\delta$ on $l^2(\mathcal{E})$ is given by

$$\Delta_1 \varphi(e) = \frac{1}{\tilde{c}(e^+)} \sum_{e_1, e_1^+ = e^+} c(e_1)\varphi(e_1) - \frac{1}{\tilde{c}(e^-)} \sum_{e_2, e_2^+ = e^-} c(e_2)\varphi(e_2).$$

In fact, we have

$$\begin{aligned} \Delta_1 \varphi(e) &= d(\delta\varphi)(e) \\ &= \delta\varphi(e^+) - \delta\varphi(e^-) \\ &= \frac{1}{\tilde{c}(e^+)} \sum_{e_1, e_1^+ = e^+} c(e_1)\varphi(e_1) - \frac{1}{\tilde{c}(e^-)} \sum_{e_2, e_2^+ = e^-} c(e_2)\varphi(e_2). \end{aligned}$$

PROPOSITION 2.1. *The operator Δ_0 is bounded and self-adjoint.*

Proof. For $f, g \in l^2(\mathcal{Y})$, we have

$$\begin{aligned}
 |\langle \Delta_0 f, g \rangle_{\mathcal{Y}}| &= \left| \sum_x \tilde{c}(x) \frac{1}{\tilde{c}(x)} \sum_{y \sim x} c(x, y) (f(x) - f(y)) g(x) \right| \\
 &\leq \sum_x \sum_{y \sim x} c(x, y) |(f(x) - f(y))| |g(x)| \\
 &\leq \sum_x \sum_{y \sim x} c(x, y) |f(x)| |g(x)| + \sum_x \sum_{y \sim x} c(x, y) |f(y)| |g(x)| \\
 &= \sum_x \tilde{c}(x) |f(x)| |g(x)| + \sum_x \sum_{y \sim x} c(x, y) |f(y)| |g(x)| \\
 &\leq \|f\|_{\mathcal{Y}} \|g\|_{\mathcal{Y}} + I
 \end{aligned} \tag{2.2}$$

where $I := \sum_x \sum_{y \sim x} c(x, y) |f(y)| |g(x)|$.

Using the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned}
 I &\leq \sum_x \left(\sum_{y \sim x} c(x, y) |f(y)|^2 \right)^{\frac{1}{2}} \left(\sum_{y \sim x} c(x, y) \right)^{\frac{1}{2}} |g(x)| \\
 &= \sum_x \left(\sum_{y \sim x} c(x, y) f^2(y) \right)^{\frac{1}{2}} (\tilde{c}(x))^{\frac{1}{2}} |g(x)| \\
 &\leq \left(\sum_x \sum_{y \sim x} c(x, y) f^2(y) \right)^{\frac{1}{2}} \left(\sum_x \tilde{c}(x) g^2(x) \right)^{\frac{1}{2}} \\
 &= \left(\sum_y \tilde{c}(y) f^2(y) \right)^{\frac{1}{2}} \left(\sum_x \tilde{c}(x) g^2(x) \right)^{\frac{1}{2}} \\
 &= \|f\|_{\mathcal{Y}} \|g\|_{\mathcal{Y}}.
 \end{aligned} \tag{2.3}$$

Therefore, (2.2) and (2.3) gives

$$|\langle \Delta_0 f, g \rangle_{\mathcal{Y}}| \leq 2 \|f\|_{\mathcal{Y}} \|g\|_{\mathcal{Y}}.$$

But by the definition of the norm of operator, we have

$$\|\Delta_0\| = \sup_{\|f\|=1} \|\Delta_0 f\|_{\mathcal{Y}} = \sup_{\|f\|=1} \sup_{\|g\|=1} \langle \Delta_0 f, g \rangle_{\mathcal{Y}}$$

So $\|\Delta_0\| \leq 2$, which shows that Δ_0 is a bounded operator.

Now, we want to prove the selfadjointness of the operator Δ_0 defined on $l^2(\mathcal{Y})$. As Δ_0 is a bounded operator on $l^2(\mathcal{Y})$, it remains to show that Δ_0 is symmetric.

As we have $\Delta_0 = \delta d$ and δ is the adjoint operator of d , we obtain for f and $g \in l^2(\mathcal{Y})$

$$\begin{aligned}
 \langle \Delta_0 f, g \rangle_{\mathcal{Y}} &= \langle \delta d f, g \rangle_{\mathcal{Y}} \\
 &= \langle d f, d g \rangle_{\mathcal{E}} \\
 &= \langle f, \delta d g \rangle_{\mathcal{Y}} \\
 &= \langle f, \Delta_0 g \rangle_{\mathcal{Y}}. \quad \square
 \end{aligned}$$

REMARK 2.2.

- The operators d and δ are bounded. Indeed, using the inequality $(a - b)^2 \leq 2(a^2 + b^2)$ and the definition of the weights on vertices: $\tilde{c}(x) = \sum_{y \sim x} c(x, y)$, we obtain

$$\begin{aligned} \|df\|_{\mathcal{E}}^2 &= \frac{1}{2} \sum_{(x,y) \in \mathcal{E}} c(x,y) (df(x,y))^2 \\ &= \frac{1}{2} \sum_{(x,y) \in \mathcal{E}} c(x,y) (f(y) - f(x))^2 \\ &\leq \sum_{(x,y) \in \mathcal{E}} c(x,y) (f^2(y) + f^2(x)) \\ &= 2 \sum_{x \in \mathcal{V}} f^2(x) \sum_{y \sim x} c(x,y) \\ &= 2 \sum_{x \in \mathcal{V}} f^2(x) \tilde{c}(x) \\ &= 2 \|f\|_{\mathcal{V}}^2. \end{aligned}$$

So d is bounded, and the same is true for the adjoint δ .

Notice that since Δ_0 is the composite operator of δ and d ; this gives another proof that Δ_0 is bounded.

- It is easy to see that Δ_0 is also positive, since $\langle \Delta_0 f, f \rangle_{\mathcal{V}} = \langle df, df \rangle_{\mathcal{E}} \geq 0$.

COROLLARY 2.1. *As the operator Δ_0 is self-adjoint and positive, its spectrum is real and lies in $[0, 2]$.*

2.5. Weyl's criterion

As our operator is bounded and self-adjoint on a Hilbert space, we can use Weyl's criterion [14] to characterize its spectrum.

Weyl's criterion: Let \mathcal{H} be a separable Hilbert space, and let Δ be a bounded self-adjoint operator on \mathcal{H} . Then λ is in the spectrum of Δ if and only if there exists a sequence $(f_n)_{n \in \mathbb{N}}$ so that $\|f_n\| = 1$ and $\lim_{n \rightarrow \infty} \|(\Delta - \lambda)f_n\| = 0$.

We denote $\sigma(\Delta)$ the spectrum of Δ and we set

- $\sigma_d(\Delta)$ is the set of $\lambda \in \sigma(\Delta)$ which is an isolated point and an eigenvalue with finite multiplicity.
- $\sigma_{ess}(\Delta) := \sigma(\Delta) \setminus \sigma_d(\Delta)$.

3. The relation between the spectrum of Δ_0 and Δ_1

3.1. The nonzero spectrum of Δ_0 and Δ_1

In this section we will prove the relation between the spectrum of Δ_0 and that of Δ_1 , by using Weyl's criterion.

Following [15] and [18] we have the next lemma.

LEMMA 3.1. *Let $\Delta_0 = \delta d$ and $\Delta_1 = d\delta$. Then we have*

1. $d\Delta_0 = \Delta_1 d$.
2. $\delta\Delta_1 = \Delta_0\delta$.

LEMMA 3.2.

1. $\ker\Delta_0 = \ker d$.
2. $\ker\Delta_1 = \ker\delta$.

Proof.

1. Clearly, we have $\ker d \subset \ker\Delta_0$.

On the other hand, if $\Delta_0 f = 0$ for $f \in l^2(\mathcal{V})$ and $f \neq 0$, we have

$$0 = \langle \Delta_0 f, f \rangle_{\mathcal{V}} = \langle df, df \rangle_{\mathcal{E}}.$$

Then $df = 0$ for $f \in l^2(\mathcal{V})$.

2. If $\varphi \in \ker\delta$, then $\varphi \in l^2(\mathcal{E})$ and $\delta\varphi = 0$. Thus, $d\delta\varphi = 0$ and we obtain $\varphi \in \ker\Delta_1$.

For the other inclusion, let $\varphi \in l^2(\mathcal{V})$, $\varphi \neq 0$ such that $\Delta_1\varphi = 0$. Then

$$0 = \langle \Delta_1\varphi, \varphi \rangle_{\mathcal{E}} = \langle \delta\varphi, \delta\varphi \rangle_{\mathcal{E}}.$$

We get $\delta\varphi = 0$ and as a result $\ker\Delta_1 \subset \ker\delta$. \square

We arrive at our main result.

THEOREM 1.

$$\sigma(\Delta_1) \setminus \{0\} = \sigma(\Delta_0) \setminus \{0\}.$$

Proof.

- Let $\lambda \neq 0$ be in the spectrum of Δ_0 . By Weyl's criterion, there exists a sequence $(f_n)_n$ of $l^2(\mathcal{V})$ such that

$$\|f_n\|_{\mathcal{V}} = 1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|(\Delta_0 - \lambda)f_n\|_{\mathcal{V}} = 0.$$

We want to find a sequence $(\varphi_n)_n$ of $l^2(\mathcal{E})$ such that

$$\|\varphi_n\|_{\mathcal{E}} = 1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|(\Delta_1 - \lambda)\varphi_n\|_{\mathcal{E}} = 0.$$

We set

$$\varphi_n := \frac{df_n}{\|df_n\|_{\mathcal{E}}}.$$

First, let us check that $\|df_n\|_{\mathcal{E}} \neq 0$. We have

$$\begin{aligned} \|df_n\|_{\mathcal{E}}^2 &= \langle \Delta_0 f_n, f_n \rangle_{\mathcal{V}} \\ &= \langle (\Delta_0 - \lambda) f_n, f_n \rangle_{\mathcal{V}} + \langle \lambda f_n, f_n \rangle_{\mathcal{V}} \\ &= \underbrace{\langle (\Delta_0 - \lambda) f_n, f_n \rangle_{\mathcal{V}}}_{\text{converges to 0}} + \lambda. \end{aligned}$$

Then, $\lim_{n \rightarrow \infty} \|df_n\|_{\mathcal{E}}^2 = \lambda$. Thus, by positivity of Δ_0 , there exists $A > 0$ and an integer n_0 such that for all $n \geq n_0$, we have $\|df_n\|_{\mathcal{E}} > A$. This implies that the sequence $(\varphi_n)_n$ is well defined.

Now, we verify that $\lim_{n \rightarrow \infty} \|(\Delta_1 - \lambda)\varphi_n\|_{\mathcal{E}} = 0$. By the first assertion of Lemma 3.1 and the fact that the operator d is bounded, we obtain for all n sufficiently large

$$\begin{aligned} \|(\Delta_1 - \lambda)\varphi_n\|_{\mathcal{E}} &= \left\| (\Delta_1 - \lambda) \frac{df_n}{\|df_n\|_{\mathcal{E}}} \right\|_{\mathcal{E}} \\ &= \frac{\|(\Delta_1 - \lambda)df_n\|_{\mathcal{E}}}{\|df_n\|_{\mathcal{E}}} \\ &= \frac{\|d(\Delta_0 - \lambda)f_n\|_{\mathcal{E}}}{\|df_n\|_{\mathcal{E}}} \\ &\leq \frac{\|d\|}{A} \|(\Delta_0 - \lambda)f_n\|_{\mathcal{V}}. \end{aligned}$$

But $\lim_{n \rightarrow \infty} \|(\Delta_0 - \lambda)f_n\|_{\mathcal{V}} = 0$. Therefore, $\lim_{n \rightarrow \infty} \|(\Delta_1 - \lambda)\varphi_n\|_{\mathcal{E}} = 0$ and we can conclude that λ is in the spectrum of $\Delta_1 \setminus \{0\}$.

- The second part of the proof follows in the same fashion, with the roles of d and δ swapped. \square

There is a second method to prove Theorem 1 when 0 is not in the spectrum of Δ_0 .

LEMMA 3.3. *If 0 is not in the spectrum of Δ_0 , then the operator d defined in $l^2(\mathcal{V})$ has a closed range.*

Proof. Let $\varphi \in \overline{\text{Im}d}$, let us check that $\varphi \in \text{Im}d$, that means we look for a function $f \in l^2(\mathcal{V})$ such that $\varphi = df$. We have $\varphi \in \overline{\text{Im}d}$, so there exists a sequence $(\varphi_n)_n$ of $\text{Im}d$ such that $\varphi_n = df_n$, for $f_n \in l^2(\mathcal{V})$. Moreover, the sequence $(\varphi_n)_n$ converges to φ in $l^2(\mathcal{E})$. On the other hand, by assumption 0 is not in the spectrum of Δ_0 which implies the existence of a positive constant C such that

$$\|f\|_{\mathcal{V}} \leq C \|\Delta_0 f\|_{\mathcal{V}}, \forall f \in l^2(\mathcal{V}).$$

But by the definition of the operator norm and Remark 2.2, we obtain

$$\|\Delta_0 f\| = \sup_{g, \|g\|_{\mathcal{Y}}=1} \langle \Delta_0 f, g \rangle_{\mathcal{Y}} \leq \|df\|_{\mathcal{E}} \sup_{g, \|g\|_{\mathcal{Y}}=1} \|dg\|_{\mathcal{E}} \leq \sqrt{2} \|df\|_{\mathcal{E}}.$$

Then

$$\|f\|_{\mathcal{Y}} \leq \sqrt{2} C \|df\|_{\mathcal{E}}, \quad \forall f \in l^2(\mathcal{Y}).$$

Thus

$$\|f_n - f_m\|_{\mathcal{Y}} \leq \sqrt{2} C \|df_n - df_m\|_{\mathcal{E}}, \quad f_n, f_m \in l^2(\mathcal{Y}).$$

And

$$\|f_n - f_m\|_{\mathcal{Y}} \leq \sqrt{2} C \|\varphi_n - \varphi_m\|_{\mathcal{E}}, \quad f_n, f_m \in l^2(\mathcal{Y}).$$

As the sequence $(\varphi_n)_n$ converges, so it is a Cauchy sequence and also $(f_n)_n$ is a Cauchy sequence in $l^2(\mathcal{Y})$ which is complete. Then, $(f_n)_n$ converges to f . By the boundedness of the operator d , we obtain $df_n = \varphi_n$ converges to df and by uniqueness of the limit, we have $df = \varphi$. So φ is in $\text{Im}d$. \square

COROLLARY 3.1. *If 0 is not in the spectrum of Δ_0 , then*

$$\sigma(\Delta_1|_{\text{Im}d}) = \sigma(\Delta_0).$$

Proof. By the first assertion of Lemma 3.1, we obtain

$$\Delta_1 d = d \Delta_0.$$

But by assumption 0 is not in the spectrum of Δ_0 . Then by the first assertion of Lemma 3.2, the operator d is invertible. So we obtain

$$\Delta_1|_{\text{Im}d} = d \Delta_0 d^{-1}.$$

Thus,

$$\sigma(\Delta_1|_{\text{Im}d}) = \sigma(\Delta_0). \quad \square$$

3.2. The 0-spectrum of Δ_0 and Δ_1

As the nonzero spectrum of Δ_0 and Δ_1 are the same, we are interested in characterizing the 0-spectrum. We give in the following an extension of a result of John Lott's [11] (Proposition 9, p. 12).

THEOREM 2. *Let (G, c) be a connected, locally finite and weighted infinite graph such that the weight on edges c is bounded, i.e., there exists a constant $\alpha > 0$ such that $\frac{1}{\alpha} \leq c(x, y) \leq \alpha$, for all $(x, y) \in \mathcal{E}$. Then*

$$0 \in \sigma(\Delta_1) \quad \text{or} \quad 0 \in \sigma(\Delta_0).$$

First, we start with preliminary results.

By [17] (page 44) and [9] (chapter 4) we have the next definition.

DEFINITION 3.1. The graph G verifies *the isoperimetric inequality* if there exists a constant $C > 0$ such that for all finite sub-graphs $G_U = (U, \mathcal{E}_U)$ of G , we have

$$|\partial \mathcal{E}_U| \geq C |U|,$$

where

$$|\partial \mathcal{E}_U| = \sum_{x \in U} \sum_{y \notin U} c(x, y) \quad \text{and} \quad |U| = \sum_{x \in U} \tilde{c}(x).$$

LEMMA 3.4. *If Δ_0 is invertible then the isoperimetric inequality holds.*

Proof. Let U a finite sub-graph of G . Let us set $g = \mathbf{1}_U$, meaning that $g(x) = 1$ if $x \in U$ and $g(x) = 0$ if $x \notin U$. Then we obtain

$$|U| = \sum_{x \in U} \tilde{c}(x) = \|g\|_{\mathcal{Y}}^2$$

and

$$|\partial \mathcal{E}_U| = \sum_{x \in U} \sum_{y \notin U} c(x, y) = \|\mathbf{d}g\|_{\mathcal{E}}^2.$$

By assumption 0 is not in the spectrum of Δ_0 . Then by the first assertion of Lemma 3.2, the operator \mathbf{d} is invertible, so there exists a positive constant λ so that

$$\|g\|_{\mathcal{Y}} \leq \lambda \|\mathbf{d}g\|_{\mathcal{E}}, \quad \forall g \in l^2(\mathcal{V}).$$

Thus, it follows that

$$|\partial \mathcal{E}_U| \geq C |U|, \quad \text{with} \quad C = \frac{1}{\lambda^2}. \quad \square$$

DEFINITION 3.2.

- A *branch* B is a finite sequence of vertices x_0, x_1, \dots, x_{m+1} such that for all j ; $1 \leq j \leq m$, we have $\deg(x_j) = 2$.
- The *length of a branch* B , denoted $\text{long}(B)$, is the number of vertices in this branch, here, $\text{long}(B) = m + 2$.
- The *interior of the branch* B is the set of vertices x_j of B satisfying the following conditions:
 - i) $\deg(x_j) = 2$.
 - ii) $\forall y \in \mathcal{V}$; $y \sim x_j \Rightarrow y \in B$.

See [5] and [19] for the definition of the interior set of vertices.

Instead of the argument of Lott [11] inspired by Gromov [10] (p. 236–237), we use the following lemma:



Figure 1: A branch of length $m + 2$

LEMMA 3.5. *We suppose that the following conditions are satisfied:*

- *The weight on edges c is bounded, i.e., there exists a constant $\alpha > 0$ such that $\frac{1}{\alpha} \leq c(x, y) \leq \alpha, \forall (x, y) \in \mathcal{E}$.*
- *The operator Δ_0 is invertible.*
- *The operator Δ_1 is injective.*

Then the graph (G, c) is a tree which contains branches with uniformly bounded lengths, that means $\exists M > 0, \forall B$ branch of $G, \text{long}(B) \leq M$.

Proof. On the one hand, the operator Δ_1 is injective which leads to the absence of cycles in the graph, so that G is a tree.

On the other hand, the operator Δ_0 is invertible, then the isoperimetric inequality is checked, by Lemma 3.4 there is a positive constant C such that for all finite sub-graphs U , we have

$$|\partial \mathcal{E}_U| \geq C|U|.$$

Let B be a branch with vertices $x_0, x_1, \dots, x_m, x_{m+1}$. We set $U = \{x_1, \dots, x_m\}$ the interior of the branch B , then

$$c(x_0, x_1) + c(x_m, x_{m+1}) \geq C \sum_{j=1}^m \tilde{c}(x_j). \tag{3.4}$$

For the sake of simplicity, we first prove the lemma for the case of the constant weight $c = 1$, before handling the case of general weights.

- *If $c = 1$, then we have $\tilde{c}(x) = \sum_{y \sim x} c(x, y) = \sum_{y \sim x} 1 = \text{deg}(x), \forall x \in \mathcal{V}$ (this is J. Lott’s case [11]). Therefore, the inequality (3.4) and Definition 3.2 gives*

$$2 \geq C \sum_{j=1}^m \text{deg}(x_j) = C \sum_{j=1}^m 2 = 2Cm.$$

We set $M := \frac{1}{C} + 2$ (independent of B), then $\text{long}(B) \leq M$. Thus the lengths of branches of G are uniformly bounded.

- *If $c \neq 1$ but c is bounded, that means there exists $\alpha > 0$ such that $\frac{1}{\alpha} \leq c(x, y) \leq \alpha$, for all $(x, y) \in \mathcal{E}$. And as we have the weight on the vertices is $\tilde{c}(x) = \sum_{y \sim x} c(x, y)$, we obtain that \tilde{c} is also bounded from below by $\frac{1}{\alpha}$.*

By the inequality (3.4) we have

$$2\alpha \geq c(x_0, x_1) + c(x_m, x_{m+1}) \geq C \sum_{j=1}^m \tilde{c}(x_j) \geq Cm \frac{1}{\alpha}.$$

Hence,

$$\frac{2\alpha^2}{C} \geq m.$$

We set $M = \frac{2\alpha^2}{C} + 2$ (independent of B), then $\text{long}(B) \leq M$. Thus, the lengths of the branches of G are uniformly bounded. \square

Now, we arrive to the proof of Theorem 2.

Proof. Taking the arguments from [11], we argue by contradiction. Suppose that both operators Δ_0 and Δ_1 are invertible. Then, by Lemma 3.5, the graph G is a tree which contains branches with uniformly bounded lengths; see Figure 2 for an example.




Figure 2: A branch tree

But the existence of such tree gives a δ -harmonic nonzero square-integral function φ . Indeed: we consider a part of the branch tree in Figure 2 as an example to simplify the understanding of the construction.

For the sake of simplicity, we first prove the theorem for the case of the constant weight $c = 1$, before handling the case of general weights.

First case: $c = 1$, we fix a vertex 0 as the origin of the tree and we set 0^- and 1 its different neighbors. Let us take

$$\varphi(0, 0^-) = \varphi(0, 1) = 1.$$

Then, we obtain $\delta\varphi(0) = 0$ (the tree is oriented).

Afterwards on the branch B_1 , φ is constant, in other words, $\varphi(j, j + 1) = 1$, for all j , such that $1 \leq j \leq 3$. And at the point 4, we have $\varphi(4, 5) = \varphi(4, 5^-) = \frac{1}{2}$. It is claimed that $\delta\varphi(4) = 0$. And for the points which are in the branch B_2 , the function φ is constant and takes the value $\frac{1}{2}$. And so on to the point 8, we have $\varphi(8, 9) = \varphi(8, 9^-) = \frac{1}{4}$, to obtain $\delta\varphi(8) = 0$. And for the points which are in the




Figure 3: An example of a branch tree




Figure 4: Another example of a branch tree

branch B_3 , the function φ is constant and takes the value $\frac{1}{4}$. And we continue in this way...

In a general way, G is a tree which contains branches with uniformly bounded lengths and we construct a functions φ in a part of G by selecting always two branches at bifurcation points and at all edges that occur on other branches, φ is set to zero, as in Figure 4. In the Figure 3, the construction of φ is done in the following way: on B_0 the function φ is constant and equals to 1. Then we add a generation, we get two branches $B_{1,1}$ and $B_{1,2}$ such that the function φ takes the value $\frac{1}{2}$. And to the generation m , we have $B_{m,k}$ branches, where $1 \leq k \leq 2^m$, then the function φ is equal to $\frac{1}{2^m}$. As a result, we show that this construction of φ is in $l^2(\mathcal{E})$. Using the fact that the lengths of the branches of the tree are uniformly bounded by a constant $M > 0$, we obtain

$$\begin{aligned} \|\varphi\|_{\mathcal{E}}^2 &= \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} \sum_{e \in B_{m,k}} (\varphi(e))^2 \\ &= \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} \sum_{e \in B_{m,k}} \left(\frac{1}{2^m}\right)^2 \\ &\leq \frac{1}{2} \sum_{m \geq 0} 2^m M \left(\frac{1}{2^m}\right)^2 \\ &= \frac{M}{2} \sum_{m \geq 0} \frac{1}{2^m} < \infty. \end{aligned}$$

Second case: $c \neq 1$ but c bounded by a positive constant. That means there exists $\alpha > 0$ such that $\frac{1}{\alpha} \leq c(x, y) \leq \alpha$, $\forall (x, y)$ in \mathcal{E} . As in Figure 2, on the branch B_1 , the vertex 0 has two neighbors denoted 0^- and 1. We want $\delta\varphi(0) = 0$, so we choose the function φ in the following way $\varphi(0, 1) = \frac{c(0, 0^-)}{c(0, 1)}\varphi(0, 0^-)$. And in the interior of B_1 , we set $\varphi(j, j+1) = \frac{c(0, 0^-)}{c(j, j+1)}\varphi(0, 0^-) \forall j, 1 \leq j \leq 3$. Next, we look at the point 4 which has two neighbors 5 and 5^- , to obtain $\delta\varphi(4) = 0$ and as we have $\varphi(3, 4) = \frac{c(0, 0^-)}{c(3, 4)}\varphi(0, 0^-)$. We choose $\varphi(4, 5) = \frac{c(0, 0^-)}{2c(4, 5)}\varphi(0, 0^-)$ and $\varphi(4, 5^-) = \frac{c(0, 0^-)}{2c(4, 5^-)}\varphi(0, 0^-)$. Therefore, in the interior of the branch B_2 ,

$$\varphi(j, j+1) = \frac{c(0, 0^-)}{2c(j, j+1)}\varphi(0, 0^-) \forall j, \quad 5 \leq j \leq 7.$$

And for the vertex 8, which has two neighbors 9 and 9^- . To have $\delta\varphi(8) = 0$ and by using that $\varphi(7, 8) = \frac{c(0, 0^-)}{2c(7, 8)}\varphi(0, 0^-)$. We choose $\varphi(8, 9) = \frac{c(0, 0^-)}{4c(8, 9)}\varphi(0, 0^-)$ and $\varphi(8, 9^-) = \frac{c(0, 0^-)}{4c(8, 9^-)}\varphi(0, 0^-)$. And in the interior of the branch B_3 ,

$$\varphi(j, j+1) = \frac{c(0, 0^-)}{4c(j, j+1)}\varphi(0, 0^-) \quad \text{for } j = 10.$$

And so on... In a general way, see Figure 3, on B_0 the function $\varphi(e_0) = \frac{c(0, 0^-)}{c(e_0)}\varphi(0, 0^-)$, where e_0 is an edge of B_0 . Then we add a generation, we get two branches $B_{1,1}$ and $B_{1,2}$ such that the function φ has a value $\varphi(e_1^k) = \frac{c(0, 0^-)}{2c(e_1^k)}\varphi(0, 0^-)$, where e_1^k denotes the edges of $B_{1,k}$ for $1 \leq k \leq 2$. And at generation m , we have $B_{m,k}$ branches, where $1 \leq k \leq 2^m$, then the function φ equals to $\varphi(e_m^k) = \frac{c(0, 0^-)}{2^m c(e_m^k)}\varphi(0, 0^-)$, where e_m^k denotes the edges of $B_{m,k}$. And to simplify the formulas, we can suppose that

$$\varphi(0, 0^-) = \frac{1}{c(0, 0^-)}.$$

Then, we obtain

$$\varphi(e_m^k) = \frac{1}{2^m c(e_m^k)}, \quad \forall m \geq 0 \quad \text{and} \quad 1 \leq k \leq 2^m.$$

Therefore, this construction gives $\varphi \in l^2(\mathcal{E})$. Using the fact that the lengths of the branches of the tree are uniformly bounded by a constant $M > 0$ and the weight c on the edges is bounded by a positive constant, we obtain

$$\begin{aligned} \|\varphi\|_{\mathcal{E}}^2 &= \frac{1}{2} \sum_e c(e) (\varphi(e))^2 \\ &\leq \sum_{m \geq 0} \sum_{k=1}^{2^m} \sum_{e \in B_{m,k}} c(e) (\varphi(e))^2 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{m \geq 0} \sum_{k=1}^{2^m} \sum_{e \in B_{m,k}} c(e) \left(\frac{1}{2^m c(e)} \right)^2 \\
 &= \sum_{m \geq 0} \sum_{k=1}^{2^m} \sum_{e \in B_{m,k}} \frac{1}{2^{2m} c(e)} \\
 &\leq \alpha M \sum_{m \geq 0} \frac{1}{2^m} \\
 &= 2M\alpha < \infty.
 \end{aligned}$$

Finally, we have φ in $l^2(\mathcal{E})$ and δ -harmonic. So, $0 \in \sigma(\Delta_1)$, which contradicts the assumption that 0 is not in the spectrum of Δ_1 . \square

REMARK 3.1. Any point of the graph can play the role of the first vertex 0 in the previous construction. It is then clear that we can construct an infinite family of independent functions φ which are in $l^2(\mathcal{E})$ and δ -harmonic. Then 0 is an eigenvalue of Δ_1 with infinite multiplicity, so $0 \in \sigma_{ess}(\Delta_1)$.

4. Examples

In this section, we will construct a δ -harmonic function φ in different examples of trees.

1) *Symmetric tree*: Following [7] we introduce the next definition:

DEFINITION 4.1. A tree T_s is symmetric around o with branching numbers $\{m_i\}_{i=0}^\infty$ if the degree of each vertex depends only on its distance from o , namely, for each $x \in T_s$, $\deg(x) = m_i$ if $d(o, x) = i$.

Example of a symmetric tree: We fix a vertex o as an origin of the tree. We set $S_n = \{x \in T_s; d(o, x) = n\}$. T_s is symmetric around o with branching numbers $\{m_n\}_{n=0}^\infty$. In Figure 4, we choose $m_n = 3 + n$ for all $n \in \mathbb{N}$ which is an increasing sequence. So, we have $m_0 = 3$ that means $\deg(o) = 3$. And for $x \in S_1$, we obtain $\deg(x) = m_1 = 4$. In the same way, if $x \in S_2$ we have $m_2 = 5$ and so on.

PROPOSITION 4.1. *If the symmetric tree T_s is simple (the edge weights are equal to 1) with $\deg(x) > 2$ for all $x \in T_s$, then there is a δ -harmonic function $\varphi \in l^2(\mathcal{E})$.*

Proof. We fix a vertex x_0 as an origin of the tree T_s , we can find an increasing sequence of finite subgraph $\{S_n\}_n$ such that $S_n = \{x \in T_s; d(x_0, x) = n\}$ and $T_s = \cup_n S_n$. By the definition of the symmetric tree, we have for all n $\deg(x_n) = m_n, \forall x_n \in S_n$. First, we construct a function φ so that $\delta\varphi = 0$ as follows: Let e_0 and b_0 denote two distinct outward edges connecting to the vertex x_0 . We define φ to be 0 excepted on these edges where $\varphi(e_0) = 1$ and $\varphi(b_0) = -1$ which gives $\delta\varphi(x_0) = 0$. And denote




Figure 5: Symmetric tree

$e_n^k, n \geq 1, 1 \leq k \leq \prod_{j=1}^n (m_j - 1)$, resp. $b_n^k, n \geq 1, 1 \leq k \leq \prod_{j=1}^n (m_j - 1)$, the outward edges emanating from e_0 , resp. b_0 , of generation n . We define

$$\varphi(e_n^k) = \frac{1}{\prod_{j=1}^n (m_j - 1)} \varphi(e_0),$$

$$\varphi(b_n^k) = \frac{1}{\prod_{j=1}^n (m_j - 1)} \varphi(b_0)$$

and φ takes value 0 on all edges other than e_n^k and b_n^k .

Second, through this construction, we look for $\varphi \in l^2(\mathcal{E})$. Using the fact that $\deg(x_n) = m_n \geq 3, \forall x_n \in S_n, \forall n$, we obtain

$$\begin{aligned} \|\varphi\|_{\mathcal{E}}^2 &= \frac{1}{2} \sum_{e \in \mathcal{E}} \varphi^2(e) \\ &= \frac{1}{2} \left(2 + \sum_{n \geq 1} \sum_{k=1}^{\prod_{j=1}^n (m_j - 1)} \varphi^2(e_n^k) + \varphi^2(b_n^k) \right) \\ &= 1 + \sum_{n \geq 1} \sum_{k=1}^{\prod_{j=1}^n (m_j - 1)} \left(\frac{1}{(m_1 - 1)(m_2 - 1) \dots (m_n - 1)} \right)^2 \\ &= 1 + \sum_{n \geq 1} \frac{1}{(m_1 - 1)(m_2 - 1) \dots (m_n - 1)} \\ &\leq 1 + \sum_{n \geq 1} \frac{1}{2^n} \\ &< \infty. \quad \square \end{aligned}$$

2) *Triadic tree with weights bounded from below:* As [2] (p. 19), we have the following definition of a triadic tree.

DEFINITION 4.2. A tree is a connected graph containing no cycles. The **triadic tree** is a tree such that all the vertices have degree 3.

PROPOSITION 4.2. *If the triadic tree has weights on the edges bounded from below by a positive constant λ , then there is a δ -harmonic function $\varphi \in l^2(\mathcal{E})$.*




Figure 6: Triadic tree

Proof. We fix a vertex o as the origin of the tree T . Define the increasing sequence of finite subgraphs $\{G_n\}_n$, $G_n = \{x \in \mathcal{V}; d(o, x) \leq n\}$ and let $G = \bigcup_n G_n$. Denote $S_n = \{x \in T; d(o, x) = n\}$.

We set x_1^1, x_1^2 and x_1^3 the different neighbors of o which are in S_1 . We suppose that $\varphi(o, x_1^i) = 0$ for all $i \in \{1, 2, 3\}$, so we have $\delta\varphi(o) = 0$.

We fix one vertex of S_1 for example $x_1 := x_1^1$, let e_0 and b_0 be the two outward edges of x_1 and define inductively $e_m^k, m \geq 1, 1 \leq k \leq 2^m$, resp. $b_m^k, m \geq 1, 1 \leq k \leq 2^m$, to be the outward edges emanating from e_0 , resp. b_0 , of generation m (the edge are oriented outward). For $m \geq 0$, we define φ to be 0 excepted on these edges where

$$\varphi(e_m^k) = \frac{1}{2^m} \frac{1}{c(e_m^k)}, \forall k; \quad 1 \leq k \leq 2^m$$

and

$$\varphi(b_m^k) = \frac{-1}{2^m} \frac{1}{c(b_m^k)}, \forall k; \quad 1 \leq k \leq 2^m.$$

With this construction, we obtain for each $x_n \in S_n$, $\delta\varphi(x_n) = 0, \forall n \geq 1$. Moreover, $\varphi \in l^2(\mathcal{E})$. Indeed: by using the assumption that the weights on the edges are bounded

from below by a positive constant λ , we obtain

$$\begin{aligned}
 \|\varphi\|_{\mathcal{E}}^2 &= \frac{1}{2} \sum_{e \in \mathcal{E}} c(e) \varphi^2(e) \\
 &= \frac{1}{2} \left(\sum_{m \geq 0} \sum_{k=1}^{2^m} c(e_m^k) \varphi^2(e_m^k) + c(b_m^k) \varphi^2(b_m^k) \right) \\
 &= \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} c(e_m^k) \frac{1}{2^{2m}} \frac{1}{c^2(e_m^k)} + \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} c(b_m^k) \frac{1}{2^{2m}} \frac{1}{c^2(b_m^k)} \\
 &= \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} \frac{1}{2^{2m}} \frac{1}{c(e_m^k)} + \frac{1}{2} \sum_{m \geq 0} \sum_{k=1}^{2^m} \frac{1}{2^{2m}} \frac{1}{c(b_m^k)} \\
 &\leq \lambda' \sum_{m \geq 0} \sum_{k=1}^{2^m} \frac{1}{2^{2m}} \\
 &= \lambda' \sum_{m \geq 0} 2^m \frac{1}{2^{2m}} \\
 &= \lambda' \sum_{m \geq 0} \frac{1}{2^m} \\
 &= 2\lambda',
 \end{aligned}$$

where $\lambda' = \frac{1}{\lambda}$. \square

REMARK 4.1.

- The construction of a δ -harmonic nonzero square-integral function depends on the edge weights.
- In the simple triadic tree, 0 is both in the spectrum of Δ_0 [4] and in the spectrum of Δ_1 [2].

Acknowledgements. I would like to express my sincere gratitude to my advisors Professors Colette Anné and Nabila Torki-Hamza for the continuous support of my Ph.D studies, for their patience, motivation, enthusiasm, and immense knowledge. I would like to thank the research unit (UR/13ES47) of Faculty of Sciences of Bizerte (University of Carthage) and the Laboratory of Mathematics Jean Leray of Nantes (LMJL) for their financial and continuous support. This work was financially supported by the “PHC Utique” program of the French Ministry of Foreign Affairs and Ministry of higher education and research and the Tunisian Ministry of higher education and scientific research in the CMCU project number 13G1501 “Graphes, Géométrie et théorie Spectrale”.

Finally, I thank my PhD’s referee: Professors Sylvain Golénia, Luc Hillairet and Aref Jeribi for their fruitful discussions, remarks and corrections to improve my PhD thesis from which this work is extracted.

I would like to thank also the anonymous referee for the careful reading of my paper and the valuable comments and suggestions.

REFERENCES

- [1] C. ANNÉ, N. TORKI-HAMZA, *The Gauss-Bonnet Operator of an Infinite Graph*, Analysis Mathematical Physics **5** (2015) 137–159.
- [2] H. AYADI, *Semi-Fredholmness of the discrete Gauss-Bonnet operator*, to appear in Filomat (2015).
- [3] H. BALOUDI, S. GOLÉNIA, A. JERIBI, *The Adjacency Matrix and the Discrete Laplacian Acting on Forms*, preprint arXiv:1505.06109(2015).
- [4] Y. COLIN DE VERDIÈRE, *Spectre de Graphes, Cours spécialisés*, volume **4** Société Mathématique de France (1998).
- [5] J. DODZIUK, *Elliptic operators on infinite graphs. Analysis, geometry and topology of elliptic operators*, World Sci. Publ., Hackensack NJ. (2006) 353–368.
- [6] D. E. DUTKAY, P. E. T. JORGENSEN, *Spectral theory for discrete Laplacians*, Complex analysis and operator theory **1** (2010) 1–38.
- [7] K. FUJIWARA, *The Laplacian on rapidly branching trees*, Duke mathematical journal **83** (1996) 191–202.
- [8] A. GRIGOR'YAN, *Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds*, Bulletin of the American Mathematical Society **36** (1999) 135–249.
- [9] A. GRIGOR'YAN, *Analysis on graphs*, Lecture Notes (2011/2012).
- [10] M. GROMOV, *Asymptotic Invariants of Infinite Groups*, Geometric Group Theory, vol. **2**, ed. by Graham Niblo and Martin Roller, London Math. Soc. Lecture Notes 182, Cambridge University Press, Cambridge (1993).
- [11] J. LOTT, *The zero-in-the spectrum question*, Enseign. Math **42** (1996) 341–376.
- [12] R. LYONS, Y. PERES, *Probability on Trees and Networks*, Cambridge University Press, New York (2016), available at <http://pages.iu.edu/~rdlyons/>.
- [13] B. MOHAR, W. WOESS, *A survey on spectra of infinite graphs*, Bull. London Math. Soc. **21**, (1989) 209–234.
- [14] M. REED, B. SIMON, *Methods of Modern Mathematical Physics I*, Academic Press (1980).
- [15] T. SHIRAI, *The spectrum of infinite regular line graphs*, Transactions of the American Mathematical Society **352** (2000) 115–132.
- [16] P. M. SOARDI, *Potential Theory on Infinite Networks*, Lecture Notes in Mathematics **1590** (1994).
- [17] P. M. SOARDI, W. WOESS, *Uniqueness of currents in infinite resistive networks*, Discrete Applied Mathematics **31** (1991) 37–49.
- [18] R. S. STRICHARTZ, *Transformation of spectra of graph laplacians*, Rocky Mountain Journal of Math, **40** (6) (2010).
- [19] N. TORKI-HAMZA, *Laplaciens de graphes infinis I-Graphes métriquement complets*, Confluentes Mathematici **2** (2010) 333–350, translated to Laplacians of infinite graphs. I: Metrically complete graphs, Arxiv:1201.4644.
- [20] W. WOESS, *Random walks on infinite graphs and groups*, Bulletin of the London Mathematical Society **26** (1994) 1–60.

(Received August 23, 2015)

Hèla Ayadi
 Unité de recherche Mathématiques et applications (UR/13ES47)
 à la faculté des sciences de Bizerte
 Laboratoire De Mathématiques Jean Leray, Université de Nantes
 e-mail: halaayadi@yahoo.fr, Hela.Ayadi@univ-nantes.fr