

## ON THE COMMUTATIVITY OF TOEPLITZ OPERATORS WITH HARMONIC SYMBOLS

HASHEM AL SABI AND ISSAM LOUHICHI

**Abstract.** In this paper we prove that if the polar decomposition of a symbol  $f$  is truncated above, i.e.,  $f(re^{i\theta}) = \sum_{k=-\infty}^N e^{ik\theta} f_k(r)$  where the  $f_k$ 's are radial functions, and if the associated Toeplitz operator  $T_f$  commutes with  $T_{z^2 + \bar{z}^2}$ , then  $T_f = Q(T_{z^2 + \bar{z}^2})$  where  $Q$  is a polynomial of degree at most 1.

*Mathematics subject classification (2010):* Primary 47B35, Secondary 47B38.

*Keywords and phrases:* Toeplitz operator, quasihomogeneous symbol, Mellin transform.

### REFERENCES

- [1] S. AXLER AND Ž. ČUČKOVIĆ, *Commuting Toeplitz operators with harmonic symbols*, Integral Equations and Operator Theory **14**, 1–12 (1991).
- [2] S. AXLER, Ž. ČUČKOVIĆ, AND N. V. RAO, *Commutants of analytic Toeplitz operators on the Bergman space*, Proc. Amer. Math. Soc. **128**, 1951–1953 (2000).
- [3] A. BROWN AND P. R. HALMOS, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. **123** (1964), 89–102.
- [4] H. HEDENMALM, B. KORENBLUM, AND K. ZHU, *Theory of Bergman spaces*, Springer Verlag, New York, 2000.
- [5] I. LOUHICHI, *Powers and roots of Toeplitz operators*, Proc. Amer. Math. Soc. **135**: 5 (2007), 1465–1475.
- [6] I. LOUHICHI, N. V. RAO, A. YOUSEF, *Two questions on the theory of Toeplitz operators on the Bergman space*, Complex Anal. Oper. Theory **3** (2009), no. 4, 881–889.
- [7] A. YOUSSEF, *Two Problems in the Theory of Toeplitz Operators on the Bergman Space*, 2009, Doctor of Philosophy, University of Toledo, Mathematics.