

SEMIGROUP GENERATIONS OF UNBOUNDED BLOCK OPERATOR MATRICES BASED ON THE SPACE DECOMPOSITION

JIE LIU, JUNJIE HUANG* AND ALATANCANG CHEN

(Communicated by I. M. Spitkovsky)

Abstract. This paper deals with the problem for unbounded block operator matrix

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

with natural domain to generate C_0 semigroups, based on the space decomposition. By describing the spectral inclusion relations between the numerical range of M and its inner entries, using the quadratic complements of M , some necessary and sufficient conditions for M to generate C_0 semigroups are given.

1. Introduction

The research of operator matrices is motivated by systems of linear evolution equations. It is well known that such systems are well-posed if and only if the corresponding operator matrix is the infinitesimal generator of a C_0 semigroup on underlying spaces^[1,2]. One usually concerns with conditions for the operator matrix $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ to generate C_0 semigroups, and obtain some conclusions^[3,4]. However, most of the results are discussed in the diagonal domain $\mathcal{D}(M) = \mathcal{D}(A) \oplus \mathcal{D}(D)$, by using the standard perturbation theorems. The change on the domain of the infinitesimal generator M has a great influence on its semigroup generation property. How can an unbounded operator matrix M with natural domain generates a semigroup? The problem is need to be discussed in other methods. In this paper, we consider the semigroup generation properties of the operator matrix M with natural domain $\mathcal{D}(M) = (\mathcal{D}(A) \cap \mathcal{D}(C)) \oplus (\mathcal{D}(B) \cap \mathcal{D}(D))$ in the different way.

As we know, the main obstacle for unbounded operators to generate semigroups is the unboundedness of their numerical range and the non-emptiness of their residual spectrum. Hence, we characterize the right boundedness of M with the quadratic numerical range of M , and consider the residual spectrum based on the space decomposition and quadratic complements.

Mathematics subject classification (2010): 47D03, 47B99, 37K99.

Keywords and phrases: Unbounded block operator matrix, generator, C_0 semigroup, space decomposition.

This work is supported by the National Natural Science Foundation of China (No. 11761029, 11761052, 11961022, 11961052), Natural Science Foundation of Inner Mongolia (No. 2018BS01005).

* Corresponding author.

2. Preliminaries

Let T be a linear operator between Hilbert spaces, and let \mathcal{Y} be a linear subspace of a Hilbert space. Then, the closure and orthogonal complement of \mathcal{Y} are denoted by $\overline{\mathcal{Y}}$ and \mathcal{Y}^\perp , respectively. Write $P_{\mathcal{Y}}$ for the orthogonal projection onto \mathcal{Y} along \mathcal{Y}^\perp (when \mathcal{Y} is closed) and $T|_{\mathcal{Y}}$ for the restriction of T to \mathcal{Y} . Also, we use $\mathcal{D}(T)$, $\mathcal{N}(T)$ and $\mathcal{R}(T)$ to denote the domain, nullspace and range of T , respectively. $n(T)$ is the dimension of $\mathcal{N}(T)$. Recall that T is said to be *dissipative*^[5], if the numerical range of T , i.e., $W(T) = \{(Tv, v) : v \in \mathcal{D}(T), \|v\| = 1\}$ is contained in the closed left half plane.

Throughout this paper, $\mathcal{X}_1, \mathcal{X}_2$ are always Hilbert spaces. In the product space $\mathcal{X}_1 \oplus \mathcal{X}_2$, we consider the unbounded closed block operator matrix

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \tag{2.1}$$

where $A : \mathcal{D}(A) \subset \mathcal{X}_1 \rightarrow \mathcal{X}_1$, $B : \mathcal{D}(B) \subset \mathcal{X}_2 \rightarrow \mathcal{X}_1$, $C : \mathcal{D}(C) \subset \mathcal{X}_1 \rightarrow \mathcal{X}_2$ and $D : \mathcal{D}(D) \subset \mathcal{X}_2 \rightarrow \mathcal{X}_2$ are densely defined closable operators. Write $\mathcal{D}_1 = \mathcal{D}(A) \cap \mathcal{D}(C)$, $\mathcal{D}_2 = \mathcal{D}(B) \cap \mathcal{D}(D)$. We suppose that the natural domain of M defined in (2.1), i.e.,

$$\mathcal{D}(M) = \mathcal{D}_1 \oplus \mathcal{D}_2 \tag{2.2}$$

is also dense in $\mathcal{X}_1 \oplus \mathcal{X}_2$. The followings are some constants

$$\begin{aligned} \alpha_0 &= \inf\{\operatorname{Re}\lambda : \lambda = (-Af, f), \|f\| = 1, f \in \mathcal{D}_1\}, \\ \beta_0 &= \inf\{\operatorname{Re}\lambda : \lambda = (-Dg, g), \|g\| = 1, g \in \mathcal{D}_2\}, \\ \delta_0 &= \min\{\alpha_0, \beta_0\}, \\ \gamma_0 &= \sup\{\operatorname{Re}\lambda : \lambda = \frac{(Cf, g) + (Bg, f)}{\|f\|^2 + \|g\|^2}, (f, g)^t \in \mathcal{D}(M)\}, \end{aligned} \tag{2.3}$$

where $\operatorname{Re}\lambda$ is the real part of the complex number λ .

DEFINITION 2.1. ^[6] Let M be the block operator matrix defined in (2.1)(2.2). For $f \in \mathcal{D}_1, g \in \mathcal{D}_2$ with $\|f\| = \|g\| = 1$, define the 2×2 matrix

$$M_{f,g} = \begin{pmatrix} (Af, f) & (Bg, f) \\ (Cf, g) & (Dg, g) \end{pmatrix}.$$

Then the set

$$W^2(M) = \bigcup_{\substack{f \in \mathcal{D}_1, g \in \mathcal{D}_2, \\ \|f\| = \|g\| = 1}} \sigma_p(M_{f,g})$$

is called the *quadratic numerical range* of M .

DEFINITION 2.2. ^[7] Let M be the block operator matrix defined in (2.1)(2.2). Suppose that either C or B is boundedly invertible. Then the quadratic operator polynomials T_1 and T_2 defined by

$$T_1(\lambda) = C - (D - \lambda)B^{-1}(A - \lambda) \text{ if } B \text{ is boundedly invertible,}$$

$$T_2(\lambda) = B - (A - \lambda)C^{-1}(D - \lambda) \text{ if } C \text{ is boundedly invertible,}$$

for $\lambda \in \mathbb{C}$ are called *quadratic complements* of M .

LEMMA 2.1. ^[8] *Let M be the block operator matrix defined in (2.1)(2.2). Then:*

- (i) $W^2(M) \subset W(M)$,
- (ii) $\dim \mathcal{X}_2 \geq 2 \implies W(\tilde{A}) \subset W^2(M)$,
- (iii) $\dim \mathcal{X}_1 \geq 2 \implies W(\tilde{D}) \subset W^2(M)$,

where $\tilde{A} = A|_{\mathcal{D}_1}, \tilde{D} = D|_{\mathcal{D}_2}$.

LEMMA 2.2. ^[5] *Let T be a closed linear operator in Hilbert space \mathcal{X} . Then for any $\lambda \notin \overline{W(T)}$, $\mathcal{N}(T - \lambda) = \{0\}$ and $\mathcal{R}(T - \lambda)$ is closed.*

LEMMA 2.3. ^[9] *Let T be a densely defined closed linear operator in a Hilbert space \mathcal{X} . Then:*

- (i) $\sigma(T) = \sigma_{app}(T) \cup \sigma_{r,1}(T)$,
- (ii) $\sigma_{app}(T) \subset \overline{W(T)}$,

where

$$\begin{aligned} \sigma_{r,1}(T) &= \{\lambda \in \mathbb{C} : T - \lambda \text{ is injective, } \overline{\mathcal{R}(T - \lambda)} \neq \mathcal{X} \text{ and } \mathcal{R}(T - \lambda) \text{ is closed}\}, \\ \sigma_{app}(T) &= \{\lambda \in \mathbb{C} : (T - \lambda)v_n \rightarrow 0, \{v_n\}_{n=1}^{+\infty} \subset \mathcal{D}(T), \|v_n\| = 1, n = 1, 2, \dots\}. \end{aligned}$$

PROPOSITION 2.1. *Let $\mathcal{X}_1, \mathcal{X}_2$ be infinite dimensional Hilbert spaces, and let M be the block operator matrix defined in (2.1)(2.2). Then the boundedness to the right with bound β ($\in \mathbb{R}$)^[10] of M , i.e., $\text{Re}(Mv, v) \leq \beta(v, v), v \in \mathcal{D}(M)$ implies those of \tilde{A} and \tilde{D} , where $\tilde{A} = A|_{\mathcal{D}_1}, \tilde{D} = D|_{\mathcal{D}_2}$.*

Proof. According to Lemma 2.1, it is easy to obtain that $W(\tilde{A}) \subset W(M)$ and $W(\tilde{D}) \subset W(M)$, since $\mathcal{X}_1, \mathcal{X}_2$ are infinite dimensional Hilbert spaces.

3. Main results

In what follows, we assume that the spaces $\mathcal{X}_1, \mathcal{X}_2$ defined in (2.1)(2.2) are infinite dimensional Hilbert spaces.

THEOREM 3.1. *Let M be the operator matrix defined in (2.1)(2.2), and let $\gamma_0 \leq \delta_0$ with γ_0, δ_0 defined as in (2.3). Write $\tilde{A} = A|_{\mathcal{D}_1}, \tilde{B} = B|_{\mathcal{D}_2}, \tilde{C} = C|_{\mathcal{D}_1}, \tilde{D} = D|_{\mathcal{D}_2}$. Suppose that \tilde{D} is a closed and $0 < n(\tilde{C}) < \infty$. Then M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$ if and only if the following statements hold:*

- (a) \tilde{A} and \tilde{D} are bounded to the right with bound β ,
- (b) $\mathcal{R}((A_1 - \lambda T_2(\lambda))) = \mathcal{X}_1$ and $\mathcal{R}(C_{11}) = \mathcal{R}(\tilde{D} - \lambda)^\perp$ for $\lambda > \beta$, where $(A_1 - \lambda T_2(\lambda))$ is a line operator, $A_1 = A|_{\mathcal{N}(\tilde{C})}$, $T_2(\lambda) = A_{22} - \lambda - \tilde{B}(\tilde{D} - \lambda)^{-1}C_{22}$, $A_{22} = A|_{\mathcal{N}(C_1)}$, $C_1 = P_{\mathcal{R}(\tilde{D}-\lambda)^\perp}C|_{\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1}$, $C_{11} = P_{\mathcal{R}(\tilde{D}-\lambda)^\perp}C|_{(\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \cap \mathcal{D}_1}$ and $C_{22} = P_{\mathcal{R}(\tilde{D}-\lambda)}C|_{\mathcal{N}(C_1)}$.

Proof. The assertion (a) implies $W(-\tilde{A}) \subset \{z \in \mathbb{C} : \operatorname{Re}z \geq -\beta\}$ and $W(-\tilde{D}) \subset \{z \in \mathbb{C} : \operatorname{Re}z \geq -\beta\}$, so α_0, β_0 are well defined. For each $v = (f g)^t \in \mathcal{D}(M)$ with $f \neq 0$ and $g \neq 0$, let $\tau = \frac{\|f\|^2}{\|f\|^2 + \|g\|^2}$. It follows from $\gamma_0 \leq \delta_0$ that

$$\begin{aligned} \frac{\operatorname{Re}(Cf, g) + \operatorname{Re}(Bg, f)}{\|f\|^2 + \|g\|^2} &\leq \gamma_0 \leq \delta_0 \leq \tau\alpha_0 + (1 - \tau)\beta_0 \\ &\leq \tau(\alpha_0 + \beta) + (1 - \tau)(\beta_0 + \beta) \\ &\leq \tau \cdot \frac{\operatorname{Re}(-Af, f) + \beta(f, f)}{\|f\|^2} + (1 - \tau) \cdot \frac{\operatorname{Re}(-Dg, g) + \beta(g, g)}{\|g\|^2} \\ &= \frac{\operatorname{Re}(-Af, f) + \beta(f, f)}{\|f\|^2 + \|g\|^2} + \frac{\operatorname{Re}(-Dg, g) + \beta(g, g)}{\|f\|^2 + \|g\|^2}. \end{aligned} \tag{3.1}$$

Hence

$$\operatorname{Re}(Cf, g) + \operatorname{Re}(Bg, f) + \operatorname{Re}(Af, f) + \operatorname{Re}(Dg, g) \leq \beta(f, f) + \beta(g, g),$$

and

$$\begin{aligned} \operatorname{Re}(Mv, v) &= \operatorname{Re} \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix}, \begin{pmatrix} f \\ g \end{pmatrix} \right) \\ &= \operatorname{Re}(Cf, g) + \operatorname{Re}(Bg, f) + \operatorname{Re}(Af, f) + \operatorname{Re}(Dg, g) \\ &\leq \beta(v, v). \end{aligned}$$

If $f = 0$ or $g = 0$, it is easy to prove $\operatorname{Re}(Mv, v) \leq \beta(v, v)$. Thus, M is bounded to the right with bound β .

To complete the proof, it suffices to show that $M - \lambda$ ($\lambda > \beta$) is surjective.

Since $0 < n(\tilde{C}) < \infty$, $\mathcal{N}(\tilde{C})$ is a closed subspace of \mathcal{X}_1 . From $\mathcal{N}(\tilde{C}) \subset \mathcal{D}(\tilde{C}) = \mathcal{D}_1$, we know

$$\mathcal{D}(\tilde{C}) = \mathcal{N}(\tilde{C}) \oplus (\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1).$$

On the other hand, from assertion (i) follows that $\lambda \notin \overline{W(\tilde{A}) \cup W(\tilde{D})}$, and $\mathcal{R}(\tilde{D} - \lambda)$ is a closed subspace of \mathcal{X}_2 by Lemma 2.2. Then, as an operator from $\mathcal{N}(\tilde{C}) \oplus (\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1) \oplus \mathcal{D}_2 \subset \mathcal{N}(\tilde{C}) \oplus \mathcal{N}(\tilde{C})^\perp \oplus \mathcal{X}_2$ to $\mathcal{X}_1 \oplus \mathcal{R}(\tilde{D} - \lambda)^\perp \oplus \mathcal{R}(\tilde{D} - \lambda)$, $M - \lambda$ admits the following block representation

$$M - \lambda = \begin{pmatrix} A_1 - \lambda & A_2 - \lambda & \tilde{B} \\ 0 & C_1 & 0 \\ 0 & C_2 & \tilde{D} - \lambda \end{pmatrix}. \tag{3.2}$$

Here $A_2 = A|_{\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1}$ and $C_2 = P_{\mathcal{R}(\tilde{D}-\lambda)}C|_{\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1}$. From assertion (i) follows $\lambda \notin \overline{W(\tilde{A})} \cup \overline{W(\tilde{D})}$, and hence $A_1 - \lambda : \mathcal{N}(\tilde{C}) \rightarrow \mathcal{X}_1$ is injective and $\tilde{D} - \lambda : \mathcal{D}_2 \rightarrow \mathcal{R}(\tilde{D} - \lambda)$ is boundedly invertible by Lemma 2.2. It follows from $0 < n(\tilde{C}) < \infty$ that $0 < n(C_1) < \infty$, hence

$$\mathcal{D}(C_1) = \mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1 = ((\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \cap \mathcal{D}_1) \oplus \mathcal{N}(C_1).$$

As an operator from $\mathcal{N}(\tilde{C}) \oplus ((\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \cap \mathcal{D}_1) \oplus \mathcal{N}(C_1) \oplus \mathcal{D}_2 \subset \mathcal{N}(\tilde{C}) \oplus (\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \oplus \mathcal{N}(C_1) \oplus \mathcal{X}_2$ to $\mathcal{X}_1 \oplus \mathcal{R}(\tilde{D} - \lambda)^\perp \oplus \mathcal{R}(\tilde{D} - \lambda)$, $M - \lambda$ has the following representation

$$M - \lambda = \begin{pmatrix} A_1 - \lambda & A_{21} - \lambda & A_{22} - \lambda & \tilde{B} \\ 0 & C_{11} & 0 & 0 \\ 0 & C_{21} & C_{22} & \tilde{D} - \lambda \end{pmatrix}.$$

Clearly, $C_{11} : (\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \cap \mathcal{D}_1 \rightarrow \mathcal{R}(\tilde{D} - \lambda)^\perp$ is injective, and hence C_{11} is left invertible, i.e., there exist C_{11}^{-1} such that $C_{11}^{-1}C_{11} = I$ holds. Hence, $M - \lambda$ ($\lambda > 0$) has the following transformation

$$E_1(M - \lambda) = N,$$

where

$$E_1 = \begin{pmatrix} I & \tilde{B}(\tilde{D} - \lambda)^{-1}C_{21}C_{11}^{-1} - (A_{21} - \lambda)C_{11}^{-1} & -\tilde{B}(\tilde{D} - \lambda)^{-1} \\ 0 & I & 0 \\ 0 & -C_{21}C_{11}^{-1} & I \end{pmatrix},$$

$$N = \begin{pmatrix} A_1 - \lambda & 0 & A_{22} - \lambda - \tilde{B}(\tilde{D} - \lambda)^{-1}C_{22} & 0 \\ 0 & C_{11} & 0 & 0 \\ 0 & 0 & C_{22} & \tilde{D} - \lambda \end{pmatrix}.$$

According to the relative boundedness of unbounded operators([7, P92]), operator $\tilde{B}(\tilde{D} - \lambda)^{-1}$, $C_{21}C_{11}^{-1}$ and $(A_{21} - \lambda)C_{11}^{-1}$ are bounded operator, hence E_1 is bijective. Thus, $M - \lambda$ is surjective if and only if so is N . Since $\tilde{D} - \lambda$ is boundedly invertible, N has a further transformation that

$$NE_2 = L,$$

where

$$E_2 = \begin{pmatrix} I & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & -(\tilde{D} - \lambda)^{-1}C_{22} & I \end{pmatrix},$$

$$L = \begin{pmatrix} A_1 - \lambda & 0 & A_{22} - \lambda - \tilde{B}(\tilde{D} - \lambda)^{-1}C_{22} & 0 \\ 0 & C_{11} & 0 & 0 \\ 0 & 0 & 0 & \tilde{D} - \lambda \end{pmatrix}.$$

Thus, N is surjective if and only if so is L . It is clear that the condition (b) implies L is surjective, so $M - \lambda$ is surjective. Therefore, M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$.

Conversely, if M is the infinitesimal generator of a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$, then M is bounded to the right with bound β and $\lambda \in \rho(M)$ for $\lambda > \beta$. By Proposition 2.1, \tilde{A} and \tilde{D} are both bounded to the right with bound β , i.e., the assertion (a) holds. According to the proof in sufficiency, $M - \lambda$, as an operator from $\mathcal{N}(\tilde{C}) \oplus ((\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \cap \mathcal{D}_1) \oplus \mathcal{N}(C_1) \oplus \mathcal{D}_2 \subset \mathcal{N}(\tilde{C}) \oplus (\mathcal{N}(\tilde{C})^\perp \ominus \mathcal{N}(C_1)) \oplus \mathcal{N}(C_1) \oplus \mathcal{X}_2$ to $\mathcal{X}_1 \oplus \mathcal{R}(\tilde{D} - \lambda)^\perp \oplus \mathcal{R}(\tilde{D} - \lambda)$, is surjective. Hence, the assertion (b) holds immediately.

THEOREM 3.2. *Let M be the operator matrix defined in (2.1)(2.2), and let $\gamma_0 \leq \delta_0$ with γ_0, δ_0 defined as in (2.3). Write $\tilde{A} = A|_{\mathcal{D}_1}$, $\tilde{B} = B|_{\mathcal{D}_2}$, $\tilde{C} = C|_{\mathcal{D}_1}$, $\tilde{D} = D|_{\mathcal{D}_2}$. Suppose that \tilde{A} is a closed and $0 < n(\tilde{B}) < \infty$, then M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$ if and only if the following statements hold:*

- (i) \tilde{A} and \tilde{D} are bounded to the right with bound β ,
- (ii) $\mathcal{R}((D_1 - \lambda \ T_1(\lambda))) = \mathcal{X}_2$ and $\mathcal{R}(B_{11}) = \mathcal{R}(\tilde{A} - \lambda)^\perp$ for $\lambda > \beta$, where $(D_1 - \lambda \ T_1(\lambda))$ is a line operator, $D_1 = D|_{\mathcal{N}(\tilde{B})}$, $T_1(\lambda) = D_{22} - \lambda - \tilde{C}(\tilde{A} - \lambda)^{-1}B_{22}$, $D_{22} = D|_{\mathcal{N}(B_1)}$, $B_1 = P_{\mathcal{R}(\tilde{A} - \lambda)^\perp}B|_{\mathcal{N}(\tilde{B})^\perp \cap \mathcal{D}_2}$ and $B_{22} = P_{\mathcal{R}(\tilde{A} - \lambda)}B|_{\mathcal{N}(B_1)}$.

Proof. The proof is similar to that of Theorem 3.1. We only need to note that the operator $M - \lambda$ ($\lambda > \beta$), discussed from $\mathcal{X}_1 \oplus \mathcal{N}(\tilde{B}) \oplus (\mathcal{N}(\tilde{B})^\perp \ominus \mathcal{N}(B_1)) \oplus \mathcal{N}(B_1)$ to $\mathcal{R}(\tilde{A} - \lambda)^\perp \oplus \mathcal{R}(\tilde{A} - \lambda) \oplus \mathcal{X}_2$ has the representation

$$M - \lambda = \begin{pmatrix} 0 & 0 & B_{11} & 0 \\ \tilde{A} - \lambda & 0 & B_{21} & B_{22} \\ \tilde{C} & D_1 - \lambda & D_{21} - \lambda & D_{22} - \lambda \end{pmatrix}.$$

Under more finer subdivision of the domain space, we may have the following results.

THEOREM 3.3. *Let M be the operator matrix defined in (2.1)(2.2), and let $\gamma_0 \leq \delta_0$ with γ_0, δ_0 defined as in (2.3). Write $\tilde{A} = A|_{\mathcal{D}_1}$, $\tilde{B} = B|_{\mathcal{D}_2}$, $\tilde{C} = C|_{\mathcal{D}_1}$, $\tilde{D} = D|_{\mathcal{D}_2}$. Suppose that \tilde{A} and \tilde{D} are closed, $0 < n(\tilde{B}) < \infty$ and $0 < n(\tilde{C}) < \infty$. Then M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$ if and only if the following statements hold:*

- (i) \tilde{A} and \tilde{D} are bounded to the right with bound β ,
- (ii) $\mathcal{R}((A_{22} - \lambda \ B_{22})) = \mathcal{R}(A_2 - \lambda)$, $\mathcal{R}((C_{22} \ D_{22} - \lambda)) = \mathcal{R}(D_2 - \lambda)$, $\mathcal{R}(B_{11}) = \mathcal{R}(\tilde{A} - \lambda)^\perp$, $\mathcal{R}(C_{11}) = \mathcal{R}(\tilde{D} - \lambda)^\perp$ for $\lambda > \beta$, where $A_2 = A|_{\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1}$, $D_2 = D|_{\mathcal{N}(\tilde{B})^\perp \cap \mathcal{D}_2}$, $C_1 = P_{\mathcal{R}(\tilde{D} - \lambda)^\perp}C|_{\mathcal{N}(\tilde{C})^\perp \cap \mathcal{D}_1}$, $A_{22} = A|_{\mathcal{N}(C_1)}$, $D_{22} = D|_{\mathcal{N}(B_1)}$, $B_1 = P_{\mathcal{R}(\tilde{A} - \lambda)^\perp}B|_{\mathcal{N}(\tilde{B})^\perp \cap \mathcal{D}_2}$, $B_{22} = P_{\mathcal{R}(\tilde{A} - \lambda)}B|_{\mathcal{N}(B_1)}$ and $C_{22} = P_{\mathcal{R}(\tilde{D} - \lambda)}C|_{\mathcal{N}(C_1)}$.

Proof. The proof is similar to that of Theorem 3.1. We only need to note that the operator $M - \lambda$ ($\lambda > \beta$), discussed from $\mathcal{N}(\tilde{C}) \oplus (\mathcal{N}(\tilde{C})^\perp \oplus \mathcal{N}(C_1)) \oplus \mathcal{N}(C_1) \oplus \mathcal{N}(\tilde{B}) \oplus (\mathcal{N}(\tilde{B})^\perp \oplus \mathcal{N}(B_1)) \oplus \mathcal{N}(B_1)$ to $\mathcal{R}(\tilde{A} - \lambda) \oplus \mathcal{R}(\tilde{D} - \lambda) \oplus \mathcal{R}(\tilde{D} - \lambda)^\perp \oplus \mathcal{R}(\tilde{A} - \lambda)^\perp$ has the representation

$$M - \lambda = \begin{pmatrix} A_1 - \lambda & A_{21} - \lambda & A_{22} - \lambda & 0 & B_{21} & B_{22} \\ 0 & C_{21} & C_{22} & D_1 - \lambda & D_{21} - \lambda & D_{22} - \lambda \\ 0 & C_{11} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & B_{11} & 0 \end{pmatrix}.$$

THEOREM 3.4. *Let M be the operator matrix defined in (2.1) with the dense domain*

$$\mathcal{D}(M) = \mathcal{D}(C) \oplus (\mathcal{D}(B) \cap \mathcal{D}(D)) := \mathcal{D}(C) \oplus \mathcal{D}_2,$$

and let $\gamma_0 \leq \delta_0$ with γ_0, δ_0 defined as in (2.3). Write $\tilde{A} = A|_{\mathcal{D}(C)}, \tilde{D} = D|_{\mathcal{D}_2}$. If C is boundedly invertible and $C^{-1}D$ is bounded on $\mathcal{D}(D)$, then M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t}$ ($t \geq 0$) for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$ if and only if the following statements hold:

- (i) \tilde{A} and \tilde{D} are bounded to the right with bound β ,
- (ii) $\sigma_{r,1}(T_2) \cap \{z \in \mathbb{C} : \text{Re} z > \beta\} = \emptyset$, where $T_2(\lambda) = B - (A - \lambda)C^{-1}(D - \lambda)$ for $\lambda \in \mathbb{C}$.

Proof. Combining (i) and assumption $\gamma_0 \leq \delta_0$, we see that M is bounded to the right with bound β . Since C is boundedly invertible, $M - \lambda$ ($\lambda \in \mathbb{C}$) has the following factorization

$$M - \lambda = \begin{pmatrix} I & (A - \lambda)C^{-1} \\ 0 & I \end{pmatrix} \begin{pmatrix} 0 & T_2(\lambda) \\ C & 0 \end{pmatrix} \begin{pmatrix} I & C^{-1}(D - \lambda) \\ 0 & I \end{pmatrix}.$$

As $\mathcal{D}(C) \subset \mathcal{D}(A)$ and $C^{-1}D$ is bounded on $\mathcal{D}(D)$, $\begin{pmatrix} I & (A - \lambda)C^{-1} \\ 0 & I \end{pmatrix}$ and $\begin{pmatrix} I & C^{-1}(D - \lambda) \\ 0 & I \end{pmatrix}$ are both bounded and invertible, and hence

$$\lambda \in \sigma_{r,1}(M) \iff \lambda \in \sigma_{r,1}(T_2). \tag{3.3}$$

By Lemma 2.3, we have

$$\sigma_{app}(M) \subset \overline{W(M)} \subset \{z \in \mathbb{C} : \text{Re} z \leq \beta\}$$

since M is bounded to the right with bound β . Hence, it follows from assertion (ii) that

$$\sigma(M) \subset \{z \in \mathbb{C} : \text{Re} z \leq \beta\},$$

and hence

$$\{z \in \mathbb{C} : \text{Re} z > \beta\} \subset \rho(M). \tag{3.4}$$

Thus, M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t} (t \geq 0)$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$.

Conversely, suppose that M is the infinitesimal generator of a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t} (t \geq 0)$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$. Then, we may have that (3.4) is valid, and that \tilde{A} and \tilde{D} are both bounded to the right with bound β . Since C is boundedly invertible and $C^{-1}D$ is bounded on $\mathcal{D}(D)$, (3.3) holds. Therefore, the assertion (ii) follows.

In the same way, we have the following symmetric result.

THEOREM 3.5. *Let M be the operator matrix defined in (2.1) with dense domain*

$$\mathcal{D}(M) = (\mathcal{D}(A) \cap \mathcal{D}(C)) \oplus \mathcal{D}(B) := \mathcal{D}_1 \oplus \mathcal{D}(B),$$

and let $\gamma_0 \leq \delta_0$ with γ_0, δ_0 defined as in (2.3). Write $\tilde{A} = A|_{\mathcal{D}_1}, \tilde{D} = D|_{\mathcal{D}(B)}$. If B is boundedly invertible and $B^{-1}A$ is bounded on $\mathcal{D}(A)$, then M generates a C_0 semigroup $(T(t))_{t \geq 0}$ with $\|T(t)\| \leq e^{\beta t} (t \geq 0)$ for some $\beta \geq 0$ on $\mathcal{X}_1 \oplus \mathcal{X}_2$ if and only if the following statements hold:

(i) \tilde{A} and \tilde{D} are bounded to the right with bound β ,

(ii) $\sigma_{r,1}(T_1) \cap \{z \in \mathbb{C} : \operatorname{Re} z > \beta\} = \emptyset$, where $T_1(\lambda) = C - (D - \lambda)B^{-1}(A - \lambda)$ for $\lambda \in \mathbb{C}$.

Proof. The proof is similar to that of Theorem 3.4. Note that $M - \lambda$ ($\lambda \in \mathbb{C}$) has the following factorization

$$M - \lambda = \begin{pmatrix} I & 0 \\ (D - \lambda)B^{-1} & I \end{pmatrix} \begin{pmatrix} 0 & B \\ T_1(\lambda) & 0 \end{pmatrix} \begin{pmatrix} I & 0 \\ B^{-1}(A - \lambda) & I \end{pmatrix}$$

as B is boundedly invertible. It follows from $\mathcal{D}(B) \subset \mathcal{D}(D)$ and $B^{-1}A$ is bounded on $\mathcal{D}(A)$ that $\begin{pmatrix} I & 0 \\ (D - \lambda)B^{-1} & I \end{pmatrix}$ and $\begin{pmatrix} I & 0 \\ B^{-1}(A - \lambda) & I \end{pmatrix}$ are both bounded and invertible, and hence

$$\lambda \in \sigma_{r,1}(M) \iff \lambda \in \sigma_{r,1}(T_1).$$

4. Example

Consider the mixed problem of partial differential equation

$$\begin{cases} \frac{\partial^4 u(x,y)}{\partial x^4} + \frac{\partial^2 u(x,y)}{\partial y^2} + 2 \frac{\partial u(x,y)}{\partial y} + u = 0, & 0 < x < 1, 0 < y < h, \\ u(0,y) = u(1,y) = 0, u_x''(0,y) = u_x''(1,y) = 0, & 0 \leq y \leq h, \\ u(x,0) = \varphi(x), u_y'(x,0) = \psi(x), & 0 \leq x \leq 1. \end{cases} \tag{4.1}$$

Let $\mathcal{X} = L^2[0, 1]$, $p = \frac{\partial^2 u}{\partial x^2}$, $q = -\frac{\partial u}{\partial y} - u$, then problem (4.1) can be transformed into abstract Cauchy problem in $\mathcal{X} \oplus \mathcal{X}$ as follows

$$\frac{\partial}{\partial y} \begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} -1 & -\frac{\partial^2}{\partial x^2} \\ \frac{\partial^2}{\partial x^2} & -1 \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix}.$$

The corresponding block operator matrix is

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} -I & -\frac{\partial^2}{\partial x^2} \\ \frac{\partial^2}{\partial x^2} & -I \end{pmatrix} : \mathcal{D}(M) = \mathcal{D}(C) \oplus \mathcal{D}(B) \rightarrow \mathcal{X} \oplus \mathcal{X},$$

and

$$\mathcal{D}(B) = \mathcal{D}(C) = \{v(x) \in \mathcal{X} : v(x)', v(x)'' \in \mathcal{X}, v(0) = v(1) = 0\}.$$

It is easy to see that $\gamma_0 = 0 \leq \delta_0$ and $0 \in \rho(B)$. Clearly, A, D are dissipative and

$$T_1(\lambda) := C - (D - \lambda)B^{-1}(A - \lambda) = \frac{\partial^2}{\partial x^2} + (\lambda + 1)^2 \left(\frac{\partial^2}{\partial x^2}\right)^{-1}, \lambda \in \mathbb{C}.$$

It follows from $0 \in \rho(B) \cap \rho(C)$ that $\mathcal{R}(T_1(\lambda)) = \mathcal{X}$, and hence $\sigma_r(T_1) = \emptyset$. Thus, M satisfies the assertions of Theorem 3.5 and hence M generates a contraction semigroup on $\mathcal{X} \oplus \mathcal{X}$.

REFERENCES

[1] GOLDSTEIN J. A., *Semigroups of Linear Operators and Applications*, Oxford University Press, New York, 1985.
 [2] ENGEL K. J., NAGEL R., *One Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics 194, Springer-Verlag, New York, 2000.
 [3] NAGEL R., *Towards a “matrix theory” for unbounded operator matrices*, Math. Z, 1989, 201, 57–68.
 [4] MUGNOLO D., *Asymptotics of semigroups generated by operator matrices*, Arab. J. Math. 2014, 3: 419–435.
 [5] KATO T., *Perturbation Theory for Linear Operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
 [6] LANGER H., MARKUS A. S., MATSAEV V. I., TRETTER C., *A new concept for block operator matrices: the quadratic numerical range*, Linear Algebra Appl., 330 (1-3), 2001, 89–112.
 [7] TRETTER C., *Spectral Theory of Block Operator Matrices and Applications*, Imperial College Press, London, 2008.
 [8] TRETTER C., *Spectral inclusion for unbounded block operator matrices*, Journal of Functional Analysis 256, 2009, 3806–3829.

- [9] WU D., CHEN A., *Spectral inclusion properties of the numerical range in a space with an indefinite metric*, *Linear Algebra Appl.* 2011, 435: 1131–1136.
- [10] KREIN S. G., *Linear Differential Equations in Banach Space*, American Mathematical Society, Providence, Rhode Island, 1971.

(Received January 14, 2018)

Jie Liu
School of Mathematics and Big Data
Hohhot Minzu College
Hohhot 010051, P.R. China
School of Mathematical Sciences
Inner Mongolia University
Hohhot 010021, P.R. China
e-mail: liujieem@163.com

Junjie Huang
School of Mathematical Sciences
Inner Mongolia University
Hohhot 010021, P.R. China
e-mail: huangjunjie@imu.edu.cn

Alatancang Chen
School of Mathematics and Big Data
Hohhot Minzu College
Hohhot 010051, P.R. China
e-mail: alatanca@imu.edu.cn