

THE ALGEBRA GENERATED BY SIMPLE ELEMENTS OF A MATRIX CENTRALIZER

RALPH JOHN DE LA CRUZ

Abstract. Let $\mathcal{C}(S)$ denote the centralizer of an arbitrary square matrix S . An element $A \in \mathcal{C}(S)$ is simple if $A - I$ is of rank 1. Let \mathcal{A}_S denote the subalgebra generated by the simple elements of $\mathcal{C}(S)$. We use the Weyr canonical form to describe the subalgebra \mathcal{A}_S , and we show that if $\lambda_1, \dots, \lambda_k$ are the distinct eigenvalues of S , and l is the number of defective eigenvalues of S , then \mathcal{A}_S is of dimension $l + \sum_{i=1}^k \text{nullity}(S - \lambda_i I)^2$.

Mathematics subject classification (2020): 15A03, 15A27.

Keywords and phrases: Simple, Weyr structure, Weyr canonical form, centralizer, defective.

REFERENCES

- [1] T. AGAPITO, A. T. PARAS, ψ_S -orthogonal matrices and ψ_S -symmetries, Linear Algebra Appl. **584** (2020) 185–196.
- [2] R. J. DE LA CRUZ, K. DELA ROSA, D. I. MERINO, A. T. PARAS, The Cartan-Diedonne-Scherk Theorems for complex S -orthogonal matrices, Linear Algebra Appl. **458** (2014) 251–260.
- [3] M. CATRAL, L. LEBTAHI, J. STUART, N. THOME, On a matrix group constructed from an $\{R, s + 1, k\}$ -potent matrix, Linear Algebra Appl. **461** (2014) 200–210.
- [4] R. J. DE LA CRUZ, D. I. MERINO, A. T. PARAS, S orthogonal matrices and S symmetries, Linear Algebra Appl. **474** (2015) 213–229.
- [5] J. DIEUDONNE, Sur les generateurs des groupes classiques, Summa Bras. Math. **3** (1955) 149–178.
- [6] R. A. HORN AND C. R. JOHNSON, Topics in Matrix Analysis, Cambridge University Press, New York, 1991.
- [7] G. HUANG, F. YIN, Constrained inverse eigenproblem and associated approximation problem for anti-Hermitian R -symmetric matrices, Applied Mathematics and Computation **186** (2007) 426–434.
- [8] G. HUANG, F. YIN, Matrix inverse problem and its optimal approximation problem for R -symmetric matrices, Applied Mathematics and Computation **189** (2007) 482–489.
- [9] K. O'MEARA, J. CLARK, C. VINSONHALER, Advanced topics in linear algebra: weaving matrix problems through the Weyr form, Oxford University Press, Oxford 2011.
- [10] H. RADJAVI, Decomposition of matrices into simple involutions, Linear Algebra Appl., **12** (1975) 247–255.
- [11] P. SCHERK, On the decomposition of orthogonalities into symmetries, Proc. Amer. Math. Soc. **1** (1950) 481–491.
- [12] W. TRENCH, Characterization and properties of matrices with generalized symmetry or skew symmetry, Linear Algebra Appl. **377** (2004) 207–218.
- [13] W. TRENCH, Minimization problems for (R, S) -symmetric and (R, S) -skew symmetric matrices, Linear Algebra Appl. **389** (2004) 23–31.
- [14] F. UHLIG, Constructive ways for generating (generalized) real orthogonal matrices as products of (generalized) symmetries, Linear Algebra Appl. **332–334** (2001) 459–467.