

THE SPECTRUM OF q -CESÀRO MATRICES ON c AND ITS VARIOUS SPECTRAL DECOMPOSITION FOR $0 < q < 1$

NUH DURNA AND MERVE ESRA TÜRKAY

(Communicated by B. Jacob)

Abstract. One of q -analogs of the Cesàro matrix of order one is the triangular matrix with nonzero entries $c_{nk} = \frac{q^{n-k}}{1+q+\dots+q^n}$, $0 \leq k \leq n$, where $q \in [0, 1]$. In this article, we will determine the spectrum of this matrix on the space of convergent sequences c . We will also obtain the fine spectral decomposition in the sense of Goldberg and a non-discrete spectral decomposition of the obtained spectrum.

1. Introduction

This section is devoted firstly to make a brief introduction to symbols of q -mathematics and q -Cesàro matrices. The subject of q -mathematics has many applications in mathematics, and the beginnings of q -mathematics date back to time of Euler. The q -analogue of the integer n , is defined by

$$[n]_q = \frac{1 - q^n}{1 - q} \quad (q \neq 1). \tag{1.1}$$

Then, the q -analogue of the factorial, i.e q -factorial, is defined as

$$[n]_q! = \begin{cases} \frac{q-1}{q-1} \frac{q^2-1}{q-1} \dots \frac{q^n-1}{q-1}, & n = 1, 2, \dots \\ 1, & n = 0. \end{cases} \tag{1.2}$$

and then q -binomial coefficients are defined as

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[n-k]_q! [k]_q!}. \tag{1.3}$$

Note that, as $q \rightarrow 1$, the q -binomial coefficients approach the usual binomial coefficients.

Mathematics subject classification (2020): Primary 40H05, 40C99; Secondary 46A35, 47A10.

Keywords and phrases: q -Hausdorff matrices, Lower bound problem, q -Cesaro matrices, spectrum, fine spectrum.

An ordinary Hausdorff matrix H has the representation $H = \delta\mu\delta$, where μ is the diagonal matrix with diagonal entries

$$\delta_{nk} = (-1)^k \begin{bmatrix} n \\ k \end{bmatrix}. \quad (1.4)$$

The matrix δ is its own inverse.

A q -Hausdorff matrix H_q can be written in the form

$$H_q = \delta_q \mu (\delta_q)^{-1} \quad (1.5)$$

where μ is again the diagonal matrix with entries μ_k , and

$$(\delta_q)_{nk} = (-1)^k \begin{bmatrix} n \\ k \end{bmatrix}_q \quad (1.6)$$

For H_q matrices, δ_q is not its own inverse. However, as is the case for ordinary Hausdorff matrices, the row sum of every q -Hausdorff matrix is μ_0 .

Cesàro matrix, C_1 is defined by

$$c_{nk} = \begin{cases} \frac{1}{n+1}, & 0 \leq k \leq n \\ 0, & k > n \end{cases}. \quad (1.7)$$

For $0 < q < 1$, $C_1(q)$, the q -Hausdorff analog of C_1 which is the lower triangular matrix, has nonzero entries

$$C_1(q) = \frac{q^{n-k}(1-q)}{1-q^{n+1}}, \quad 0 \leq k \leq n. \quad (1.8)$$

For $q > 1$, entries of C_q which is the lower triangular matrix, have nonzero entries

$$C_1(q) = \frac{q^k(q-1)}{q^{n+1}-1}, \quad 0 \leq k \leq n. \quad (1.9)$$

Also, if we take the limit for $q \rightarrow 1^-$ in (1.6), then the Cesàro matrix C_1 is obtained.

A remarkable production of research involving q -series and q -differences in [28] has seen in the last thirty years.

Akgün and Rhoades show that the $C_1(q)$ matrices, the q -Hausdorff analogs of the Cesàro matrix of order 1, are all equivalent to convergence, and that the same is true of the q -Hölder matrices in [5].

The Fourier theory containing specific q -analogs of trigonometric functions was first discussed in [12] and [14]. A complete development of q -Fourier theory should include a suitable summability theory. An introduction was made to q -Fourier theory using the q -analog of the Cesàro summability in [13].

Recently, there are many studies on q -analogs of matrix methods such as Cesàro, Hölder, Euler and Hausdorff methods [5], [13], [42], [10]. There are also new studies such as q -density and q -statistical convergence [3], [16].

1.1. The spectrum

Let X and Y be the Banach spaces, and $L : X \rightarrow Y$ also be a bounded linear operator. By $R(L)$, we denote the range of L , i.e.,

$$R(L) = \{y \in Y : y = Lx, x \in X\}.$$

By $B(X)$, we also denote the set of all bounded linear operators on X into itself. If X is any Banach space and $L \in B(X)$ then the adjoint L^* of L is a bounded linear operator on the dual X^* of X defined by $(L^*f)(x) = f(Lx)$ for all $f \in X^*$ and $x \in X$.

Let $L : D(L) \rightarrow X$ be a linear operator, defined on $D(L) \subset X$, where $D(L)$ denote the domain of L and X is a complex normed linear space. For $L \in B(X)$ we associate a complex number α with the operator $(\alpha I - L)$ denoted by L_α defined on the same domain $D(L)$, where I is the identity operator. The inverse $(\alpha I - L)^{-1}$, denoted by L_α^{-1} is known as the resolvent operator of L .

A regular value of L is a complex number α of L such that L_α^{-1} exists, is bounded and, is defined on a set which is dense in X .

The resolvent set of L is the set of all such regular values of L , denoted by $\rho(L, X)$. Its complement is given by $\mathbb{C} \setminus \rho(L, X)$ in the complex plane \mathbb{C} is called the spectrum of L , denoted by $\sigma(L, X)$. Thus the spectrum $\sigma(L, X)$ consist of those values of $\alpha \in \mathbb{C}$, for which L_α is not invertible.

The spectrum $\sigma(L, X)$ is union of three disjoint sets as follows: The point (discrete) spectrum $\sigma_p(L, X)$ is the set such that L_α^{-1} does not exist. Further $\alpha \in \sigma_p(L, X)$ is called the eigenvalue of L . We say that $\alpha \in \mathbb{C}$ belongs to the continuous spectrum $\sigma_c(L, X)$ of L if the resolvent operator L_α^{-1} is defined on a dense subspace of X and is unbounded. Furthermore, we say that $\alpha \in \mathbb{C}$ belongs to the residual spectrum $\sigma_r(L, X)$ of L if the resolvent operator L_α^{-1} exists, but its domain of definition (i.e. the range $R(\alpha I - L)$ of $\alpha I - L$) is not dense in X ; in this case L_α^{-1} may be bounded or unbounded. Together with the point spectrum, these two subspectra form a disjoint subdivision

$$\sigma(L, X) = \sigma_p(L, X) \cup \sigma_c(L, X) \cup \sigma_r(L, X) \tag{1.10}$$

of the spectrum of L .

1.2. Subdivision of the spectrum

The spectrum $\sigma(L, X)$ is partitioned into three sets which are not necessarily disjoint as follows:

If there exists a sequence (x_n) in X such that $\|x_n\| = 1$ and $\|Lx_n\| \rightarrow 0$ as $n \rightarrow \infty$ then (x_n) is called Weyl sequence for L .

We call the set

$$\sigma_{ap}(L, X) := \{\alpha \in \mathbb{C} : \text{there exists a Weyl sequence for } \alpha I - L\} \tag{1.11}$$

as the approximate point spectrum of L . Moreover, the set

$$\sigma_\delta(L, X) := \{\alpha \in \sigma(L, X) : \alpha I - L \text{ is not surjective}\} \tag{1.12}$$

is called defect spectrum of L . Finally, the set

$$\sigma_{co}(L, X) = \{\alpha \in \mathbb{C} : \overline{R(\alpha I - L)} \neq X\} \tag{1.13}$$

is called compression spectrum in the literature.

The following Proposition is quietly useful for calculating the separation of the spectrum of linear operator in Banach spaces.

PROPOSITION 1.1. ([4], Proposition 1.3) *The spectra and subspectra of an operator $L \in B(X)$ and its adjoint $L^* \in B(X^*)$ are related by the following relations:*

- (a) $\sigma(L^*, X^*) = \sigma(L, X)$, (b) $\sigma_c(L^*, X^*) \subseteq \sigma_{ap}(L, X)$,
- (c) $\sigma_{ap}(L^*, X^*) = \sigma_\delta(L, X)$, (d) $\sigma_\delta(L^*, X^*) = \sigma_{ap}(L, X)$,
- (e) $\sigma_p(L^*, X^*) = \sigma_{co}(L, X)$, (f) $\sigma_{co}(L^*, X^*) \supseteq \sigma_p(L, X)$,
- (g) $\sigma(L, X) = \sigma_{ap}(L, X) \cup \sigma_p(L^*, X^*) = \sigma_p(L, X) \cup \sigma_{ap}(L^*, X^*)$.

1.3. Goldberg’s classification of spectrum

If $T \in B(X)$, then there are three cases for $R(T)$:

(I) $R(T) = X$, (II) $\overline{R(T)} = X$, but $R(T) \neq X$, (III) $\overline{R(T)} \neq X$ and three cases for T^{-1} :

(1) T^{-1} exists and continuous, (2) T^{-1} exists but discontinuous, (3) T^{-1} does not exist.

If these cases are combined in all possible ways, nine different states are created. These are labelled by: $\mathcal{I}_1, \mathcal{I}_2, \mathcal{I}_3, \mathcal{II}_1, \mathcal{II}_2, \mathcal{II}_3, \mathcal{III}_1, \mathcal{III}_2, \mathcal{III}_3$ (see [29]).

$\sigma(L, X)$ can be divided into subdivisions $\mathcal{I}_2\sigma(L, X) = \emptyset, \mathcal{I}_3\sigma(L, X), \mathcal{II}_2\sigma(L, X), \mathcal{II}_3\sigma(L, X), \mathcal{III}_1\sigma(L, X), \mathcal{III}_2\sigma(L, X), \mathcal{III}_3\sigma(L, X)$. For example, if $T = \alpha I - L$ is in a given state, \mathcal{III}_2 (say), then we write $\alpha \in \mathcal{III}_2\sigma(L, X)$.

By the definitions given above, we can write following table

		1	2	3
		L_α^{-1} exists and is bounded	L_α^{-1} exists and is unbounded	L_α^{-1} does not exist
\mathcal{I}	$R(\alpha I - L) = X$	$\alpha \in \rho(L, X)$	–	$\alpha \in \sigma_p(L, X)$ $\alpha \in \sigma_{ap}(L, X)$
\mathcal{II}	$\overline{R(\alpha I - L)} = X$	$\alpha \in \rho(L, X)$	$\alpha \in \sigma_c(L, X)$ $\alpha \in \sigma_{ap}(L, X)$ $\alpha \in \sigma_\delta(L, X)$	$\alpha \in \sigma_p(L, X)$ $\alpha \in \sigma_{ap}(L, X)$ $\alpha \in \sigma_\delta(L, X)$
\mathcal{III}	$\overline{R(\alpha I - L)} \neq X$	$\alpha \in \sigma_r(L, X)$ $\alpha \in \sigma_\delta(L, X)$ $\alpha \in \sigma_{co}(L, X)$	$\alpha \in \sigma_r(L, X)$ $\alpha \in \sigma_{ap}(L, X)$ $\alpha \in \sigma_\delta(L, X)$ $\alpha \in \sigma_{co}(L, X)$	$\alpha \in \sigma_p(L, X)$ $\alpha \in \sigma_{ap}(L, X)$ $\alpha \in \sigma_\delta(L, X)$ $\alpha \in \sigma_{co}(L, X)$

Table 1: Subdivision of the spectrum

The spectrum of matrix transformations on sequence spaces has also been found by various authors in [11], [15], [37].

The discrete spectral decomposition of the spectrum of bounded linear operators (in the sense of Goldberg classification or residual, point and continuous spectra) has been made by many authors to this day in [1], [24], [26], [27], [30], [31], [32], [34], [35], [38], [39], [40], [41], [43], [46], [49]. Durna and Yildirim gave the table above in [19] and [20]. After this date, many authors have used this table to perform spectral decomposition of the spectrum of bounded linear operators (in terms of approximate point spectrum, defect spectrum and compression spectrum) in [2], [6], [7], [8], [9], [17], [18], [21], [22], [23], [25], [33], [44], [45], [47] and [48].

2. Generalized

In this section, boundedness, spectra, fine spectra and various spectral separations of spectrum of $C_1(q) = (a_{nk}(q))$ on the sequence space c have been determined.

The following theorem gives us necessary and sufficient condition to have a bounded linear transformation of an infinite matrix $A = (a_{nk})$ on c .

THEOREM 2.1. (Kojima-Schur (Maddox [36, p. 166])) $A = (a_{nk}) \in B(c)$ if and only if i) $\|A\| = \sup_n \sum_k |a_{nk}| < \infty$ and ii) for each $p \in \mathbb{N}$ there exists $\lim_{n \rightarrow \infty} \sum_{k=p}^{\infty} a_{nk} = a_p$.

Let us now show that the q -analog of the Cesàro operator has a bounded linear transformation on c :

LEMMA 2.2. $C_1(q) \in B(c)$ and $\|C_1(q)\| = 1$.

Proof. For each k

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{k=p}^{\infty} c_{nk} &= \lim_{n \rightarrow \infty} \sum_{k=p}^n \frac{q^{n-k}}{1+q+\dots+q^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1+q+\dots+q^n} \sum_{k=p}^n q^{n-k} \\ &= \lim_{n \rightarrow \infty} \frac{1-q^{n-p+1}}{1-q} \\ &= \begin{cases} \left(\frac{1}{q}\right)^p, & q > 1 \\ 1, & 0 < q < 1 \end{cases} \end{aligned}$$

and

$$\|C_1(q)\| = \sup_n \sum_{k=0}^{\infty} |c_{nk}| = \sup_n \left\{ \frac{1}{1+q+\dots+q^n} \sum_{k=0}^n q^{n-k} \right\} = 1.$$

From the above statements and Theorem 2.1, the desired result is obtained. \square

It can be easily seen that there are two different matrices for states $0 < q < 1$ and $q > 1$.

Throughout this article we will cover the matrix obtained with case $0 < q < 1$, the matrix obtained with case $q > 1$ can be further studied in another study.

We need the following Lemma:

LEMMA 2.3. [50, p. 267] *If $T : c \rightarrow c$ is a linear transformation and $T^* : \ell_1 \rightarrow \ell_1$, $T^*g = g \circ T$, $g \in c^* \cong \ell_1$, then T and T^* have matrix representations, also $T^* : \ell_1 \rightarrow \ell_1$ is given by*

$$T^* = A^* = \begin{pmatrix} \chi(\lim A) (\vartheta_n)_{n=0}^\infty \\ (a_k)_{k=0}^\infty & A^t \end{pmatrix} = \begin{pmatrix} \chi(\lim A) & \vartheta_0 & \vartheta_1 & \vartheta_2 & \cdots \\ a_0 & a_{00} & a_{10} & a_{20} & \cdots \\ a_1 & a_{01} & a_{11} & a_{21} & \cdots \\ a_2 & a_{02} & a_{12} & a_{22} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

where

$$\begin{aligned} a_k &= \lim_n a_{nk} \\ \chi(A) &= \lim A e - \sum_{k=0}^\infty \lim A e_k = \lim_n \sum_k a_{nk} - \sum_k \lim_n a_{nk} \\ \vartheta_n &= \chi(P_n \circ T) = (P_n \circ T) e - \sum_k a_{nk}, \\ a_{nk} &= P_n(T(e_k)) = (T(e_k))_n. \end{aligned}$$

If $0 < q < 1$, then one can get the following result from the above lemma for q -Cesàro matrix.

Making use of the relations given above and Lemma 2.3, we easily arrive at the following Lemma:

LEMMA 2.4. *The adjoint of $C_1(q)$ on c is given by*

$$C_1^*(q) = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & & & \\ 0 & C_1^t(q) & & \\ \vdots & & & \end{pmatrix}. \tag{2.1}$$

Proof. Since $0 < q < 1$, we have

$$c_k = \lim_n c_{nk} = \lim_{n \rightarrow \infty} \frac{q^{n-k}(1-q)}{1-q^{n+1}} = q^{-k}(1-q) \lim_{n \rightarrow \infty} \frac{q^n}{1-q^{n+1}} = 0$$

and

$$\sum_{k=0}^n c_{nk} = \frac{(1-q)}{1-q^{n+1}} \sum_{k=0}^n q^{n-k} = \frac{(1-q)}{1-q^{n+1}} \sum_{k=0}^n q^k = \frac{(1-q)}{1-q^{n+1}} \frac{1-q^{n+1}}{(1-q)} = 1.$$

Hence

$$\chi(C_1(q)) = \lim_n \sum_k c_{nk} - \sum_k \lim_n c_{nk} = 1.$$

Also, since

$$(P_n \circ C_1(q))e = \left\{ \sum_{k=0}^n c_{nk} x_k \right\}_{x=e} = \sum_{k=0}^n c_{nk} = 1,$$

we have

$$\vartheta_n = (P_n \circ C_1(q))e - \sum_{k=0}^n c_{nk} = 1 - 1 = 0$$

Hence, by Lemma 2.3, we have

$$C_1^*(q) = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & & & \\ 0 & C_1^t(q) & & \\ \vdots & & & \end{pmatrix}.$$

This proves Lemma. \square

In the following theorems, let's calculate $\sigma_p(C_1(q), c)$, $\sigma_p(C_1^*(q), \ell_1)$, $\sigma(C_1(q), c)$ and $\sigma_r(C_1(q), c)$, respectively.

THEOREM 2.5. $\sigma_p(C_1(q), c) = \{1\}$.

Proof. Let $\sigma_p(C_1(q), c) \neq \emptyset$. Then, in this case there is $0 \neq x \in c$, such that $C_1(q)x = \alpha x$. Hence

$$\begin{aligned} x_0 &= \alpha x_0 \\ \frac{1}{1+q}(qx_0 + x_1) &= \alpha x_1 \\ \frac{1}{1+q+q^2}(q^2x_0 + qx_1 + x_2) &= \alpha x_2 \\ &\vdots \\ \frac{1}{1+q+q^2+\cdots+q^n}(q^n x_0 + q^{n-1}x_1 + \cdots + qx_{n-1} + x_n) &= \alpha x_n \\ &\vdots \end{aligned}$$

the above equalities are obtained. If x_m , $m \geq 1$ is the first component that is different from zero of sequence $(x_n) = x$, then

$$\alpha = \frac{1}{1+q+\cdots+q^m}$$

is obtained. Furthermore, the statement

$$x_{m+1} = \frac{1+q+\cdots+q^m}{q^m} x_m$$

is provided. Similarly we obtain

$$\begin{aligned} x_{m+2} &= \frac{(1+q+\dots+q^m)(1+q+\dots+q^{m+1})}{q^{2m}(1+q)}x_m, \\ x_{m+3} &= \frac{(1+q+\dots+q^m)(1+q+\dots+q^{m+1})(1+q+\dots+q^{m+2})}{q^{3m}(1+q)(1+q+q^2)}x_m, \\ x_{m+4} &= \frac{(1+q+\dots+q^m)(1+q+\dots+q^{m+1})(1+q+\dots+q^{m+2})(1+q+\dots+q^{m+3})}{q^{4m}(1+q)(1+q+q^2)(1+q+q^2+q^3)}x_m. \end{aligned}$$

If we continue in this way then we have

$$x_n = \frac{(1+q+\dots+q^m)(1+q+\dots+q^{m+1})\dots(1+q+\dots+q^{n-1})}{q^{(n-m)m}(1+q)(1+q+q^2)\dots(1+q+\dots+q^{n-m-1})} \tag{2.2}$$

is obtained for $n > m$ with $0 < q < 1$, so $q^m > q^n$. Thereby for $n > m$, $1 - q^m < 1 - q^n$ and so $\frac{1-q^n}{1-q^m} > 1$. In that case, from (2.2)

$$\begin{aligned} x_n &= \frac{\frac{1-q^{m+1}}{1-q} \frac{1-q^{m+2}}{1-q} \dots \frac{1-q^n}{1-q}}{q^{(n-m)m} \frac{1-q^2}{1-q} \frac{1-q^3}{1-q} \dots \frac{1-q^{n-m}}{1-q}} \\ &= \frac{(1-q^{m+1})(1-q^{m+2})\dots(1-q^n)}{(1-q^2)(1-q^3)\dots(1-q^{n-m})} \frac{1}{q^{(n-m)m}} \\ &= \frac{1-q^{m+1}}{1-q} \frac{1-q^{m+2}}{1-q^2} \dots \frac{1-q^n}{1-q^{n-m}} \frac{1-q}{q^{(n-m)m}} \\ &\geq \frac{1-q}{q^{(n-m)m}} \rightarrow \infty, n \rightarrow \infty. \end{aligned}$$

That is, $x = (x_n) \notin c$. Also, for $\alpha = 1$, $C_1(q)x = \alpha x$ with $x = (1, 1, 1, \dots) \in c$. This indicates that $\sigma_p(C_1(q), c) = \{1\}$. \square

THEOREM 2.6. $\sigma_p(C_1^*(q), \ell_1) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q} \right\} \cup \{1\}$.

Proof. Let $x \neq 0$ and $C_1^*(q)x = \alpha x$. Then, the below equalities are obtained

$$\begin{aligned} x_0 &= \alpha x_0 \\ x_1 + \frac{q}{1+q}x_2 + \frac{q^2}{1+q+q^2}x_3 + \frac{q^3}{1+q+q^2+q^3}x_4 + \dots &= \alpha x_1 \\ \frac{1}{1+q}x_2 + \frac{q}{1+q+q^2}x_3 + \frac{q^2}{1+q+q^2+q^3}x_4 + \dots &= \alpha x_2 \\ &\vdots \end{aligned}$$

Thus

$$x_n = \frac{x_1}{q^n} \prod_{k=1}^n \left(1 - \frac{1-q}{1-q^k} \frac{1}{\alpha} \right), \quad n = 2, 3, \dots$$

Suppose $\alpha \in \left\{ 1, \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$.

If $\alpha = 1$, then $x = (x_0, x_1, 0, \dots) \neq 0$ and $C_1^*(q)x = x$ are obtained so $1 \in \sigma_p(C_1^*(q), \ell_1)$.

If $\alpha = \frac{1}{1+q}$, then $x = (x_0, x_1, -x_1, 0, 0, \dots) \neq 0$ and $C_1^*(q)x = \frac{1}{1+q}x$ so $\frac{1}{1+q} \in \sigma_p(C_1^*(q), \ell_1)$.

If $\alpha = \frac{1}{1+q+q^2}$, then $x = (x_0, x_1, -(1+q)x_1, qx_1, 0, 0, \dots) \neq 0$ and $C_1^*(q)x = \frac{1}{1+q+q^2}x$ hence $\frac{1}{1+q+q^2} \in \sigma_p(C_1^*(q), \ell_1)$.

In a similar way, $\left\{ \frac{1}{1+q+q^2+q^3}, \frac{1}{1+q+q^2+q^3+q^4}, \dots \right\} \subset \sigma_p(C_1^*(q), \ell_1)$ is obtained. Is there any eigenvalues other than $\left\{ 1, \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$?

Now, let us accept $\alpha \notin \left\{ 1, \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$. By the ratio test

$$\lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = \lim_{n \rightarrow \infty} \frac{1}{q} \left| 1 - \frac{1-q}{1-q^{n+1}} \frac{1}{\alpha} \right| = \frac{1}{q} \left| 1 - \frac{1-q}{\alpha} \right|$$

is obtained. Move from here


$$\frac{1}{q} \left| 1 - \frac{1-q}{\alpha} \right| < 1 \Leftrightarrow \left| 1 - \frac{1-q}{\alpha} \right| < q$$

Take $\alpha = \lambda + i\mu$ in the above equation we get

$$\begin{aligned} &\Leftrightarrow \left| 1 - \frac{1-q}{\lambda^2 + \mu^2} \lambda + \frac{1-q}{\lambda^2 + \mu^2} \mu i \right| < q \\ &\Leftrightarrow 1 - \frac{2(1-q)\lambda}{\lambda^2 + \mu^2} + \frac{(1-q)^2}{\lambda^2 + \mu^2} < q^2 \\ &\Leftrightarrow 1 - q^2 < (1-q) \left(\frac{2\lambda}{\lambda^2 + \mu^2} - \frac{1-q}{\lambda^2 + \mu^2} \right) \\ &\Leftrightarrow 1 + q < \frac{2\lambda}{\lambda^2 + \mu^2} - \frac{1-q}{\lambda^2 + \mu^2} \\ &\Leftrightarrow \lambda^2 + \mu^2 < \frac{2\lambda}{1+q} - \frac{1-q}{1+q} \\ &\Leftrightarrow \left(\lambda - \frac{1}{1+q} \right)^2 + \mu^2 < \frac{1}{(1+q)^2} - \frac{1-q}{1+q} = \frac{q^2}{(1+q)^2} \end{aligned}$$

$$\Leftrightarrow \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q}$$

has the same meaning. So from ratio test, if $\left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q}$ then $\sigma_p(C_1^*(q), \ell_1) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q} \right\} \cup \{1\}$ with $x_n \in \ell_1$ is obtained. \square



THEOREM 2.7. $\sigma(C_1(q), c) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| \leq \frac{q}{1+q} \right\}$.

Proof. We will show the proof of Theorem by showing that complex numbers α with $\left| \alpha - \frac{1}{1+q} \right| > \frac{q}{1+q}$ are present in $\rho(C_1(q), c)$. From Theorem 2.5, $\alpha I - C_1(q)$ is one to one. Let us now show that $(\alpha I - C_1(q))^{-1} \in B(c)$ for complex numbers α with $\left| \alpha - \frac{1}{1+q} \right| > \frac{q}{1+q}$.

From the equality $(\alpha I - C_1(q))x = y$,

$$x_n = \frac{1}{\alpha - \frac{1}{1+q+\dots+q^n}} y_n + \frac{1}{1+q+\dots+q^n} \sum_{k=0}^{n-1} \frac{y_k}{q} \frac{1}{\alpha^2} \prod_{i=k}^n \frac{\alpha q}{\alpha - \frac{1}{1+q+\dots+q^i}}$$

is obtained. Now we have to find the value of

$$\begin{aligned} & \lim_{n \rightarrow \infty} x_n \\ &= \lim_{n \rightarrow \infty} \left[\frac{1}{\alpha - \frac{1}{1+q+\dots+q^n}} y_n + \frac{1}{1+q+\dots+q^n} \sum_{k=0}^{n-1} \frac{y_k}{q} \frac{1}{\alpha^2} \prod_{i=k}^n \frac{\alpha q}{\alpha - \frac{1}{1+q+\dots+q^i}} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{\alpha - \frac{1}{1+q+\dots+q^n}} y_n + \lim_{n \rightarrow \infty} \frac{1}{1+q+\dots+q^n} \sum_{k=0}^{n-1} \frac{y_k}{q} \frac{1}{\alpha^2} \prod_{i=k}^n \frac{\alpha q}{\alpha - \frac{1}{1+q+\dots+q^i}} \end{aligned}$$

$$\begin{aligned}
 &= \frac{\ell}{\alpha + q - 1} + \lim_{n \rightarrow \infty} \frac{1 - q}{1 - q^{n+1}} \sum_{k=0}^{n-1} \frac{y_k}{q} \frac{1}{\alpha^2} \prod_{i=k}^n \frac{\alpha q}{\alpha - \frac{1 - q}{1 - q^{i+1}}} \\
 &= \frac{\ell}{\alpha + q - 1} + \lim_{n \rightarrow \infty} \alpha^{n-1} \frac{1 - q}{1 - q^{n+1}} \frac{\sum_{k=0}^{n-1} \frac{y_k \prod_{i=k}^n (1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{(\alpha q)^k} \frac{1}{\left(\frac{1}{q}\right)^n}
 \end{aligned}$$

where $\ell = \lim_{n \rightarrow \infty} y_n$ for $y_n \in c$.

Let $a_n = \sum_{k=0}^n \frac{y_k \prod_{i=k}^n \frac{1 - q}{1 - q - \alpha(1 - q^{i+1})}}{q^k}$ and $b_n = \left(\frac{1}{q}\right)^n$. That is; let

$$x_n = \frac{a_n}{b_n}. \tag{2.3}$$

So, since

$$\begin{aligned}
 a_{n+1} - a_n &= \sum_{k=0}^n \frac{y_k \prod_{i=k}^{n+1} \frac{(1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{(\alpha q)^k} - \sum_{k=0}^{n-1} \frac{y_k \prod_{i=k}^n \frac{(1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{(\alpha q)^k} \\
 &= \frac{y_n \frac{(1 - q^{n+1})}{\alpha (1 - q^{n+1}) - (1 - q)} \frac{(1 - q^{n+2})}{\alpha (1 - q^{n+2}) - (1 - q)}}{(\alpha q)^n} \\
 &\quad + \sum_{k=0}^{n-1} \frac{\frac{(1 - q^{n+2})}{\alpha (1 - q^{n+2}) - (1 - q)} \prod_{i=k}^n \frac{(1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{q^k} \\
 &\quad - \sum_{k=0}^{n-1} \frac{y_k \prod_{i=k}^n \frac{(1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{(\alpha q)^k} \\
 &= \frac{y_n \frac{(1 - q^{n+1})}{\alpha (1 - q^{n+1}) - (1 - q)} \frac{(1 - q^{n+2})}{\alpha (1 - q^{n+2}) - (1 - q)}}{(\alpha q)^n} \\
 &\quad + \sum_{k=0}^{n-1} \frac{\left[\frac{(1 - q^{n+2})}{\alpha (1 - q^{n+2}) - (1 - q)} - 1 \right] \prod_{i=k}^n \frac{(1 - q^{i+1})}{\alpha (1 - q^{i+1}) - (1 - q)}}{q^k} \\
 &= \frac{y_n \frac{(1 - q^{n+1})}{\alpha (1 - q^{n+1}) - (1 - q)} \frac{(1 - q^{n+2})}{\alpha (1 - q^{n+2}) - (1 - q)}}{(\alpha q)^n} \\
 &\quad + \frac{(1 - \alpha) (1 - q^{n+2}) + (1 - q)}{\alpha (1 - q^{n+2}) - (1 - q)} a_n
 \end{aligned}$$

we get

$$\begin{aligned} & \frac{a_{n+1} - a_n}{b_{n+1} - b_n} \\ &= \frac{y_n \frac{(1 - q^{n+1})}{\alpha(1 - q^{n+1}) - (1 - q)} \frac{(1 - q^{n+2})}{\alpha(1 - q^{n+2}) - (1 - q)}}{\frac{1}{(\alpha q)^n}} + \frac{(1 - \alpha)(1 - q^{n+2}) + (1 - q)}{\alpha(1 - q^{n+2}) - (1 - q)} a_n \\ &= \frac{\frac{1}{q^{n+1}} - \frac{1}{q^n}}{\frac{1}{q^{n+1}} - \frac{1}{q^n}} \\ &= \frac{y_{n+1}}{1 - q - \alpha(1 - q^{n+2})} + \frac{q}{1 - q} \left[\frac{\alpha(1 - q^{n+2})}{1 - q - \alpha(1 - q^{n+2})} \right] \frac{a_n}{b_n}. \tag{2.4} \end{aligned}$$

From (b_n) increasing and Stolz theorem, $\lim_{n \rightarrow \infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = \lim_{n \rightarrow \infty} \frac{a_n}{b_n}$ is valid. Let's ap-
 prove $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$. Also, $\lim_{n \rightarrow \infty} y_n = \ell$ with $y \in c$. In (2.4), if we switch to the limit for
 $n \rightarrow \infty$

$$L = \frac{\ell}{1 - q - \alpha} + \frac{q}{1 - q} \left[\frac{\alpha}{1 - q - \alpha} \right] L$$

is obtained. Because of the fact that $\alpha \neq 1 - q$ and $\alpha \neq (1 - q)^2$ for complex numbers
 α with $\left| \alpha - \frac{1}{1 + q} \right| > \frac{q}{1 + q}$

$$L = \frac{(1 - q)\ell}{(1 - q)^2 - \alpha}$$

is given. In (2.4), if we switch to the limit for $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{(1 - q)\ell}{(1 - q)^2 - \alpha}$$

is valid. So, for complex numbers α with $\left| \alpha - \frac{1}{1 + q} \right| > \frac{q}{1 + q}$, $x = (x_n) \in c$ is ob-
 tained. Now let's find $\|x_n\|_c$. Since $y \in c$, $\|x_n\|_c$ is bounded. Hence, for each $k \in \mathbb{N}$,
 $|y_k| \leq M$ with $M > 0$. Also,

$$\begin{aligned} \left| \alpha - \frac{1}{1 + q} \right| > \frac{q}{1 + q} &\Leftrightarrow \frac{1}{\left| \alpha(1 + q) - 1 \right|} < \frac{1}{q} \\ &\Leftrightarrow \left| \frac{(1 - q)q}{\alpha(1 - q^{i+1}) - (1 - q)} \right| < 1 \end{aligned}$$

the above is the equivalent $\left| \frac{(1 - q)q}{\alpha(1 - q^{i+1}) - (1 - q)} \right| = p_i$ with $0 < p_i < 1$. If, we choose

$p := \max_{k=0}^n p_k$, then

$$\begin{aligned} \|x_n\|_{c_0} &= \sup_n |x_n| = \sup_n \left| \sum_{k=0}^n \frac{y_k}{q} \prod_{i=k}^n \frac{(1-q)q}{\alpha(1-q^{i+1}) - (1-q)} \right| \\ &\leq \frac{M}{q} \sup_n \sum_{k=0}^n \prod_{i=k}^n \left| \frac{(1-q)q}{\alpha(1-q^{i+1}) - (1-q)} \right| \\ &\leq \frac{M}{q} \sup_n \sum_{k=0}^n \prod_{i=k}^n p = \frac{M}{q} \sup_n \sum_{k=0}^n p^{n-k+1} \\ &\leq \frac{Mp}{(1-p)q} \sup_n (1-p^{n+1}) = \frac{Mp}{q} \end{aligned}$$

is obtained so $(\alpha I - C_1(q))^{-1} \in B(c)$. Consequently from equality $\sigma(C_1(q), c) = \mathbb{C} \setminus \rho(C_1(q), c)$, the proof is completed. \square

REMARK 2.8. As shown in Table 1, $\sigma(C_1(q_1), c) \subset \sigma(C_1(q_2), c)$ is valid for $0 < q_1 < q_2 < 1$. Also

$$\lim_{q \rightarrow 1^-} \sigma(C_1(q), c) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{2} \right| \leq \frac{1}{2} \right\}$$

is valid and this is equal to the set of $\sigma(C_1, c)$ from [37]. Moreover, let's note that $\lim_{q \rightarrow 0^+} \sigma(C_1(q), c) = \{1\}$. So the spectrum collapses for $q \rightarrow 0^+$.

THEOREM 2.9. $\sigma_r(C_1(q), c) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q} \right\} \cup \{1\}$.

Proof. Since $\sigma_r(C_1(q), c) = \sigma_p(C_1^*(q), \ell_1) \setminus \sigma_p(C_1(q), c)$, Theorems 2.5 and 2.6 give us required result. \square

Next, we appoint the two parts $III_3\sigma(C_1(q), c)$ and $III_2\sigma(C_1(q), c)$; it gives a finer subdivision of the spectrum. For this, we need the following lemmas.

LEMMA 2.10. (Golberg [29, p. 60]) *T has a bounded inverse if and only if T^* is onto.*

LEMMA 2.11. ([29], Theorem II 3.7) *A linear operator T has a dense range if and only if the adjoint operator T^* is one to one.*

THEOREM 2.12. $III_2\sigma(C_1(q), c) = \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$.

Proof. From Theorem 2.9, $\left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \subseteq \sigma_r(C_1(q), c)$ is valid and it is clearly known that $\left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \subseteq III$, where *III* is in Table 1.

Now we have to show whether this cluster belongs to (1) or (2). For this we will use the Lemma 2.10.

For every $y \in \ell_1$, is there $x \in \ell_1$ such that $(C_1(q) - \alpha I)^* x = y$?

Let $(C_1(q) - \alpha I)^* x = y$. In this case

$$(1 - \alpha)x_0 = y_0$$

$$(1 - \alpha)x_1 + \frac{q}{1+q}x_2 + \frac{q^2}{1+q+q^2}x_3 + \frac{q^3}{1+q+q^2+q^3}x_4 + \dots = y_1$$

$$\left(\frac{1}{1+q} - \alpha \right)x_2 + \frac{q}{1+q+q^2}x_3 + \frac{q^2}{1+q+q^2+q^3}x_4 + \dots = y_2$$

$$\vdots$$

is obtained. If we take x_n from this equation system,

$$x_0 = \frac{1}{1-\alpha}y_0$$

$$x_2 = \frac{1}{q\alpha} [(\alpha - 1)x_1 + y_1 - qy_2]$$

$$x_3 = \frac{1}{q\alpha} \left[\frac{1}{\alpha q} (\alpha - 1) \left(\alpha - \frac{1}{1+q} \right) x_1 + \frac{1}{\alpha q} \left(\alpha - \frac{1}{1+q} \right) y_1 + \frac{1}{\alpha(1+q)} y_1 - qy_2 \right]$$

$$x_4 = \frac{1}{q\alpha} \left[\frac{1}{(\alpha q)^2} (\alpha - 1) \left(\alpha - \frac{1}{1+q} \right) \left(\alpha - \frac{1}{1+q+q^2} \right) x_1 \right.$$

$$+ \frac{1}{(\alpha q)^2} \left(\alpha - \frac{1}{1+q} \right) \left(\alpha - \frac{1}{1+q+q^2} \right) y_1$$

$$+ \frac{1}{\alpha q} \left(\alpha - \frac{1}{1+q+q^2} \right) \frac{1}{\alpha(1+q)} y_2$$

$$\left. + \left(1 - \frac{1}{\alpha} \left(\alpha - \frac{1}{1+q+q^2} \right) \right) y_2 - qy_3 \right]$$

$$\vdots$$

are obtained. So, for $x_0 = \frac{1}{1-\alpha}y_0$ and $n = 1, 2, \dots$

$$\begin{aligned}
 x_{n+1} = & \frac{x_1}{(q\alpha)^n} \prod_{v=0}^n \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right) + \frac{y_1}{(q\alpha)^n} \prod_{v=1}^n \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right) \\
 & + \sum_{i=1+1}^n \frac{1}{\alpha \sum_{k=0}^i q^k} \frac{y_i}{(q\alpha)^{n-1}} \prod_{v=i+1}^n \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right) - \frac{1}{\alpha} y_{n+1}
 \end{aligned} \tag{2.5}$$

is obtained. If $\alpha \in \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$, then $x_{n+1} = -\frac{1}{\alpha}y_{n+1}$ with $x \in \ell_1$. This means that the operator $(C_1(q) - \alpha I)^*$ is onto. So, from Lemma 2.10, when

$$\alpha \in \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\},$$

$C_1(q) - \alpha I$ has bounded inverse and so

$$\left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \subseteq (2) \sigma(C_1(q), c)$$

is obtained, where (2) is n , Table 1.

Let us accept

$$\alpha \notin \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}.$$

In this case, considering that $y \in \ell_1$ from (2.5), sequence of (x_n) is convergent if and only if infinite product

$$\prod_{v=0}^{\infty} \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right)$$

is convergent. The limit of the general term of this product is as follows

$$\lim_{v \rightarrow \infty} \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right) = \alpha - \frac{1}{\frac{1}{1-q}} = \alpha - 1 + q$$

with $\alpha - 1 + q \neq 1$, that is, if $\alpha \neq 2 - q$ then the infinite product becomes divergent. Hence, if $\alpha \neq 2 - q$, then $x \notin c$ and so also $x \notin \ell_1$. This is, when

$$\alpha \notin \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \cup \{2 - q\},$$

it means that $(C_1(q) - \alpha I)^*$ is not onto operator. So from Lemma 2.10, when

$$\alpha \notin \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \cup \{2 - q\},$$

$C_1(q) - \alpha I$ has not bounded inverse. Thus,

$$\left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \subseteq (2) \sigma(C_1(q), c) \subseteq \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\} \cup \{2-q\}$$

is obtained. Finally, let's accept that $\alpha = 2 - q$. From (2.5), the first component of sequence (x_n) which is

$$\frac{x_1}{(q(2-q))^n} \prod_{v=0}^n \left(\alpha - \frac{1}{\sum_{k=0}^v q^k} \right),$$

infinite product, even if bounded, must be $\frac{1}{(q(2-q))^n} \rightarrow \infty$. However, we know that $0 < q < 1$, so $0 < q(2-q) < 1$ is valid. Hence, $(q(2-q))^n \rightarrow 0$ that happens if $\alpha = 2 - q$, $x \notin c$ and so also $x \notin \ell_1$. Hence, when $\alpha = 2 - q$, it means that the operator of $(C_1(q) - \alpha I)^*$ is not onto. Thus, from Lemma 2.10, when $\alpha = 2 - q$, $C_1(q) - \alpha I$ has not bounded inverse. Thus

$$III_2 \sigma(C_1(q), c) = \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$$

is obtained. \square

COROLLARY 2.13.

$$III_1 \sigma(C_1(q), c) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q} \right\} \setminus \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}.$$

Proof. Since $\sigma_r(C_1(q), c) = III_1 \sigma(C_1(q), c) \cup III_2 \sigma(C_1(q), c)$, Theorems 2.9 and 2.12 give us required result. \square

THEOREM 2.14. $III_3 \sigma(C_1(q), c) = \{1\}$.

Proof. Since $\{1\} = \sigma_p(C_1(q), c) = I_3 \sigma(C_1(q), c) \cup II_3 \sigma(C_1(q), c) \cup III_3 \sigma(C_1(q), c)$ from Table 1 and Theorem 2.5, $III_3 \sigma(C_1(q), c) \subseteq \{1\}$. Also, from (2.5), we know that

$$\ker(C_1(q) - I)^* = \{(x_0, x_1, 0, 0, \dots) : x_0, x_1 \in \mathbb{R}\} \neq \{(0, 0, \dots)\},$$

so this means that the operator of $(C_1(q) - I)^*$ is not one to one. From Lemma 2.11, the operator of $C_1(q) - \alpha I$ does not have intense image. Consequently, $III_3 \sigma(C_1(q), c) = \{1\}$ is obtained. \square

COROLLARY 2.15. $I_3 \sigma(C_1(q), c) = II_3 \sigma(C_1(q), c) = \emptyset$.

Proof. Since

$$\sigma_p(C_1(q), c) = \mathcal{I}_3\sigma(C_1(q), c) \cup \mathcal{II}\mathcal{I}_3\sigma(C_1(q), c) \cup \mathcal{III}\mathcal{I}_3\sigma(C_1(q), c)$$

from Table 1, the required result is obtained from Theorems 2.5 and 2.14. \square

In the theorems below, σ_c , σ_{ap} , σ_δ , σ_{co} spectrum types of $C_1(q)$ spectral operator on c will be examined in order.

THEOREM 2.16. $\sigma_c(C_1(q), c) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| = \frac{q}{1+q} \right\} \setminus \{1\}$.

Proof. Since $\sigma_c(C_1(q), c) = \sigma(C_1(q), c) \setminus [\sigma_r(C_1(q), c) \cup \sigma_p(C_1(q), c)]$ from Table 1, the required result is obtained from Theorems 2.5, 2.7 ve 2.9. \square

THEOREM 2.17. *The following statements are hold*

$$\begin{aligned} (a) \sigma_{ap}(C_1(q), c) &= \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| = \frac{q}{1+q} \right\} \cup \left\{ \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}, \\ (b) \sigma_\delta(C_1(q), c) &= \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| \leq \frac{q}{1+q} \right\}, \\ (c) \sigma_{co}(C_1(q), c) &= \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| < \frac{q}{1+q} \right\} \cup \{1\}. \end{aligned}$$

Proof. (a) Since

$$\sigma_{ap}(C_1(q), c) = \sigma(C_1(q), c) \setminus \mathcal{III}\mathcal{I}_1\sigma(C_1(q), c)$$

from Table 1, give us required result Theorem 2.7 and Corollary 2.13.

(b) Since

$$\sigma_\delta(C_1(q), c) = \sigma(C_1(q), c) \setminus \mathcal{I}_1\sigma(C_1(q), c)$$

from Table 1, give us required result Theorem 2.7 and Corollary 2.15.

(c) Since

$$\sigma_{co}(C_1(q), c) = \sigma_r(C_1(q), c) \cup \mathcal{III}\mathcal{I}_3\sigma(C_1(q), c)$$

from Table 1, give us required result Theorem 2.9 and Corollary 2.15. \square

COROLLARY 2.18. (a) $\sigma_{ap}(C_1^*(q), \ell_1) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| \leq \frac{q}{1+q} \right\}$,

(b) $\sigma_\delta(C_1^*(q), \ell_1) = \left\{ \alpha \in \mathbb{C} : \left| \alpha - \frac{1}{1+q} \right| = \frac{q}{1+q} \right\} \cup \left\{ 1, \frac{1}{1+q}, \frac{1}{1+q+q^2}, \dots \right\}$.

Proof. Using Proposition 1.1 (c) and (d), we have

$$\sigma_{ap}(C_1^*(q), \ell_1) = \sigma_\delta(C_1(q), c)$$

and

$$\sigma_\delta(C_1^*(q), \ell_1) = \sigma_{ap}(C_1(q), c).$$

Using Theorem 2.17 (a) and (b) we get the required results. \square

REFERENCES

- [1] A. M. AKHMEDOV AND S. R. EL-SHABRAWY, *Spectra and fine spectra of lower triangular double-band matrices as operators on ℓ_p ($1 \leq p < \infty$)*, Math. Slovaca **65**, 5 (65), 1137–1152.
- [2] R. KH. AMIROV, N. DURNA AND M. YILDIRIM, *Subdivisions of the spectra for cesaro, rhalý and weighted mean operators on c_0 , c and ℓ_p* , Iran. J. Sci. Technol. Trans. A Sci. **A3**, (2011), 175–183.
- [3] H. AKTUĞLU AND Ş. BEKAR, *q -Cesáro matrix and q -statistical convergence*, J. Comput. Appl. Math. **235**, 16 (2011), 4717–4723.
- [4] J. APPELL, E. DE PASCALE AND A. VIGNOLI, *Nonlinear Spectral Theory*, Walter de Gruyter, Berlin, New York, 2004.
- [5] A. F. AYDIN AND B. E. RHOADES, *Properties of some q -Hausdorff matrices*, Appl. Math. Comput. **219**, 14 (2013), 7392–7397.
- [6] F. BAŞAR, N. DURNA AND M. YILDIRIM, *Subdivisions of the spectra for generalised difference operator Δ_ν on the sequence space ℓ_1* , AIP Conf. Proc. **1309**, (2010), 254–260.
- [7] F. BAŞAR, N. DURNA AND M. YILDIRIM, *Subdivisions of the Spectra for Generalized Difference Operator over Certain Sequence Spaces*, Thai J. Math. **9**, 1 (2011), 285–295.
- [8] F. BAŞAR, N. DURNA AND M. YILDIRIM, *Subdivisions of the Spectra for the Triple Band Matrix over Certain Sequence Spaces*, Gen. Math. Notes, **1**, 1 (2011), 35–48.
- [9] F. BAŞAR, N. DURNA AND M. YILDIRIM, *Subdivision of the spectra for difference operator over certain sequence space*, Malays. J. Math. Sci. **6**, (2012), 151–165.
- [10] Ş. BEKAR, *q -Matrix Summability Methods*, Eastern Mediterranean University, Ph. D. Dissertation, 2010.
- [11] D. W. BOYD, *The spectrum of the Cesaro operator*, Acta Sci. Math. **29**, (1968), 31–34.
- [12] J. BUSTOZ AND J. L. CARDOSO, *Basic Analog of Fourier Series on a q -Linear Grid*, J. Approx. Theory **112**, (2001), 134–157.
- [13] J. BUSTOZ, L. GORDILLO AND F. LUIS, *q -Hausdorff summability*, J. Comput. Anal. Appl. **7**, 1 (2005), 135–48.
- [14] J. BUSTOZ AND S. K. SUSLOV, *Basic Analog of Fourier Series on a q -Quadratic Grid*, Methods Appl. Anal. **4**, (1998), 1–38.
- [15] J. M. CARDLIDGE, *Weighted Mean Matrices as operator on ℓ_p* , Indiana University, Ph. D. Thesis 1978.
- [16] R. ÇOLAK, B. C. TRIPATHY AND M. ET, *Lacunary strongly summable sequences and q -lacunary almost statistical convergence*, Vietnam J. Math. **34**, 2 (2006), 129–138.
- [17] R. DAS, B. TRIPATHY, *The spectrum and fine spectrum of the lower triangular matrix $B(r,s,t)$ on the sequence space cs* , Songklanakarin J. Sci. Technol. **38**, 3 (2016), 265–274.
- [18] R. DAS, *On the fine spectrum of the lower triangular matrix $B(r,s)$ over the Hahn sequence space*, Kyungpook Math. J. **57**, 3 (2017), 441–455.
- [19] N. DURNA AND M. YILDIRIM, *Subdivision of the Spectra for Factorable Matrices on c_0* , Gazi Univ. J. Sci. **24**, 1 (2011), 45–49.
- [20] N. DURNA AND M. YILDIRIM, *Subdivision of the spectra for factorable matrices on c and ℓ^p* , Math. Commun. **16**, 2 (2011), 519–530.
- [21] N. DURNA, M. YILDIRIM AND Ç. ÜNAL, *On The Fine Spectrum of Generalized Lower Triangular Double Band Matrices Δ_{iv} Over The Sequence Space c_0* , Cumhuriyet Sci. J. **37**, 3 (2016), 281–291.
- [22] N. DURNA, M. YILDIRIM AND R. KILIÇ, *Partition of the spectra for the generalized difference operator $B(r,s)$ on the sequence space cs* , Cumhuriyet Sci. J. **379**, 1 (2018), 7–15.
- [23] N. DURNA, *Subdivision of spectra for some lower triangular double-band matrices as operators on c_0* , Ukrain. Mat. Zh. **70**, 7 (2018), 913–922.
- [24] S. R. EL-SHABRAWY, *On the fine spectrum of the generalized difference operator $\Delta_{a,b}$ over the sequence space ℓ_p , ($1 < p < \infty$)*, Appl. Math. Inf. Sci. **6**, 1S (2012), 111S–118S.
- [25] S. R. EL-SHABRAWY AND S. H. ABU-JANAH, *Spectra of the generalized difference operator on the sequence spaces bv_0 and h* , Linear Multilinear Algebra **8**, (2018), 1691–1708.
- [26] S. J. FATHI AND L. RAHMATOLLAH, *On the fine spectra of the generalized difference operator Δ_{iv} over the sequence space ℓ_p* , JMMRC **1**, 1 (2012), 1–12.
- [27] H. FURKAN, H. BILGIÇ AND F. BAŞAR, *On the fine spectrum of the operator $B(r,s,t)$ over the sequence spaces ℓ_p and bv_p , ($1 < p < \infty$)*, Comput. Math. Appl. **60**, 7 (2010), 2141–2152.

- [28] G. GASPER AND M. RAHMAN, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, 1990.
- [29] S. GOLDBERG, *Unbounded Linear Operators*, McGraw Hill, New York, 1966.
- [30] M. GONZALEZ, *The fine spectrum of Cesàro operator in ℓ_p ($1 < p < \infty$)*, Arch. Math. **44**, (1985), 355–358.
- [31] V. KARAKAYA, M. MANAFOV AND N. ŞİMŞEK, *On the fine spectrum of the second order difference operator over the sequence spaces ℓ_p and bv_p , ($1 < p < \infty$)*, Math. Comput. Modelling **55**, 3–4 (2012), 426–431.
- [32] V. KARAKAYA AND E. ERDOĞAN, *Notes on the spectral properties of the weighted mean difference operator $G(u, v; \Delta)$ over the sequence space ℓ_1^** , Acta Math. Sci. **36**,2 (2016), 477–486.
- [33] V. KARAKAYA, M. DZH. MANAFOV AND N. ŞİMŞEK, *On fine spectra and subspectrum (approximate point, defect and compression) of operator with periodic coefficients*, J. Nonlinear Convex Anal. **18**, 4 (2017), 709–717.
- [34] R. LASHKARIPOUR AND J. FATHI, *On the Fine Spectra of the Generalized Forward Difference Operator Δ^v Over the Sequence Space ℓ_1* , Thai J. Math. **9**, 3 (2012), 605–617.
- [35] R. LASHKARIPOUR AND J. FATHI, *On the Fine Spectra of the Zweier Matrix as an Operator Over the Weighted Sequence Space $\ell_p(w)$* , Int. J. Nonlinear Anal. Appl. **3**, 1 (2012), 31–39.
- [36] I. J. MADDIX, *Elements of Functional Analysis*, Cambridge University Press, 1970.
- [37] J. I. OKUTOYI, *On the spectrum of Cesàro operator*, Ph. D. Thesis, Birmingham University, 1986.
- [38] B. L. PANIGRAHI AND P. D. SRIVASTAVA, *Spectrum and fine spectrum of generalized second order forward difference operator Δ_{mv}^2 on sequence space ℓ^1* , Demonstratio Math. **45**, 3 (2012), 593–609.
- [39] A. PAUL AND B. C. TRIPATHY, *The spectrum of the operator $D(r, 0, 0, s)$ over the sequence spaces ℓ_p and bv_p* , Hacet. J. Math. Stat. **43**, 3 (2014), 425–434.
- [40] B. E. RHOADES, *The fine spectra for weighted mean operators in $B(\ell^p)$* , Integral Equations Operator Theory **12**, 1 (1989), 82–98.
- [41] B. E. RHOADES AND M. YILDIRIM, *Spectra and fine spectra for factorable matrices*, Integral Equations Operator Theory **53**, 1 (2005), 127–144.
- [42] T. SELMANOGULLARI, E. SAVAŞ AND B. E. RHOADES, *On q -Hausdorff matrices*, Taiwanese J. Math. **15**, 6 (2011), 2429–2437.
- [43] B. C. TRIPATHY AND R. DAS, *On Fine spectrum of the upper triangular matrix $U(r, 0, 0, s)$ over the sequence spaces c_0 and c* , Proyecciones **37**, 1 (2018), 85–101.
- [44] B. C. TRIPATHY AND R. DAS, *Spectrum and Fine Spectrum of the Lower Triangular Matrix $B(r, 0, s)$ over the Sequence Space cs* , Appl. Math. Inf. Sci. **9**, 4 (2015), 2139–2145.
- [45] M. YESILKAYAGIL AND M. KIRISCI, *On the fine spectrum of the forward difference operator on the Hahn space*, Gen. Math. Notes. **33**, 2 (2016), 1–16.
- [46] M. YILDIRIM, *The spectrum and fine spectrum of the compact Rhaly operator*, Indian J. pure appl. math. **27**, 8 (1996), 779–784.
- [47] M. YILDIRIM AND N. DURNA, *The spectrum and some subdivisions of the spectrum of discrete generalized Cesàro operators on ℓ_p , ($1 < p < \infty$)*, J. Inequal. Appl. **193**, (2017), 1–13.
- [48] M. YILDIRIM, M. MURSALEEN AND Ç. DOĞAN, *The Spectrum and fine spectrum of generalized Rhaly-Cesàro matrices on c_0 and c* , Oper. Matrices **12**, 4 (2018), 955–975.
- [49] R. B. WENGER, *The fine spectra of the Hölder summability operator*, Indian J. Pure Appl. Math. **6**, 6 (1975), 695–712.
- [50] A. WILANSKY, *Summability Through Functional Analysis*, North Holland, 1984.

(Received March 21, 2019)

Nuh Durna

Department of Mathematics, Faculty of Science
University of Sivas Cumhuriyet
58140, Sivas, Turkey
e-mail: ndurna@cumhuriyet.ed.tr

Merve Esra Türkay

Department of Mathematics, Faculty of Science
University of Sivas Cumhuriyet
58140, Sivas, Turkey
e-mail: mesra@cumhuriyet.ed.tr