

## SPHERICAL SYMMETRY OF SOME UNITARY INVARIANTS FOR COMMUTING TUPLES

SAMEER CHAVAN AND KAIS FEKI

**Abstract.** We discuss spherical and Euclidean analogues of joint spectral radius, joint operator norm and joint numerical radius associated with commuting  $d$ -tuples of Hilbert space operators. In particular, we deduce their invariance under the action of the group  $\mathcal{U}(d)$  of  $d \times d$  unitary matrices. Unlike spectral and numerical radii, the analogues of joint operator norm differ in dimension  $d > 1$ . The joint hyponormality ensures that these analogues of joint operator norm agree in all dimensions. However, the separate hyponormality fails to ensure so.

*Mathematics subject classification (2020):* 47A13, 47B20.

*Keywords and phrases:* Unitary invariants, Taylor spectrum, joint hyponormality.

### REFERENCES

- [1] W. ARVESON, *Subalgebras of  $C^*$ -algebras, III. Multivariable operator theory*, Acta Math. **181** (1998), 159–228.
- [2] A. ATHAVALE, *On joint hyponormality of operators*, Proc. Amer. Math. Soc. **103** (1988), 417–423.
- [3] H. BAKLOUTI, K. FEKI, *On joint spectral radius of commuting operators in Hilbert spaces*, Linear Algebra Appl. **557** (2018) 455–463.
- [4] H. BAKLOUTI, K. FEKI, O. A. M. SID AHMED, *Joint numerical ranges of operators in semi-Hilbertian spaces*, Linear Algebra Appl. **555** (2018) 266–284.
- [5] J. W. BUNCE, *Models for  $n$ -tuples of noncommuting operators*, J. Funct. Anal. **57** (1984), 21–30.
- [6] S. CHAVAN, V. M. SHOLAPURKAR, *Completely hyperexpansive tuples of finite order*, J. Math. Anal. Appl. **447** (2017) 1009–1026.
- [7] S. CHAVAN, D. YAKUBOVICH, *Spherical tuples of Hilbert space operators*, Indiana Univ. Math. J. **64** (2015), 577–612.
- [8] M. CHŌ, M. TAKAGUCHI, *Boundary points of joint numerical ranges*, Pacific journal of math **95**, (1981), 27–35.
- [9] M. CHŌ, W. ŽELAZKO, *On geometric spectral radius of commuting  $n$ -tuples of operators*, Hokkaido Math. J., **21** (1992), 251–258.
- [10] R. E. CURTO, *Applications of several complex variables to multiparameter spectral theory. In Surveys of some recent results in operator theory*, Vol. II, volume 192 of Pitman Res. Notes Math. Ser., pages 25–90. Longman Sci. Tech., Harlow, 1988.
- [11] R. E. CURTO, *Joint hyponormality: a bridge between hyponormality and subnormality*, Operator theory: operator algebras and applications, Part 2 (Durham, NH, 1988), 69–91, Proc. Sympos. Pure Math., 51, Part 2, Amer. Math. Soc., Providence, RI, 1990.
- [12] C. GU, *Examples of  $m$ -isometric tuples of operators on a Hilbert space*, J. Korean Math. Soc. **55** (2018), 225–251.
- [13] K. FEKI, *A note on doubly commuting tuples of hyponormal operators on Hilbert spaces*, Results Math 75, 93 (2020), <https://doi.org/10.1007/s00025-020-01220-5>.
- [14] P. R. HALMOS, *A Hilbert space problem book*, Vol. 19, Springer Science and Business Media, 2012.
- [15] V. MÜLLER, A. SOLTYSIAK, *Spectral radius formula for commuting Hilbert space operators*, Studia Math. **103** (1992), 329–333.
- [16] G. POPESCU, *Unitary invariants in multivariable operator theory*, Memoirs of the American Mathematical Society, **200** (2009), vi+91 pp.

- [17] J. L. TAYLOR, *A joint spectrum for several commuting operators*, J. Funct. Anal. **6** (1970) 172–191.