

SURGERY OF FRAMES IN HILBERT SPACES

DONGWEI LI AND JING JIANG

(Communicated by D. Han)

Abstract. Frames which are tight or full spark might be considered optimally conditioned in applications. This leads to the question of perfect preconditioning of frames. In this paper, we consider the surgery of frames such that given frames can be manipulated to tight or full spark frames by removing and adding some elements. We give a necessary and sufficient condition such that a (r, k) -surgery on a frame results in a tight frame. We also provide a necessary and sufficient condition such that a $(1, k)$ -surgery on a tight frame resulting in a tight frame with same bound. Finally, we characterize that a (r, k) -surgery on a frame resulting in a full spark frame is possible. We obtain a necessary and sufficient condition such that a (r, k) -surgery on a given frame results in a full spark frame.

1. Introduction

A frame is a sequence of vectors in \mathcal{H} such that every element in \mathcal{H} has a representation as a linear combination of the frame elements and its element are not necessarily linearly independent. We also call a frame a redundant basis. Due to redundant, frames have wide applications in coding theory [11], signal processing [6, 13], quantum information [3, 5], filter bank theory [8] and neural networks [17].

Frames with nice geometric structures, such as tight or full spark, play an important role in signal processing because they provide more robust to erasures, additive noise, distortions and they also provide stable reconstruction formula. Hence, it is necessary to manipulate a given frame to a tight frame or full spark frame. There are several methods to manipulate frames to tight frames in Hilbert spaces. Kutyniok et al. introduced and characterized scalable frames such that the vectors can be rescaled to yield a tight frame [9]. However, not every frame is scalable. For example, a basis in \mathbb{R}^2 which is not an orthogonal basis is not scalable. An another technique for modification of a given frame to a tight frame is surgery which is first provided by Han et al. [4]. It is possible to generate a new tight frame by adding or removing some elements of a given frame. Li and Sun proved that every frame can be expanded to a tight frame by adding an element [10]. Sivaram et al. expanded a frame to a tight frame from the view of length of elements [14]. They called this surgery the length surgery. Copenhaver et al. generalized the results on surgery from [14] and answered the question of when length surgery resulting in a tight frame for a finite dimensional Hilbert space is possible [2].

Mathematics subject classification (2020): 42C15, 15A03, 05B20.

Keywords and phrases: Frames, surgery, full spark, tight.

Full spark frames are increasing interest in applications because they provide maximum robustness to erasures [1, 16]. However, at least to the author, there are no convenient techniques such that a frame can be modified to a full spark frames.

Our goal of this paper is to study surgery on frames such that a frame can be modified to a tight or full spark frame by removing some vectors and replacing this set with other a set of vectors.

Throughout this paper, let \mathcal{H} be an n -dimensional Hilbert space. A sequence $\{x_i\}_{i=1}^m$ is called a frame for \mathcal{H} if there exist constants $0 < A \leq B < \infty$ such that

$$A\|x\|^2 \leq \sum_{i=1}^m |\langle x, x_i \rangle|^2 \leq B\|f\|^2, \quad \forall f \in \mathcal{H}.$$

Here A is the greatest lower frame bound and B is the least upper frame bound. When $A = B$, $\{x_i\}_{i=1}^m$ is called a tight frame with bound A . A uniform frame is a frame in which all the vectors have equal norm. A unit-norm frame is a frame such that each frame element has norm one.

Let $\{x_i\}_{i=1}^m$ be a sequence of vectors in \mathcal{H} . The linear map $\Theta: \mathcal{H} \rightarrow \ell^2(\{1, \dots, m\})$ defined by $(\Theta x)_i = \langle x, x_i \rangle$ is called the analysis operator. The adjoint Θ^* such that $\Theta: \ell^2(\{1, \dots, m\}) \rightarrow \mathcal{H}$ is called the synthesis operator. The frame operator S of a sequence of vectors $\{x_i\}_{i=1}^m$ (not necessarily a frame) is defined as $\Theta^*\Theta$. For all $f \in \mathcal{H}$,

$$Sx = \Theta^*\Theta x = \sum_{i=1}^m \langle x, x_i \rangle x_i.$$

We can verify that $S = \sum_{i=1}^m x_i \otimes x_i$, where $x \otimes y$ is the elementary tensor rank-one operator defined by $(x \otimes y)z = \langle z, y \rangle x$ for $z \in \mathcal{H}$.

If a sequence of vectors $\{x_i\}_{i=1}^m$ is a frame for \mathcal{H} , its frame operator S is a positive invertible bounded linear operator on \mathcal{H} . Let $\{\lambda_i\}_{i=1}^m$ (counted with multiplicity and arranged in non-increasing order) be the eigenvalues of S , then $B = \lambda_1 \geq \dots \geq \lambda_m = A$. The frame operator $S = AI$ if and only if $\{x_i\}_{i=1}^m$ is a tight frame with bound A .

2. Surgery for tight frames

We first give a simple definition of the (r, k) -surgery.

DEFINITION 2.1. A (r, k) -surgery on a finite sequence of vectors X in \mathcal{H} removes r vectors from X and replaces them with k vectors.

For two sequences of vectors $X = \{x_i\}_{i=1}^m$, $Y = \{y_i\}_{i=1}^k$, we say that a (r, k) -surgery on X with Y means removing k vectors from X and replacing them with Y .

For the convenience, we always assume that the last r vectors are removed from X throughout this paper.

LEMMA 2.1. *Given two sequences of vectors $X = \{x_i\}_{i=1}^m$, $Y = \{y_i\}_{i=1}^k$, a (r, k) -surgery on X with Y resulting in a tight frame is possible if*

$$\frac{1}{n} \left(\sum_{i=1}^{m-r} \|x_i\|^2 + \sum_{i=1}^k \|y_i\|^2 \right) \geq \lambda_1,$$

where λ_1 is the largest eigenvalue of the frame operator of $\{x_i\}_{i=1}^{m-r}$.

Proof. Let S_X and S_Y be the frame operators of $X = \{x_i\}_{i=1}^{m-r}$ and $Y = \{y_i\}_{i=1}^k$, respectively, and let

$$S = S_X + S_Y = \sum_{i=1}^{m-r} x_i \otimes x_i + \sum_{i=1}^k y_i \otimes y_i.$$

Next, we show that it is possible to find a sequence of vectors $Y = \{y_i\}_{i=1}^k$ and a non-negative constant A such that $S = AI$.

We first consider trace of S . Since trace is additive, we have

$$nA = \text{trace}(S) = \text{trace}(S_X) + \text{trace}(S_Y) = \sum_{i=1}^{m-r} \|x_i\|^2 + \sum_{i=1}^k \|y_i\|^2. \quad (2.1)$$

Let $S_Y = AI - S_X$. In this case, we calculate eigenvalues of S_Y as following:

$$\det(S_Y - \lambda_y I) = \det(AI - S_X - \lambda_y I) = \det((A - \lambda_y)I - S_X) = \det(\lambda_x I - S_X) = 0, \quad (2.2)$$

where $\lambda_x = A - \lambda_y$. From the right side of (2.2), $\det(\lambda_x I - S_X) = 0$, we have that the eigenvalues of S_X are $\{\lambda_x\}$.

We assume that the eigenvalues of S_X are $\{\lambda_x\}$, From (2.2), we have $\lambda_y = A - \lambda_x$. Next, we show $A > 0$. Since S is positive semi-definite, S_X has n nonnegative real eigenvalues (counted with multiplicity and arranged in non-increasing order). Hence, λ_y is always real. Therefore, we can choose a nonnegative constant A such that $\lambda_y = A - \lambda_x \geq 0$. Thus there exists a constant A such that $A \geq \lambda_1$. From (2.1), we have that

$$\frac{1}{n} \left(\sum_{i=1}^{m-r} \|x_i\|^2 + \sum_{i=1}^k \|y_i\|^2 \right) \geq \lambda_1. \quad \square$$

Note that a (r, k) -surgery on X with Y resulting in a tight frame with bound A in Lemma 2.1 if and only if $S_Y = AI - S_X$. Next we give a necessary and sufficient condition such that $S_Y = AI - S_X$ from the view of majorization. We first give the following conception.

DEFINITION 2.2. [7] Let $a = \{a_i\}_{i=1}^m$, $b = \{b_i\}_{i=1}^k$ be non-increasing summable sequences of nonnegative numbers, and let $t = \min\{m, k\}$. We say that b majorizes a , noted $b \succ a$, if

$$\sum_{i=1}^j b_i \geq \sum_{i=1}^j a_i, \quad \text{for } 1 \leq i \leq j \text{ and } \sum_{i=1}^m b_i = \sum_{i=1}^k a_i.$$

THEOREM 2.1. [12] *Let $a = \{a_i\}_{i=1}^m$ be a non-increasing sequence of positive numbers and let T be a bounded positive semi-definite operator on \mathcal{H} with eigenvalues (counted with multiplicity and arranged in non-increasing order) $\Lambda = \{\lambda_i\}_{i=1}^n$. Then the following statements are equivalent:*

1. $a \prec \Lambda$.
2. *There exists a Bessel sequence $Y = \{y_i\}_{i=1}^m \subset \mathcal{H}$ such that $\|y_i\|^2 = a_i$ for $1 \leq i \leq m$ and $S_Y = T$.*

THEOREM 2.2. *Given two sequences of vectors $X = \{x_i\}_{i=1}^m$, $Y = \{y_i\}_{i=1}^k$ (arranged in non-increasing order under norm), a (r, k) -surgery on X with Y resulting in a tight frame with bound A if and only if $a \prec \Lambda$, where $a = \{\|y_i\|^2\}_{i=1}^k$, $\Lambda = \{A - \lambda_{n-i+1}\}_{i=1}^n$ and $\{\lambda_i\}_{i=1}^n$ are the eigenvalues (arranged in non-increasing order) of $S_X = \sum_{i=1}^{m-r} x_i \otimes x_i$.*

Proof. Let S_X and S_Y be the frame operators of $X = \{x_i\}_{i=1}^{m-r}$ and $Y = \{y_i\}_{i=1}^k$, respectively. Assume that $\{x_i\}_{i=1}^{m-r} \cup \{y_i\}_{i=1}^k$ is a tight frame with bound A . Let S be the frame operator of $\{x_i\}_{i=1}^{m-r} \cup \{y_i\}_{i=1}^k$, we have $S = S_X + S_Y = AI$. Then $S - S_X = S_Y \geq 0$. We see that the eigenvalues of S_Y arranged in non-increasing order are $A - \lambda_n \geq \dots \geq A - \lambda_1 \geq 0$. By Theorem 2.1 we have

$$(A - \lambda_n \geq \dots \geq A - \lambda_1) \succ (\|y_1\|^2 \cdots \|y_k\|^2),$$

thus $a \prec \Lambda$.

Conversely, let $a \prec \Lambda$, by Definition 2.2, we have

$$A = \frac{1}{n} \left(\sum_{i=1}^{m-r} \|x_i\|^2 + \sum_{i=1}^k \|y_i\|^2 \right) \geq \lambda_1.$$

From (2) of Theorem 2.1, we know that there exists a sequence $S_Y = \{y_i\}_{i=1}^k$ with frame operator $S_Y = S - S_X$ and we are done. \square

Note that if $a \prec \Lambda$ in Theorem 2.1, by Definition 2.2, we have

$$A = \frac{1}{n} \left(\sum_{i=1}^{m-r} \|x_i\|^2 + \sum_{i=1}^k \|y_i\|^2 \right) \geq \lambda_1,$$

and

$$A \geq \frac{1}{t} \sum_{i=1}^t (\|y_i\|^2 + \lambda_{n-i+1}), \quad 1 \leq t \leq \min\{n, k\}.$$

Then we can get a result for a (r, k) -surgery on a sequence of unit norm vectors.

COROLLARY 2.3. *Given two sequences of unit norm vectors $X = \{x_i\}_{i=1}^m$, $Y = \{y_i\}_{i=1}^k$. A (r, k) -surgery on X with Y resulting in a tight frame with bound A if and only if*

$$A \geq \max\left\{\lambda_1, 1 + \frac{1}{t} \sum_{i=1}^t \lambda_{n-i+1}\right\},$$

where $1 \leq t \leq \min\{n, k\}$ and $\{\lambda_i\}_{i=1}^n$ are the eigenvalues (arranged in non-increasing order) of $S_X = \sum_{i=1}^{m-r} x_i \otimes x_i$.

Proof. The proof is straightforward. \square

The author in [15, Lemma 2.3] considered a (r, k) -surgery ($r > k$) on a unit norm tight frame resulting in a unit norm tight frame. Next, we provide a necessary and sufficient condition such that a $(1, k)$ -surgery on a tight frame resulting in a tight frame with same bound.

THEOREM 2.4. *Let $X = \{x_i\}_{i=1}^m$ be a tight frame with bound A for \mathcal{H} and let $Y = \{y_i\}_{i=1}^k \subset \mathcal{H}$. A $(1, k)$ -surgery on X with Y resulting in a tight frame with bound A if and only if*

$$\|x_m\|^2 = \sum_{i=1}^k \|y_i\|^2 \quad \text{and} \quad \text{span}\{x_m\} = \text{span}\{y_i\}_{i=1}^k.$$

Proof. If X and $\{x_i\}_{i=1}^{m-1} \cup \{y_i\}_{i=1}^k$ are A -tight frames for \mathcal{H} , we have

$$AI = \sum_{i=1}^m x_i \otimes x_i, \tag{2.3}$$

and

$$AI = \sum_{i=1}^{m-1} x_i \otimes x_i + \sum_{i=1}^k y_i \otimes y_i. \tag{2.4}$$

By (2.3)–(2.4) and changing sides, we have

$$x_m \otimes x_m = \sum_{i=1}^k y_i \otimes y_i. \tag{2.5}$$

By taking trace of both sides of (2.5), we have

$$\|x_m\|^2 = \sum_{i=1}^k \|y_i\|^2.$$

We now prove $\text{span}\{x_m\} = \text{span}\{y_i\}_{i=1}^k$. Let

$$T = x_m \otimes x_m = \sum_{i=1}^k y_i \otimes y_i.$$

We have $\text{rank}(T) = 1 \leq n$. As T is positive, there exists a basis for \mathcal{H} with respect to which T is an $n \times n$ diagonal matrix with entries $\lambda_1 > \lambda_2 = \dots = \lambda_n = 0$ along its diagonal. The a -th entry along the diagonal of T is given by

$$\lambda_a = x_{i_a}^2 = \sum_{i=1}^k y_{i_a}^2,$$

where x_{ia} and y_{ia} are the a -th entries of x_i and y_i , respectively. As $\lambda_a = 0$ for $2 \leq a \leq n$, $x_{ia} = y_{ia} = 0$ for $2 \leq a \leq n$. Hence, $\dim \text{span}\{x_m\} \leq 1$ and $\dim \text{span}\{y_i\}_{i=1}^k \leq 1$. Since T is diagonal, there exists a vector $z \in \mathcal{H}$ such that $Tz \neq 0$. As $Tz \in \text{span}\{x_m\}$ and $Tz \in \text{span}\{y_i\}_{i=1}^k$,

$$\text{span}\{x_m\} = \text{span}\{Tz\} = \text{span}\{y_i\}_{i=1}^k.$$

Conversely, if we want to prove that $\{x_i\}_{i=1}^{m-1} \cup \{y_i\}_{i=1}^k$ is a tight frame with bound A , we only need to show

$$x_m \otimes x_m = \sum_{i=1}^k y_i \otimes y_i.$$

Assume that

$$\text{span}\{x_m\} = \text{span}\{y_i\}_{i=1}^k,$$

then $y_1 \cdots y_k \in \text{span}\{x_m\}$. Therefore, there exist $c_1, \dots, c_k \in \mathbb{R}$ such that for $a = 1, \dots, n$, $y_{ia} = c_i x_{ma}$ ($i = 1, \dots, k$). From $\|x_m\|^2 = \sum_{i=1}^k \|y_i\|^2$, we have

$$\|x_m\|^2 = \sum_{i=1}^k c_i^2 \|x_m\|^2,$$

and then $c_1^2 + c_2^2 + \cdots + c_k^2 = 1$. The (p, q) entry of the matrix form of $x \otimes x$ is $x_p x_q$. Then $x_m \otimes x_m = \sum_{i=1}^k y_i \otimes y_i$ is equivalent to following,

$$x_{mp} x_{mq} = y_{1p} y_{1q} + y_{2p} y_{2q} + \cdots + y_{kp} y_{kq}.$$

From $y_{ia} = c_i x_{ma}$, we have

$$\begin{aligned} y_{1p} y_{1q} + y_{2p} y_{2q} + \cdots + y_{kp} y_{kq} &= c_1^2 x_{mp} x_{mq} + c_2^2 x_{mp} x_{mq} + \cdots + c_k^2 x_{mp} x_{mq} \\ &= (c_1^2 + \cdots + c_k^2) x_{mp} x_{mq} \\ &= x_{mp} x_{mq}. \end{aligned}$$

Thus $x_m \otimes x_m = \sum_{i=1}^k y_i \otimes y_i$. Hence $\{x_i\}_{i=1}^{m-1} \cup \{y_i\}_{i=1}^k$ is a tight frame for \mathcal{H} with bound A . \square

EXAMPLE 1. Let

$$x_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad x_2 = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix}, \quad x_3 = \begin{bmatrix} -\frac{1}{2} \\ -\frac{\sqrt{3}}{2} \end{bmatrix}.$$

It is easy to verify that $\{x_1, x_2, x_3\}$ is a tight frame with bound $\frac{3}{2}$. Let

$$y_1 = \begin{bmatrix} -\frac{\sqrt{3}}{6} \\ -\frac{1}{2} \end{bmatrix}, \quad y_2 = \begin{bmatrix} -\frac{\sqrt{6}}{6} \\ -\frac{3\sqrt{2}}{6} \end{bmatrix}.$$

By removing x_3 from $\{x_1, x_2, x_3\}$ and adding y_1, y_2 , it is easy to verify that $\|x_3\|^2 = \|y_1\|^2 + \|y_2\|^2$ and $\text{span}\{x_3\} = \text{span}\{y_1, y_2\}$, and $\{x_1, x_2, y_1, y_2\}$ is also a tight frame for \mathbb{R}^2 with bound $\frac{3}{2}$.

From Theorem 2.4, the following result is straightforward.

COROLLARY 2.5. *Let $X = \{x_i\}_{i=1}^m$ be a tight frame with bound A for \mathcal{H} and let $y \in \mathcal{H}$. A $(r, 1)$ -surgery on X with y resulting in a tight frame with bound A if and only if*

$$\sum_{i=m-r+1}^m \|x_i\|^2 = \|y\|^2 \quad \text{and} \quad \text{span}\{y\} = \text{span}\{x_i\}_{i=m-r+1}^m.$$

3. Surgery for full spark frames

DEFINITION 3.1. [1] A sequence $\{x_i\}_{i=1}^m$ is called a full spark frame for \mathcal{H} , if for any $\sigma \subset \{1, \dots, m\}$ with $|\sigma| = n$, $\{x_i\}_{i \in \sigma}$ is a frame for \mathcal{H} .

The following proposition is straightforward but important.

PROPOSITION 3.1. *A sequence $X = \{x_i\}_{i=1}^m$ is a full spark frame for \mathcal{H} if and only if every n elements of X are linearly independent.*

We can see that a frame is full spark if any n of its members make up a basis for \mathcal{H} . Hence it is possible to take surgery on a basis resulting in a full spark frame.

PROPOSITION 3.2. *A $(r, 0)$ -surgery on a full spark frame results a full spark frame if $r \leq m - n$.*

Proof. Let $X = \{x_i\}_{i=1}^m$ be a full spark frame, by Proposition 3.1, every n elements of X are linearly independent. If $r \leq m - n$, then every n elements of $\{x_i\}_{i=1}^{m-r}$ are linearly independent, and then $\text{span}\{x_i\}_{i=1}^{m-r} = \mathcal{H}$. Thus $\{x_i\}_{i=1}^{m-r}$ is a full spark frame for \mathcal{H} . \square

LEMMA 3.1. *A (r, k) -surgery on a frame $\{x_i\}_{i=1}^m$ resulting in a full spark frame is always possible if the $\{x_i\}_{i=1}^{m-r}$ is linear independent and $k \geq n + r - m$.*

Proof. Since any linear independent set can be expanded to a basis for \mathcal{H} , and $k \geq n + r - m$, then it is possible to expand $\text{span}\{x_i\}_{i=1}^{m-r}$ to a full spark frame. \square

COROLLARY 3.1. *A (r, k) -surgery on a frame $\{x_i\}_{i=1}^m$ resulting in a full spark frame is impossible if the $\{x_i\}_{i=1}^{m-r}$ is a linear dependent set.*

Next, we give a condition such that a (r, k) -surgery on a frame $\{x_i\}_{i=1}^m$ results in a full spark frame.

THEOREM 3.2. *Let $X = \{x_i\}_{i=1}^m$ be a frame and let $Y = \{y_i\}_{i=1}^k \subset \mathcal{H}$. A $(m - n, k)$ -surgery on X with Y resulting in a full spark frame if and only if $T_{X'}$ is invertible and all minors of $T_{X'}^{-1}T_Y$ are nonzero, where $T_{X'}$ and T_Y are the synthesis operators of $\{x_i\}_{i=1}^n$ and $\{y_i\}_{i=1}^k$, respectively.*

Proof. (\Leftarrow) Let F be an $n \times (n+k)$ matrix consisting of $\{x_1, \dots, x_n, y_1, \dots, y_k\}$, thus $F = [T_{X'} | T_Y]$. If we want to prove that $\{x_1, \dots, x_n, y_1, \dots, y_k\}$ is a full spark frame, by Proposition 3.1, it is equivalent to proving that every n vectors of $\{x_1, \dots, x_n, y_1, \dots, y_k\}$ are linearly independent. If $T_{X'}$ is invertible, we have

$$T_{X'}^{-1}F = [E | T_{X'}^{-1}T_Y].$$

Next, we show that every n columns of $T_{X'}^{-1}F = [E | T_{X'}^{-1}T_Y]$ are linearly independent if and only if all minors of $T_{X'}^{-1}T_Y$ are nonzero. The idea of the proof is simple and we shall just illustrate it by proving that the top right $l \times l$ submatrix B of $T_{X'}^{-1}T_Y$ is nonsingular, where $1 \leq l \leq \min\{n, k\}$. Take the matrix F' consisting of the last $n-l$ columns of E and the first l columns of $T_{X'}^{-1}T_Y$:

$$F' = \left[\begin{array}{c|c} 0 & B \\ \hline 1 & \\ & 1 \\ & \ddots \\ & 1 \end{array} \right]. \quad (3.1)$$

Then $\det(F') = \det(B)$. Hence, $\det(F')$ is nonzero if and only if $\det(B)$ is nonzero. This means that every n columns of $T_{X'}^{-1}F = [E | T_{X'}^{-1}T_Y]$ are linearly independent if and only if all minors of $T_{X'}^{-1}T_Y$ are nonzero.

(\Rightarrow) If $\{x_1, \dots, x_n, y_1, \dots, y_k\}$ is a full spark frame for \mathcal{H} , then $T_{X'}$ is invertible. From [16, Lemma], we deduce that $T_{X'}^{-1}F = [E | T_{X'}^{-1}T_Y]$ is also a full spark frame for \mathcal{H} . From (3.1), we know that all minors of $T_{X'}^{-1}T_Y$ are nonzero.

Hence, $\{x_1, \dots, x_n, y_1, \dots, y_k\}$ is a full spark frame for \mathcal{H} if and only if $T_{X'}$ is invertible and all minors of $T_{X'}^{-1}T_Y$ are nonzero. \square

EXAMPLE 2. We now give an example for Theorem 3.2. Let $\{x_i\}_{i=1}^5$ be a frame for \mathbb{R}^3 , where

$$x_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad x_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

Let

$$y_1 = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{3} \\ \frac{1}{4} \end{bmatrix}, \quad y_2 = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{4} \\ \frac{1}{5} \end{bmatrix}, \quad y_3 = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{5} \\ \frac{1}{6} \end{bmatrix}, \quad y_4 = \begin{bmatrix} \frac{1}{5} \\ \frac{1}{6} \\ \frac{1}{7} \end{bmatrix}.$$

It is easy to verify that all minors of T are nonzero, where columns of T are consisting of $\{y_1\}_{i=1}^4$. Then a $(2, 4)$ -surgery on $\{x_i\}_{i=1}^5$ with $\{y_1\}_{i=1}^4$ results in a full spark frame for \mathbb{R}^3 . In fact, we can compute that:

(a) x_i, y_j, y_k are linearly independent, where $i = 1, 2, 3, j, k = 1, 2, 3, 4$ and $j \neq k$;

(b) x_i, x_j, y_k are linearly independent, where $i, j = 1, 2, 3$ and $j \neq i, k = 1, 2, 3, 4$;

(c) y_i, y_j, y_k are linearly independent, where $i, j, k = 1, 2, 3, 4$ and $i \neq j \neq k$;

(d) x_1, x_2, x_3 are linearly independent.

Note that if $r > m - n$ in Theorem 3.2, and $\{x_i\}_{i=1}^{m-r}$ are unit vectors, then the minors of T_Y may be not all non-zero.

EXAMPLE 3. Let $X = \{x_i\}_{i=1}^m$ be a frame for \mathbb{R}^n , where $x_1 = (1, 0 \cdots, 0)^T$ and $x_2 = (0, 1 \cdots, 0)^T$. Let T be a Vandermonde matrix as follow:

$$T = \begin{bmatrix} 1 & 1 & \dots & 1 \\ a_1 & a_2 & \dots & a_k \\ a_1^2 & a_2^2 & \dots & a_k^2 \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{n-1} & a_2^{n-1} & \dots & a_k^{n-1} \end{bmatrix} = [y_1, y_2, \dots, y_k].$$

We will show that a $(m - 2, k)$ -surgery on X with $\{y_i\}_{i=1}^k$ results in a full spark frame if a_1, \dots, a_k are k distinct nonzero elements. Let

$$F = [x_1, x_2, y_1, \dots, y_k] = \begin{bmatrix} 1 & 0 & 1 & 1 & \dots & 1 \\ 0 & 1 & a_1 & a_2 & \dots & a_k \\ 0 & 0 & a_1^2 & a_2^2 & \dots & a_k^2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & a_1^{n-1} & a_2^{n-1} & \dots & a_k^{n-1} \end{bmatrix}.$$

Any $n \times n$ matrix, say U , consists of any n columns of F . If U is a submatrix of T , then $\det U \neq 0$ because a_1, \dots, a_k are k distinct elements. Next, we will prove that the matrix U contains first p ($0 < p \leq 2$) columns of F . Without loss of generality, assume that U consists of first p columns of F and first $n - p$ columns of T .

Case 1. $p = 1$

Assume that U contains x_1 , then

$$\det U = \begin{vmatrix} 1 & 1 & 1 & \dots & 1 \\ 0 & a_1 & a_2 & \dots & a_{n-1} \\ 0 & a_1^2 & a_2^2 & \dots & a_{n-1}^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & a_1^{n-1} & a_2^{n-1} & \dots & a_{n-1}^{n-1} \end{vmatrix} = \prod_{l=1, \dots, n-1} a_l \prod_{\substack{i, j=1, \dots, n-1 \\ i > j}} (a_i - a_j) \neq 0.$$

Assume that U contains x_2 , then

$$\det U = \begin{vmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & a_1 & a_2 & \dots & a_{n-1} \\ 0 & a_1^2 & a_2^2 & \dots & a_{n-1}^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & a_1^{n-1} & a_2^{n-1} & \dots & a_{n-1}^{n-1} \end{vmatrix} = - \sum_{i=1}^{n-1} a_1 \dots a_{i-1} a_{i+1} \dots a_{n-1} \prod_{\substack{i, j=1, \dots, n-1 \\ i > j}} (a_i - a_j) \neq 0.$$

Case 2. $p = 2$

Assume that U contains x_1, x_2 , then

$$\det U = \begin{vmatrix} 1 & 0 & 1 & 1 & \dots & 1 \\ 0 & 1 & a_1 & a_2 & \dots & a_{n-2} \\ 0 & 0 & a_1^2 & a_2^2 & \dots & a_{n-2}^2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & a_1^{n-1} & a_2^{n-1} & \dots & a_{n-2}^{n-1} \end{vmatrix} = \prod_{l=1, \dots, n-2} a_l^2 \prod_{\substack{i,j=1, \dots, n-2 \\ i>j}} (a_i - a_j) \neq 0.$$

Therefore, all $n \times n$ submatrices of F are non-singular. Thus every n vectors of $\{x_1, x_2, y_1, \dots, y_k\}$ are linearly independent. Hence, $\{x_1, x_2, y_1, \dots, y_k\}$ is a full spark frame for \mathcal{H} . But there exists a minor of T is zero. Let $a_2 = -2$ and $a_3 = 2$. We can see that a minor of order 2 of T , such as $\begin{vmatrix} 1 & 1 \\ 4 & 4 \end{vmatrix}$, is zero.

Acknowledgements. The research is supported by the Natural Science Foundation of Anhui Province (1908085MF175) and the Fundamental Research Funds for the Central Universities (JZ2019HGBZ0129).

REFERENCES

- [1] BORIS ALEXEEV, JAMESON CAHILL, AND DUSTIN G. MIXON, *Full spark frames*, Journal of Fourier Analysis and Applications, **18** (6): 1167–1194, 2012.
- [2] MARTIN S. COPENHAVER, YEON HYANG KIM, CORTNEY LOGAN, KYANNE MAYFIELD, SIVARAM K. NARAYAN, AND JONATHAN SHEPHERD, *Maximum robustness and surgery of frames in finite dimensions*, arXiv preprint arXiv:1303.1163, 2013.
- [3] DEGUANG HAN, QIANFENG HU, AND RUI LIU, *Injective continuous frames and quantum detections*, Banach Journal of Mathematical Analysis, **15** (1): 1–24, 2021.
- [4] DEGUANG HAN, KERI KORNELSON, ERIC WEBER, AND DAVID LARSON, *Frames for undergraduates*, vol. **40**, American Mathematical Soc., 2007.
- [5] DEGUANG HAN, PENG TONG LI, BIN MENG, AND WAISHING TANG, *Operator valued frames and structured quantum channels*, Science China Mathematics, **54** (11): 2361–2372, 2011.
- [6] DEGUANG HAN, AND WENCHANG SUN, *Reconstruction of signals from frame coefficients with erasures at unknown locations*, IEEE transactions on information theory, **60** (7): 4013–4025, 2014.
- [7] ROGER A. HORN, AND CHARLES R. JOHNSON, *Matrix analysis*, Cambridge university press, 2012.
- [8] JELENA KOVACEVIC, PIER LUIGI DRAGOTTI, AND VIVEK K. GOYAL, *Filter bank frame expansions with erasures*, IEEE Transactions on Information Theory, **48** (6): 1439–1450, 2002.
- [9] GITTA KUTYNIK, KASSO A. OKOUDJOU, FRIEDRICH PHILIPP, AND TULEY, ELIZABETH K, *Scalable frames*, Linear Algebra and its Applications, **438** (5): 2225–2238, 2013.
- [10] DENG FENG LI, AND WEN CHANG SUN, *Expansion of frames to tight frames*, Acta Mathematica Sinica, English Series, **25** (2): 287–292, 2009.
- [11] DONGWEI LI, JINSONG LENG, TINGZHU HUANG, AND QING GAO, *Frame expansions with probabilistic erasures*, Digital Signal Processing, **72**: 75–82, 2018.
- [12] P. G. MASSEY, AND M. A. RUIZ, *Tight frame completions with prescribed norms*, Sampling Theory in Signal and Image Processing, **7** (1): 1–14, 2008.
- [13] R. RAMU NAIDU, AND CHANDRA R. MURTHY, *Construction of unimodular tight frames for compressed sensing using majorization-minimization*, Signal Processing, **172**: 107516, 2020.
- [14] SIVARAM K. NARAYAN, EILEEN L. RADZWION, SARA P. RIMER, RACHAEL L. TOMASINO, JENNIFER L. WOLFE, AND ANDREW M. ZIMMER, *Robustness and surgery of frames*, Linear algebra and its applications, **434** (8): 1893–1901, 2011.
- [15] NGA QUYNH NGUYEN, *Surgery on frames*, Texas A&M University, 2008.

- [16] MARKUS PUSCHEL, AND JELENA KOVACEVIC, *Real, tight frames with maximal robustness to erasures*, in Data Compression Conference, pages 63–72, IEEE, 2005.
- [17] URI SHAHAM, ALEXANDER CLONINGER, AND RONALD R. COIFMAN, *Provable approximation properties for deep neural networks*, Applied and Computational Harmonic Analysis, **44** (3): 537–557, 2018.

(Received May 1, 2021)

Dongwei Li
School of Mathematics
HeFei University of Technology
230009, P. R. China
e-mail: dongweili@hufu.edu.cn

Jing Jiang
School of Mathematics
HeFei University of Technology
230009, P. R. China
e-mail: jiangjing20121449@163.com