

NUMERICAL RANGES OF SOME FOGUEL OPERATORS

MUYAN JIANG AND ILYA M. SPITKOVSKY

Abstract. The Foguel operator is defined as $F_T = \begin{bmatrix} S^* & T \\ 0 & S \end{bmatrix}$, where S is the right shift on a Hilbert space \mathcal{H} and T is an arbitrary bounded linear operator acting on \mathcal{H} . Obviously, the numerical range $W(F_0)$ of F_T with $T = 0$ is the open unit disk, and it was suggested by Gau, Wang and Wu in their Linear Algebra and Applications (2021) paper that $W(F_{aI})$ for non-zero $a \in \mathbb{C}$ might be an elliptical disk. In this paper, we described $W(F_{aI})$ explicitly and, as it happens, it is not.

Mathematics subject classification (2020): 47A12, 30H10, 47A08, 47B02, 47B35, 47B37.

Keywords and phrases: Numerical range, Foguel operator, Toeplitz operator, Schur complement.

REFERENCES

- [1] A. BÖTTCHER AND B. SILBERMANN, *Analysis of Toeplitz operators*, second ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, Prepared jointly with A. Karlovich.
- [2] H.-L. GAU, K.-Z. WANG, AND P. Y. WU, *Numerical ranges of Foguel operators*, Linear Algebra Appl. **610** (2021), 766–784.
- [3] N. NIKOLSKI, *Hardy spaces*, French ed., Cambridge Studies in Advanced Mathematics, vol. 179, Cambridge University Press, Cambridge, 2019.
- [4] P. Y. WU AND H.-L. GAU, *Numerical ranges of Hilbert space operators*, Encyclopedia of Mathematics and its Applications, vol. 179, Cambridge University Press, Cambridge, 2021.