

## MAXIMAL NUMERICAL RANGE OF THE BIMULTIPLICATION $M_{2,A,B}$

EL HASSAN BENABDI, MOHAMED KAADOUD CHRAIBI  
AND ABDERRAHIM BAGHDAD \*

**Abstract.** Let  $\mathcal{B}(\mathcal{H})$  denote the  $C^*$ -algebra of all bounded linear operators acting on a complex Hilbert space  $\mathcal{H}$ . For  $A, B \in \mathcal{B}(\mathcal{H})$ , define the bimultiplication operator  $M_{2,A,B}$  on the class of Hilbert-Schmidt operators by  $M_{2,A,B}(X) = AXB$ . It is known [5] that if either  $A$  or  $B$  is hyponormal, then

$$\overline{W(M_{2,A,B})} = \overline{\text{co}(W(A)W(B))},$$

where the bar and  $\text{co}$  stand for the closure and the convex hull, respectively and  $W(\cdot)$  denotes the numerical range. In this paper, we give some conditions satisfied by  $A$  and  $B$  to have the following equality

$$W_0(M_{2,A,B}) = \text{co}(W_0(A)W_0(B)),$$

where  $W_0(\cdot)$  denotes the maximal numerical range.

*Mathematics subject classification (2020):* 47A12, 47B20, 47B47.

*Keywords and phrases:* Maximal numerical range, Hilbert-Schmidt operators, bimultiplication.

## REFERENCES

- [1] A. BAGHDAD AND M. C. KAADOUD, *On the maximal numerical range of a hyponormal operator*, Oper. Matrices **13** (4) (2019), 1163–1171.
- [2] F. F. BONSALL AND J. DUNCAN, *Numerical ranges of operators on normed spaces and of elements of normed algebras*, London-New York: Cambridge University Press; (London mathematical society lecture note series; 2), (1971).
- [3] F. F. BONSALL AND J. DUNCAN, *Numerical ranges II*, New York-London, Cambridge University Press; (London mathematical society lecture notes series; 10), (1973).
- [4] R. BOULDIN, *The numerical range of a product*, J. Math. Anal. Appl. **32** (1970), 459–467.
- [5] M. BOUMAZGOUR, H. A. NABWEY, *A note concerning the numerical range of a basic elementary operator*, Ann. Funct. Ann. **7** (3) (2016), 434–441.
- [6] J. B. CONWAY, *A Course In Operator Theory*, American Mathematical Society, (2000).
- [7] L. FIALKOW, *Structural properties of elementary operators*, Proc. Inter. Workshop, (1991).
- [8] C. K. FONG, *On the essential maximal numerical range*, Acta Sci. Math. **41** (1979), 307–315.
- [9] H. L. GAU, K. Z. WANG, P. Y. WU, *Numerical radii for tensor products of operators*, Integral Equ Oper Theory **78** (3) (2014), 375–382.
- [10] K. E. GUSTAFSON AND D. K. M. RAO, *Numerical range: The field of values of linear operators and matrices*, Springer, New York, Inc, (1997).
- [11] P. R. HALMOS, *A Hilbert space problem book*, New York: Van Nostrand, (1967).
- [12] G. JI, N. LIU AND Z. E. LI, *Essential numerical range and maximal numerical range of the Aluthge transform*, Linear Multilinear Algebra **55** (4) (2007), 315–322.
- [13] M. C. KAADOUD, *Dominé numérique de l'opérateur produit  $M_{2,A,B}$  et de la dérivation généralisée  $\delta_{2,A,B}$* , Extracta Mathematicae **17** (1) (2002), 59–68.
- [14] M. MATHIEU, *Spectral theory for multiplication operators on  $C^*$ -algebras*, Proc. Royal Irish Acad. **83** A (2) (1983), 231–249.
- [15] I. M. SPITKOVSKY, *A note on the maximal numerical range*, Oper. Matrices **13** (3) (2019), 601–605.
- [16] J. G. STAMPFLI, *The norm of derivation*, Pacific J. Math. **33** (1970), 737–747.