

## A NORM INEQUALITY FOR THREE REAL MATRICES

FEN WANG

(Communicated by S. Fallat)

*Abstract.* In this paper, we prove a norm inequality for three real  $2 \times 2$  matrices conjectured by L. László [3] recently, which is a generalization of the famous Böttcher-Wenzel inequality.

### 1. Introduction

Let  $M_n(\mathbb{R})$  be the set of real  $n \times n$  matrices. For  $A, B \in M_n(\mathbb{R})$ , Böttcher-Wenzel [1] conjectured

$$\|AB - BA\|_F^2 \leq 2\|A\|_F^2\|B\|_F^2, \quad (1.1)$$

where  $\|\cdot\|_F$  means the Frobenius norm. This conjecture was proved by many authors, please see the survey [2] and the references therein. Motivated by (1.1), László [3] considered three real  $n \times n$  matrices and proved a new norm inequality as follows.

**THEOREM 1.1.** [3, Theorem 3.1] *If  $A, B, C \in M_n(\mathbb{R})$ , then*

$$\|ABC - CBA\|_F^2 \leq \|A\|_F^2\|B\|_F^2\|C\|_F^2 - \|B\|_F^2 \operatorname{tr}^2(A^T C). \quad (1.2)$$

At the same time, László [3] introduced the commutator for  $A, B, C \in M_n(\mathbb{R})$ :

$$D := (ABC + BCA + CAB) - (CBA + ACB + BAC),$$

and proposed the following conjecture:

**CONJECTURE 1.2.** [3] For any  $A, B, C \in M_n(\mathbb{R})$ , it holds

$$\|D\|_F^2 \leq \frac{3}{2} \left( \|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2 \right).$$

László [3] himself verified the conjecture for  $n = 2$  and  $A, B, C$  being upper triangular. In this paper, we will consider the László's conjecture for  $n = 2$ . To be precise, we will prove the following theorem.

**THEOREM 1.3.** *Let  $A, B, C \in M_2(\mathbb{R})$  and one of them be symmetric. Then*

$$\|D\|_F^2 \leq \frac{3}{2} \left( \|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2 \right). \quad (1.3)$$

We will also show that the constant  $\frac{3}{2}$  is optimal.

---

*Mathematics subject classification* (2020): 15A60, 15A45.

*Keywords and phrases:* Frobenius norm, Böttcher-Wenzel inequality, norm inequality.

The author was supported by the Foundation of Hubei Provincial Department of Education (No. Q20233003), the Scientific Research Fund of Hubei Provincial Department of Education (No. B2022207) and Hubei University of Education, Bigdata Modeling and Intelligent Computing Research Institute.

## 2. Proof of Theorem 1.3

*Proof of Theorem 1.3.* Step 1: Without loss of generality, we can assume

$$A = \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix}, \quad B = \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix}, \quad C = \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix},$$

where  $a_i, b_i, \lambda_j \in \mathbb{R}$  for  $1 \leq i \leq 4$ ,  $1 \leq j \leq 2$ .

Step 2: Computation of  $AB - BA$ ,  $BC - CB$ ,  $CA - AC$  and their norms.

$$\begin{aligned} AB - BA &= \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} - \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \\ &= \begin{pmatrix} a_1b_1 + a_2b_3 & a_1b_2 + a_2b_4 \\ a_3b_1 + a_4b_3 & a_3b_2 + a_4b_4 \end{pmatrix} - \begin{pmatrix} a_1b_1 + a_3b_2 & a_2b_1 + a_4b_2 \\ a_1b_3 + a_3b_4 & a_2b_3 + a_4b_4 \end{pmatrix} \\ &= \begin{pmatrix} a_2b_3 - a_3b_2 & (a_1b_2 - a_2b_1) + (a_2b_4 - a_4b_2) \\ (a_3b_1 - a_1b_3) + (a_4b_3 - a_3b_4) & -(a_2b_3 - a_3b_2) \end{pmatrix}, \end{aligned} \quad (2.1)$$

$$\begin{aligned} BC - CB &= \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix} - \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix} \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1b_1 & \lambda_2b_2 \\ \lambda_1b_3 & \lambda_2b_4 \end{pmatrix} - \begin{pmatrix} \lambda_1b_1 & \lambda_1b_2 \\ \lambda_2b_3 & \lambda_2b_4 \end{pmatrix} \\ &= (\lambda_1 - \lambda_2) \begin{pmatrix} 0 & -b_2 \\ b_3 & 0 \end{pmatrix}, \end{aligned} \quad (2.2)$$

$$\begin{aligned} CA - AC &= \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix} \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} - \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1a_1 & \lambda_1a_2 \\ \lambda_2a_3 & \lambda_2a_4 \end{pmatrix} - \begin{pmatrix} \lambda_1a_1 & \lambda_2a_2 \\ \lambda_1a_3 & \lambda_2a_4 \end{pmatrix} \\ &= (\lambda_1 - \lambda_2) \begin{pmatrix} 0 & a_2 \\ -a_3 & 0 \end{pmatrix}. \end{aligned} \quad (2.3)$$

Therefore,

$$\begin{aligned}\|AB - BA\|_F^2 &= 2(a_2b_3 - a_3b_2)^2 + [(a_1b_2 - a_2b_1) + (a_2b_4 - a_4b_2)]^2 \\ &\quad + [(a_3b_1 - a_1b_3) + (a_4b_3 - a_3b_4)]^2, \\ \|BC - CB\|_F^2 &= (b_2^2 + b_3^2)(\lambda_1 - \lambda_2)^2, \\ \|CA - AC\|_F^2 &= (a_2^2 + a_3^2)(\lambda_1 - \lambda_2)^2.\end{aligned}\tag{2.4}$$

*Step 3:* Computation of  $C(AB - BA)$ ,  $A(BC - CB)$ ,  $B(CA - AC)$ .

By (2.1), we have

$$\begin{aligned}C(AB - BA) &= \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \begin{pmatrix} a_2b_3 - a_3b_2 & (a_1b_2 - a_2b_1) + (a_2b_4 - a_4b_2) \\ (a_3b_1 - a_1b_3) + (a_4b_3 - a_3b_4) & -(a_2b_3 - a_3b_2) \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1(a_2b_3 - a_3b_2) & \lambda_1(a_1b_2 - a_2b_1) + \lambda_1(a_2b_4 - a_4b_2) \\ \lambda_2(a_3b_1 - a_1b_3) + \lambda_2(a_4b_3 - a_3b_4) & -\lambda_2(a_2b_3 - a_3b_2) \end{pmatrix}.\end{aligned}\tag{2.5}$$

By (2.2), we have

$$A(BC - CB) = \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} (\lambda_1 - \lambda_2) \begin{pmatrix} 0 & -b_2 \\ b_3 & 0 \end{pmatrix} = (\lambda_1 - \lambda_2) \begin{pmatrix} a_2b_3 & -a_1b_2 \\ a_4b_3 & -a_3b_2 \end{pmatrix}.\tag{2.6}$$

By (2.3), we have

$$B(CA - AC) = \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} (\lambda_1 - \lambda_2) \begin{pmatrix} 0 & a_2 \\ -a_3 & 0 \end{pmatrix} = (\lambda_1 - \lambda_2) \begin{pmatrix} -a_3b_2 & a_2b_1 \\ -a_3b_4 & a_2b_3 \end{pmatrix}.\tag{2.7}$$

*Step 4:* Computation of  $\|D\|_F^2$ .

By (2.5)–(2.7), we get

$$\begin{aligned}D &= C(AB - BA) + A(BC - CB) + B(CA - AC) \\ &= \begin{pmatrix} (2\lambda_1 - \lambda_2)(a_2b_3 - a_3b_2) & \lambda_2(a_1b_2 - a_2b_1) + \lambda_1(a_2b_4 - a_4b_2) \\ \lambda_2(a_3b_1 - a_1b_3) + \lambda_1(a_4b_3 - a_3b_4) & (\lambda_1 - 2\lambda_2)(a_2b_3 - a_3b_2) \end{pmatrix},\end{aligned}\tag{2.8}$$

therefore,

$$\begin{aligned}
\|D\|_F^2 &= [(2\lambda_1 - \lambda_2)(a_2b_3 - a_3b_2)]^2 + [(\lambda_1 - 2\lambda_2)(a_2b_3 - a_3b_2)]^2 \\
&\quad + [\lambda_2(a_1b_2 - a_2b_1) + \lambda_1(a_2b_4 - a_4b_2)]^2 \\
&\quad + [\lambda_2(a_3b_1 - a_1b_3) + \lambda_1(a_4b_3 - a_3b_4)]^2 \\
&= [(2\lambda_1 - \lambda_2)^2 + (\lambda_1 - 2\lambda_2)^2](a_2b_3 - a_3b_2)^2 \\
&\quad + [\lambda_2(a_1b_2 - a_2b_1) + \lambda_1(a_2b_4 - a_4b_2)]^2 \\
&\quad + [\lambda_2(a_3b_1 - a_1b_3) + \lambda_1(a_4b_3 - a_3b_4)]^2 \\
&= (5\lambda_1^2 + 5\lambda_2^2 - 8\lambda_1\lambda_2)(a_2b_3 - a_3b_2)^2 \\
&\quad + [(a_2b_4 - a_4b_2)^2 + (a_4b_3 - a_3b_4)^2]\lambda_1^2 \\
&\quad + [(a_1b_2 - a_2b_1)^2 + (a_3b_1 - a_1b_3)^2]\lambda_2^2 \\
&\quad + [2(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2) + 2(a_3b_1 - a_1b_3)(a_4b_3 - a_3b_4)]\lambda_1\lambda_2 \\
&= [5(a_2b_3 - a_3b_2)^2 + (a_2b_4 - a_4b_2)^2 + (a_4b_3 - a_3b_4)^2]\lambda_1^2 \\
&\quad + [5(a_2b_3 - a_3b_2)^2 + (a_1b_2 - a_2b_1)^2 + (a_3b_1 - a_1b_3)^2]\lambda_2^2 \\
&\quad + [2(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2) + 2(a_3b_1 - a_1b_3)(a_4b_3 - a_3b_4) \\
&\quad - 8(a_2b_3 - a_3b_2)^2]\lambda_1\lambda_2
\end{aligned} \tag{2.9}$$

*Step 5:* Computation of

$$\|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2.$$

By (2.4), we get

$$\begin{aligned}
&\|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2 \\
&= \|A\|_F^2(b_2^2 + b_3^2)(\lambda_1 - \lambda_2)^2 + \|B\|_F^2(a_2^2 + a_3^2)(\lambda_1 - \lambda_2)^2 \\
&\quad + (\lambda_1^2 + \lambda_2^2)\|AB - BA\|_F^2 \\
&= [\|A\|_F^2(b_2^2 + b_3^2) + \|B\|_F^2(a_2^2 + a_3^2) + \|AB - BA\|_F^2]\lambda_1^2 \\
&\quad + [\|A\|_F^2(b_2^2 + b_3^2) + \|B\|_F^2(a_2^2 + a_3^2) + \|AB - BA\|_F^2]\lambda_2^2 \\
&\quad - 2[\|A\|_F^2(b_2^2 + b_3^2) + \|B\|_F^2(a_2^2 + a_3^2)]\lambda_1\lambda_2.
\end{aligned} \tag{2.10}$$

*Step 6:* Computation of

$$3(\|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2) - 2\|D\|_F^2.$$

Denote

$$\begin{aligned}\mathbf{I}_1 &= 3 \left[ \|A\|_F^2 (b_2^2 + b_3^2) + \|B\|_F^2 (a_2^2 + a_3^2) \right], \\ \mathbf{J}_1 &= 5(a_2 b_3 - a_3 b_2)^2 + (a_2 b_4 - a_4 b_2)^2 + (a_4 b_3 - a_3 b_4)^2, \\ \mathbf{J}_2 &= 5(a_2 b_3 - a_3 b_2)^2 + (a_1 b_2 - a_2 b_1)^2 + (a_3 b_1 - a_1 b_3)^2, \\ \mathbf{J}_3 &= 2(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 2(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4) - 8(a_2 b_3 - a_3 b_2)^2.\end{aligned}\tag{2.11}$$

Then, from (2.9)–(2.11), we obtain that

$$\begin{aligned}3 (\|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2) - 2 \|D\|_F^2 \\ = (\mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_1) \lambda_1^2 - 2(\mathbf{I}_1 + \mathbf{J}_3) \lambda_1 \lambda_2 + (\mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_2) \lambda_2^2 \\ = p \lambda_1^2 + q \lambda_1 \lambda_2 + r \lambda_2^2,\end{aligned}\tag{2.12}$$

where

$$\begin{aligned}p &= \mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_1, \\ q &= -2(\mathbf{I}_1 + \mathbf{J}_3), \\ r &= \mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_2.\end{aligned}\tag{2.13}$$

*Step 7:* Estimation of  $p$  and computation of  $\Delta = q^2 - 4pr$ .

(1) Estimation of  $p$ . By (2.4), (2.11) and (2.13), we have

$$\begin{aligned}p &= \mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_1 \\ &= 3 \left[ (a_1^2 + a_2^2 + a_3^2 + a_4^2) (b_2^2 + b_3^2) + (b_1^2 + b_2^2 + b_3^2 + b_4^2) (a_2^2 + a_3^2) \right] \\ &\quad + 3 [2(a_2 b_3 - a_3 b_2)^2 + (a_1 b_2 - a_2 b_1)^2 + (a_2 b_4 - a_4 b_2)^2 + (a_3 b_1 - a_1 b_3)^2 \\ &\quad + (a_4 b_3 - a_3 b_4)^2 + 2(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 2(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)] \\ &\quad - 2 [5(a_2 b_3 - a_3 b_2)^2 + (a_2 b_4 - a_4 b_2)^2 + (a_4 b_3 - a_3 b_4)^2] \\ &= 6(a_2^2 + a_3^2)(b_2^2 + b_3^2) - 4(a_2 b_3 - a_3 b_2)^2 \\ &\quad + 3(a_1^2 b_2^2 + a_2^2 b_1^2) + 3(a_1 b_2 - a_2 b_1)^2 + 3(a_2^2 b_4^2 + a_4^2 b_2^2) + (a_2 b_4 - a_4 b_2)^2 \\ &\quad + 6(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 3(a_1^2 b_3^2 + a_3^2 b_1^2) + 3(a_1 b_3 - a_3 b_1)^2 \\ &\quad + 3(a_3^2 b_4^2 + a_4^2 b_3^2) + (a_3 b_4 - a_4 b_3)^2 + 6(a_1 b_3 - a_3 b_1)(a_3 b_4 - a_4 b_3) \\ &\geq 2(a_2^2 + a_3^2)(b_2^2 + b_3^2) \\ &\quad + \frac{9}{2}(a_1 b_2 - a_2 b_1)^2 + \frac{5}{2}(a_2 b_4 - a_4 b_2)^2 + 6(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) \\ &\quad + \frac{9}{2}(a_1 b_3 - a_3 b_1)^2 + \frac{5}{2}(a_3 b_4 - a_4 b_3)^2 + 6(a_1 b_3 - a_3 b_1)(a_3 b_4 - a_4 b_3)\end{aligned}$$

$$\begin{aligned}
&\geq 2(a_2^2 + a_3^2)(b_2^2 + b_3^2) \\
&\quad + 2 \cdot \sqrt{\frac{9}{2}(a_1b_2 - a_2b_1)^2 \cdot \frac{5}{2}(a_2b_4 - a_4b_2)^2 + 6(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2)} \\
&\quad + 2 \cdot \sqrt{\frac{9}{2}(a_1b_3 - a_3b_1)^2 \cdot \frac{5}{2}(a_3b_4 - a_4b_3)^2 + 6(a_1b_3 - a_3b_1)(a_3b_4 - a_4b_3)} \\
&\qquad\qquad\qquad (\text{by AM-GM Inequality}) \\
&= 2(a_2^2 + a_3^2)(b_2^2 + b_3^2) \\
&\quad + \sqrt{45}|(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2)| + 6(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2) \\
&\quad + \sqrt{45}|(a_1b_3 - a_3b_1)(a_3b_4 - a_4b_3)| + 6(a_1b_3 - a_3b_1)(a_3b_4 - a_4b_3) \\
&\geq 0,
\end{aligned}$$

where the first inequality follows from

$$\begin{aligned}
(a_2^2 + a_3^2)(b_2^2 + b_3^2) - (a_2b_3 - a_3b_2)^2 &= (a_2b_2 + a_3b_3)^2 \geq 0, \\
2(a_1^2b_2^2 + a_2^2b_1^2) - (a_1b_2 - a_2b_1)^2 &= (a_1b_2 + a_2b_1)^2 \geq 0, \\
2(a_1^2b_3^2 + a_3^2b_1^2) - (a_1b_3 - a_3b_1)^2 &= (a_1b_3 + a_3b_1)^2 \geq 0, \\
2(a_2^2b_4^2 + a_4^2b_2^2) - (a_2b_4 - a_4b_2)^2 &= (a_2b_4 + a_4b_2)^2 \geq 0, \\
2(a_3^2b_4^2 + a_4^2b_3^2) - (a_3b_4 - a_4b_3)^2 &= (a_3b_4 + a_4b_3)^2 \geq 0.
\end{aligned}$$

If  $p = 0$ , it is easy to check from all inequalities in the estimation of  $p$  that

$$\begin{aligned}
(a_2^2 + a_3^2)(b_2^2 + b_3^2) &= 0, \\
a_1b_2 + a_2b_1 &= 0, \\
a_1b_3 + a_3b_1 &= 0, \\
a_2b_4 + a_4b_2 &= 0, \\
a_3b_4 + a_4b_3 &= 0,
\end{aligned}$$

which implies

$$\begin{aligned}
(a_2^2 + a_3^2)(b_2^2 + b_3^2) &= 0, \\
(b_1^2 + b_4^2)(a_2^2 + a_3^2) &= (a_1^2 + a_4^2)(b_2^2 + b_3^2).
\end{aligned}$$

Then it follows that  $A = O$  or  $B = O$  or  $a_2 = a_3 = b_2 = b_3 = 0$ . Whatever,  $q = r = 0$ .

(2) Computation of  $\Delta = q^2 - 4pr$ . By (2.13), we have

$$\begin{aligned} \frac{1}{4}\Delta &= \frac{1}{4}(q^2 - 4pr) \\ &= (\mathbf{I}_1 + \mathbf{J}_3)^2 - (\mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_1) \cdot (\mathbf{I}_1 + 3\|AB - BA\|_F^2 - 2\mathbf{J}_2) \quad (2.14) \\ &= \mathbf{I}_1 \cdot (2\mathbf{J}_1 + 2\mathbf{J}_2 + 2\mathbf{J}_3 - 6\|AB - BA\|_F^2) - 9\|AB - BA\|^4 \\ &\quad + 6\|AB - BA\|_F^2(\mathbf{J}_1 + \mathbf{J}_2) + \mathbf{J}_3^2 - 4\mathbf{J}_1\mathbf{J}_2. \end{aligned}$$

Denote

$$\begin{aligned} \mathbf{I}_2 &= 2\mathbf{J}_1 + 2\mathbf{J}_2 + 2\mathbf{J}_3 - 6\|AB - BA\|_F^2 \\ \mathbf{II} &= 9\|AB - BA\|^4 \\ \mathbf{III} &= 6\|AB - BA\|_F^2(\mathbf{J}_1 + \mathbf{J}_2) \quad (2.15) \\ \mathbf{IV} &= \mathbf{J}_3^2 \\ \mathbf{V} &= 4\mathbf{J}_1\mathbf{J}_2, \end{aligned}$$

then

$$\frac{1}{4}\Delta = (\mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III}) + (\mathbf{IV} - \mathbf{V}). \quad (2.16)$$

*Step 8:* Estimation of  $\mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III}$ .

First, we compute  $\mathbf{I}_2$ . From (2.15), (2.11) and (2.4), we obtain

$$\begin{aligned} \mathbf{I}_2 &= 2\mathbf{J}_1 + 2\mathbf{J}_2 + 2\mathbf{J}_3 - 6\|AB - BA\|_F^2 \\ &= 2[2(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2) \\ &\quad + 2(a_3b_1 - a_1b_3)(a_4b_3 - a_3b_4) - 8(a_2b_3 - a_3b_2)^2] \\ &\quad + 2[5(a_2b_3 - a_3b_2)^2 + (a_2b_4 - a_4b_2)^2 + (a_4b_3 - a_3b_4)^2] \\ &\quad + 2[5(a_2b_3 - a_3b_2)^2 + (a_1b_2 - a_2b_1)^2 + (a_3b_1 - a_1b_3)^2] - 6\|AB - BA\|_F^2 \\ &= 4(a_2b_3 - a_3b_2)^2 + 2(a_2b_4 - a_4b_2)^2 + 2(a_1b_2 - a_2b_1)^2 \\ &\quad + 4(a_1b_2 - a_2b_1)(a_2b_4 - a_4b_2) + 2(a_4b_3 - a_3b_4)^2 + 2(a_3b_1 - a_1b_3)^2 \\ &\quad + 4(a_3b_1 - a_1b_3)(a_4b_3 - a_3b_4) - 6\|AB - BA\|_F^2 \\ &= 2\|AB - BA\|_F^2 - 6\|AB - BA\|_F^2 \\ &= -4\|AB - BA\|_F^2. \quad (2.17) \end{aligned}$$

Next, we give the estimation of  $\mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III}$ . From (2.11), (2.15) and (2.17),

we obtain

$$\begin{aligned}
& \mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III} \\
&= 3 [\|A\|_F^2 (b_2^2 + b_3^2) + \|B\|_F^2 (a_2^2 + a_3^2)] \cdot (-4\|AB - BA\|_F^2) \\
&\quad - 9\|AB - BA\|^4 + 6\|AB - BA\|_F^2 (\mathbf{J}_1 + \mathbf{J}_2) \\
&= 3\|AB - BA\|_F^2 \cdot [-4\|A\|_F^2 (b_2^2 + b_3^2) - 4\|B\|_F^2 (a_2^2 + a_3^2) \\
&\quad - 3\|AB - BA\|_F^2 + 2(\mathbf{J}_1 + \mathbf{J}_2)].
\end{aligned} \tag{2.18}$$

$$\begin{aligned}
& -4\|A\|_F^2 (b_2^2 + b_3^2) - 4\|B\|_F^2 (a_2^2 + a_3^2) - 3\|AB - BA\|_F^2 + 2(\mathbf{J}_1 + \mathbf{J}_2) \\
&= -4(a_1^2 + a_2^2 + a_3^2 + a_4^2)(b_2^2 + b_3^2) - 4(b_1^2 + b_2^2 + b_3^2 + b_4^2)(a_2^2 + a_3^2) \\
&\quad - 3\|AB - BA\|_F^2 + 20(a_2b_3 - a_3b_2)^2 + 2(a_2b_4 - a_4b_2)^2 \\
&\quad + 2(a_4b_3 - a_3b_4)^2 + 2(a_1b_2 - a_2b_1)^2 + 2(a_3b_1 - a_1b_3)^2 \\
&= [-4(a_1^2 + a_4^2)(b_2^2 + b_3^2) - 4(b_1^2 + b_4^2)(a_2^2 + a_3^2) + 2(a_2b_4 - a_4b_2)^2 \\
&\quad + 2(a_4b_3 - a_3b_4)^2 + 2(a_1b_2 - a_2b_1)^2 + 2(a_3b_1 - a_1b_3)^2] \\
&\quad + [-3\|AB - BA\|_F^2 - 8(a_2^2 + a_3^2)(b_2^2 + b_3^2) + 20(a_2b_3 - a_3b_2)^2] \\
&= : \mathbf{K}_1 + \mathbf{K}_2,
\end{aligned} \tag{2.19}$$

where

$$\begin{aligned}
\mathbf{K}_1 &= -4(a_1^2 + a_4^2)(b_2^2 + b_3^2) - 4(b_1^2 + b_4^2)(a_2^2 + a_3^2) + 2(a_2b_4 - a_4b_2)^2 \\
&\quad + 2(a_4b_3 - a_3b_4)^2 + 2(a_1b_2 - a_2b_1)^2 + 2(a_3b_1 - a_1b_3)^2 \\
&\leq -4(a_1^2 + a_4^2)(b_2^2 + b_3^2) - 4(b_1^2 + b_4^2)(a_2^2 + a_3^2) + 4(a_2^2b_4^2 + a_4^2b_2^2) \\
&\quad + 4(a_4^2b_3^2 + a_3^2b_4^2) + 4(a_1^2b_2^2 + a_2^2b_1^2) + 4(a_3^2b_1^2 + a_1^2b_3^2) \\
&= 0
\end{aligned} \tag{2.20}$$

and

$$\begin{aligned}
\mathbf{K}_2 &= -3\|AB - BA\|_F^2 - 8(a_2^2 + a_3^2)(b_2^2 + b_3^2) + 20(a_2b_3 - a_3b_2)^2 \\
&\leq -6(a_2b_3 - a_3b_2)^2 - 8(a_2^2 + a_3^2)(b_2^2 + b_3^2) + 20(a_2b_3 - a_3b_2)^2 \\
&= -8(a_2^2 + a_3^2)(b_2^2 + b_3^2) + 14(a_2b_3 - a_3b_2)^2 \\
&\leq 6(a_2b_3 - a_3b_2)^2, \quad (\text{by AM-GM Inequality})
\end{aligned} \tag{2.21}$$

Therefore, by (2.19), (2.20) and (2.21), we get

$$\begin{aligned}
& \mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III} \\
& \leq 18 \|AB - BA\|_F^2 \cdot (a_2 b_3 - a_3 b_2)^2 \\
& = 18 [2(a_2 b_3 - a_3 b_2)^2 + (a_1 b_2 - a_2 b_1)^2 + (a_2 b_4 - a_4 b_2)^2 \\
& \quad + (a_3 b_1 - a_1 b_3)^2 + (a_4 b_3 - a_3 b_4)^2 \\
& \quad + 2(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) \\
& \quad + 2(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)] \cdot (a_2 b_3 - a_3 b_2)^2 \\
& = 36(a_2 b_3 - a_3 b_2)^4 \\
& \quad + (a_2 b_3 - a_3 b_2)^2 [18(a_1 b_2 - a_2 b_1)^2 + 18(a_2 b_4 - a_4 b_2)^2 \\
& \quad + 18(a_3 b_1 - a_1 b_3)^2 + 18(a_4 b_3 - a_3 b_4)^2 \\
& \quad + 36(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) \\
& \quad + 36(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)].
\end{aligned} \tag{2.22}$$

*Step 9:* Estimation of  $\mathbf{IV} - \mathbf{V}$ .

From (2.15) and (2.11), we obtain

$$\begin{aligned}
\mathbf{IV} &= \mathbf{J}_3^2 \\
&= [(2(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 2(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4) \\
&\quad - 8(a_2 b_3 - a_3 b_2)^2)]^2 \\
&= 64(a_2 b_3 - a_3 b_2)^4 + 4(a_1 b_2 - a_2 b_1)^2 (a_2 b_4 - a_4 b_2)^2 \\
&\quad + 4(a_3 b_1 - a_1 b_3)^2 (a_4 b_3 - a_3 b_4)^2 \\
&\quad + 8(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2)(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4) \\
&\quad - 32(a_2 b_3 - a_3 b_2)^2 [(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) \\
&\quad + (a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)],
\end{aligned} \tag{2.23}$$

and

$$\begin{aligned}
\mathbf{V} &= 4\mathbf{J}_1\mathbf{J}_2 \\
&= 4 [5(a_2 b_3 - a_3 b_2)^2 + (a_2 b_4 - a_4 b_2)^2 + (a_4 b_3 - a_3 b_4)^2] \\
&\quad \cdot [5(a_2 b_3 - a_3 b_2)^2 + (a_1 b_2 - a_2 b_1)^2 + (a_3 b_1 - a_1 b_3)^2] \\
&= 100(a_2 b_3 - a_3 b_2)^4 \\
&\quad + 20(a_2 b_3 - a_3 b_2)^2 [(a_2 b_4 - a_4 b_2)^2 + (a_4 b_3 - a_3 b_4)^2]
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
& + (a_1 b_2 - a_2 b_1)^2 + (a_3 b_1 - a_1 b_3)^2] \\
& + 4(a_2 b_4 - a_4 b_2)^2 (a_1 b_2 - a_2 b_1)^2 + 4(a_2 b_4 - a_4 b_2)^2 (a_3 b_1 - a_1 b_3)^2 \\
& + 4(a_4 b_3 - a_3 b_4)^2 (a_1 b_2 - a_2 b_1)^2 + 4(a_4 b_3 - a_3 b_4)^2 (a_3 b_1 - a_1 b_3)^2.
\end{aligned}$$

Therefore, from (2.23) and (2.24), we obtain

$$\begin{aligned}
& \mathbf{IV} - \mathbf{V} \\
& = -36(a_2 b_3 - a_3 b_2)^4 \\
& \quad - (a_2 b_3 - a_3 b_2)^2 [20(a_2 b_4 - a_4 b_2)^2 + 20(a_4 b_3 - a_3 b_4)^2 \\
& \quad + 20(a_1 b_2 - a_2 b_1)^2 + 20(a_3 b_1 - a_1 b_3)^2 \\
& \quad + 32(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 32(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)] \\
& \quad - 4[(a_2 b_4 - a_4 b_2)(a_3 b_1 - a_1 b_3) - (a_4 b_3 - a_3 b_4)(a_1 b_2 - a_2 b_1)]^2 \\
& \leq -36(a_2 b_3 - a_3 b_2)^4 \\
& \quad - (a_2 b_3 - a_3 b_2)^2 [20(a_2 b_4 - a_4 b_2)^2 + 20(a_4 b_3 - a_3 b_4)^2 \\
& \quad + 20(a_1 b_2 - a_2 b_1)^2 + 20(a_3 b_1 - a_1 b_3)^2 \\
& \quad + 32(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) + 32(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)]. \tag{2.25}
\end{aligned}$$

*Step 10:* Estimation of  $\Delta$ . By (2.16), (2.22) and (2.25), we get

$$\begin{aligned}
\frac{1}{4}\Delta &= (\mathbf{I}_1 \cdot \mathbf{I}_2 - \mathbf{II} + \mathbf{III}) + (\mathbf{IV} - \mathbf{V}) \\
&\leq - (a_2 b_3 - a_3 b_2)^2 [2(a_2 b_4 - a_4 b_2)^2 + 2(a_4 b_3 - a_3 b_4)^2 \\
&\quad + 2(a_1 b_2 - a_2 b_1)^2 + 2(a_3 b_1 - a_1 b_3)^2 \\
&\quad - 4(a_1 b_2 - a_2 b_1)(a_2 b_4 - a_4 b_2) - 4(a_3 b_1 - a_1 b_3)(a_4 b_3 - a_3 b_4)] \tag{2.26} \\
&= -2(a_2 b_3 - a_3 b_2)^2 \left\{ [(a_1 b_2 - a_2 b_1) - (a_2 b_4 - a_4 b_2)]^2 \right. \\
&\quad \left. + [(a_3 b_1 - a_1 b_3) - (a_4 b_3 - a_3 b_4)]^2 \right\} \\
&\leq 0.
\end{aligned}$$

*Step 11:* From Step 7(1), we know  $p \geq 0$ .

- If  $p = 0$ , then from Step 7(1), we know  $q = r = 0$ . Hence

$$p\lambda_1^2 + q\lambda_1\lambda_2 + r\lambda_2^2 = 0.$$

- If  $p > 0$ , then from (2.26), we know  $\Delta \leq 0$ . Hence

$$p\lambda_1^2 + q\lambda_1\lambda_2 + r\lambda_2^2 \geq 0.$$

By (2.12), it means that

$$\|D\|_F^2 \leq \frac{3}{2} \left( \|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2 \right).$$

The proof is completed.  $\square$

**REMARK 2.1.** Let

$$A = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix},$$

then it is easy to check that

$$\|D\|_F^2 = \frac{3}{2} \left( \|A\|_F^2 \cdot \|BC - CB\|_F^2 + \|B\|_F^2 \cdot \|CA - AC\|_F^2 + \|C\|_F^2 \cdot \|AB - BA\|_F^2 \right) = 18,$$

which shows that the constant  $\frac{3}{2}$  in (1.3) is optimal.

#### REFERENCES

- [1] A. BÖTTCHER AND D. WENZEL, *How big can the commutator of two matrices be and how big is it typically?*, Linear Algebra Appl., 403: 216–228, 2005.
- [2] C.-M. CHENG, X.-Q. JIN AND S.-W. VONG, *A survey on the Böttcher-Wenzel conjecture and related problems*, Oper. Matrices, vol. 9, no. 3, 659–673, 2015.
- [3] L. LÁSZLÓ, *A norm inequality for three matrices*, Electronic Journal of Linear Algebra, vol. 38, 221–226, 2022.

(Received April 18, 2023)

Fen Wang  
Hubei University of Education  
School of Mathematics and Statistics  
430205, Wuhan, P. R. China  
e-mail: wangfen1@hue.edu.cn