

OSCILLATIONS OF EVEN ORDER LINEAR IMPULSIVE DELAY DIFFERENTIAL EQUATIONS

LIJUN PAN AND JINDE CAO

(Communicated by L. Berezansky)

Abstract. In this paper, we devote to investigation of even order impulsive delay differential equations, new oscillation criteria for every solution of equations are established.

1. Introduction

In this paper we study the oscillatory behavior of even order linear impulsive delay differential equations of the form

$$\begin{cases} x^{(n)}(t) + p(t)x(t - \tau) = 0, & t \geq t_0, t \neq t_k, \\ x^{(i)}(t_k^+) = a_k^{(i)}x^{(i)}(t_k), & i = 0, 1, \dots, n-1, k = 1, 2, \dots \\ x^{(i)}(t_0^+) = x_0^{(i)}, \end{cases} \quad (1)$$

where $x^{(0)}(t) = x(t)$, $\tau > 0$, n is even, $0 \leq t_0 < t_1 < t_2 < \dots < t_k < \dots$ such that $t_k \rightarrow \infty$ as $k \rightarrow \infty$, $p(t)$ is positive and continuous in $[t_0, +\infty)$ and for $k = 1, 2, \dots, a_k^{(i)}$ are positive numbers. We also adopt the definitions that

$$x^{(i)}(t_k) = \lim_{h \rightarrow -0} \frac{x^{(i-1)}(t_k + h) - x^{(i-1)}(t_k)}{h}$$

and

$$x^{(i)}(t_k^+) = \lim_{h \rightarrow +0} \frac{x^{(i-1)}(t_k + h) - x^{(i-1)}(t_k^+)}{h}.$$

Let $\varphi(t) : [t_0 - \tau, t_0] \rightarrow R$. Both φ and φ' have at most finite number of the first class discontinuous points on $[t_0 - \tau, t_0]$. By a solution $x = x(t)$, we mean a real function on $[t_0 - \tau, \infty)$ such that $x^{(i)}(t_0^+) = x_0^{(i)}$ for $i = 0, 1, \dots, n-1$ and for $t \in [t_0 - \tau, t_0]$, $x(t) = \varphi(t)$, and $x(t)$ satisfies $x^{(n)}(t) + p(t)x(t - \tau) = 0$ at each point $t \in [t_0 - \tau, \infty)$ with the possible exception of the points $t \neq t_k, t_k + \tau$ and $x^{(i)}(t_k^+) =$

Mathematics subject classification (2010): 34K06, 34K11.

Keywords and phrases: even order, linear, impulsive delay differential equations, oscillation.

This work is supported by the Natural Science Foundation of Guangdong Province of P. R. China (No.9151008002000012).

$a_k^{(i)}x^{(i)}(t_k)$ for any t_k . A solution of (1) is said to be nonoscillatory if this solution is eventually positive or eventually negative. Otherwise, it is said to be oscillatory.

Impulsive differential equations are mathematical apparatus for simulation of process and phenomena observed in control theory, physics, chemistry, population dynamics, biotechnologies, industrial robotics, economics, etc. So there have been quite a few results on properties of their solutions in recent years [8]-[13]. For example, in [5], K. Gopalsamy and B. G. Zhang investigated oscillation of first order delay differential equations with impulses. [14] generalized the results of [5]. In [13] J. R. Yan established Oscillation criteria for nonlinear several delays impulsive differential equations. The oscillatory property of second order ordinary differential equations with impulses had been studied in [4]-[9]. In [3], L. Berezansky and E. Braverman obtained explicit conditions of oscillation and nonoscillation for sufficiently general class of second order impulsive linear delay differential equation. These results were base on the corresponding equations without impulses in [2]. In [10], even order nonlinear differential equations with impulses were studied. But papers devoted to study of the oscillations of even order impulsive delay differential equations are quite rare.

In this paper, appling some known Lemmas and some new Lemmas, we investigate the oscillatory property of (1). We will also provide examples to show that although even order delay differential equations without impulses may have nonoscillatory solutions, adding impulses may lead to oscillatory solutions. That is, impulses may change the oscillatory behavior of an equation. Finally, we also point out that our results may be generalized to even order several delays differential equations with impulses

$$\begin{cases} x^{(n)}(t) + \sum_{j=1}^m p_j(t)x(t - \tau_j) = 0, & t \geq t_0, t \neq t_k, \\ x^{(i)}(t_k^+) = a_k^{(i)}x^{(i)}(t_k), & i = 0, 1, \dots, n-1, k = 1, 2, \dots \end{cases}$$

where n is even, $\tau_j > 0$.

For background material on oscillation of high order differential equations without impulses, we may see the references [7]-[12].

2. Main results

We will establish oscillatory results based on combinations of the following conditions:

- (i) $\lim_{t \rightarrow +\infty} \int_{t_0}^t \prod_{t_0 < t_k < s} \frac{a_k^{(i)}}{a_k^{(i-1)}} ds = +\infty, i = 1, 2, \dots, n-1$;
- (ii) there exists a positive integer k_0 such that for $k \geq k_0$ and for natural number any $l \in \{1, 3, \dots, n-1\}$

$$a_k^{(j)} \geq c_k^{(l)} (j = 0, 1, \dots, l-1),$$

where $c_k^{(l)} = \max\{a_k^{(l)}, a_k^{(l+1)}, \dots, a_k^{(n-1)}\}$;

- (iii) $\limsup_{t \rightarrow +\infty} \prod_{t-2\tau < t_k < t} a_k < +\infty$, where $a_k = \max\{a_k^{(n-1)}, 1\}$;

- (iv) $c = \liminf_{t \rightarrow +\infty} \int_{t-\tau}^t (s-\tau)^{n-1} p(s) ds > 0$.

The main results of the paper are as follows.

THEOREM 1. *Assume that the conditions (i), (ii) hold. If*

$$\limsup_{t \rightarrow +\infty} \int_t^{t+\tau} \prod_{s-\tau \leq t_k < s} \frac{1}{(n-1)} (s-\tau)^{n-1} p(s) ds > (n-1)!, \tag{2}$$

then every solution of (1) is oscillatory.

THEOREM 2. *Assume that the conditions (i), (ii), (iii) and (iv) hold. If*

$$\liminf_{t \rightarrow +\infty} \int_{t-\tau}^t \prod_{s-\tau \leq t_k < s} \frac{1}{(n-1)} (s-\tau)^{n-1} p(s) ds > \frac{(n-1)!}{e}, \tag{3}$$

then every solution of (1) is oscillatory.

To prove Theorem 1 and 2, we need the following Lemmas.

LEMMA 1. (Lakshmikantham et al. [8]) *Assume that:*

(A₀) $m \in PC^1(R_+, R)$ and $m(t)$ is left-continuous at $t_k, k = 1, 2, \dots,$

(A₁) $k = 1, 2, \dots, t \geq t_0,$

$$m'(t) \leq p(t)m(t) + q(t), t \neq t_k, \tag{4}$$

$$m(t_k^+) \leq d_k m(t_k) + b_k, \tag{5}$$

where $p, q \in PC^1(R_+, R), d_k \geq 0$ and b_k are real constants. Then for $t \geq t_0$

$$\begin{aligned} m(t) \leq m(t_0) \prod_{t_0 \leq t_k < t} d_k \exp\left(\int_{t_0}^t p(s) ds\right) + \sum_{t_0 < t_k < t} \left(\prod_{t_0 < t_j < t} d_j \exp\left(\int_{t_k}^t p(s) ds\right)\right) b_k \\ + \int_{t_0}^t \prod_{s < t_k < t} d_k \exp\left(\int_s^t p(\sigma) d\sigma\right) q(s) ds. \end{aligned} \tag{6}$$

REMARK 1. If inequalities (4) and (5) are reversed, then as an conclusion, inequality (6) is also reversed.

LEMMA 2. [10] *Let $x(t)$ be a solution of (1). Suppose there exists some $T \geq t_0$ such that $x(t) > 0$ for $t \geq T$. If condition (i) holds, then there exist $T' \geq T$ and an integer $l \in \{1, 3, \dots, n-1\}$ such that for $t \geq T'$:*

$$x^{(i)}(t) > 0, \quad i = 0, 1, \dots, l,$$

$$(-1)^{i-l} x^{(i)}(t) > 0, \quad i = l+1, l+2, \dots, n-1.$$

LEMMA 3. *Let $x(t)$ be a solution of (1). Suppose there exists some $T \geq t_0$ such that $x(t) > 0$ for $t \geq T$. If conditions (i), (ii) hold, then for any $0 < \theta < 1$, there exists some $T' \geq T$ such that for $t \geq T', x(t) \geq \frac{\theta}{(n-1)!} t^{n-1} x^{(n-1)}(t)$.*

Proof. Without loss of generality, we may assume that $k_0 = 1$. Since $x(t) > 0 (t \geq T)$, by Lemma 2 there exists some $T_1 \geq T$ such that for $t \geq T_1$,

$$x'(t) > 0 \cdots x^{(l)}(t) > 0, (-1)^{i-l} x^{(i)}(t) > 0, i = l + 1, \dots, n - 1.$$

In order to finish our proof, we consider two cases as follows.

Case 1: $l = 1$. Let $u(t) = -\frac{x^{(n-2)}(t)}{x^{(n-1)}(t)}$. Then $u(t) > 0, t \geq T_1$,

$$u'(t) = \frac{-(x^{(n-1)}(t))^2 + x^{(n-2)}(t)x^{(n)}(t)}{(x^{(n-1)}(t))^2} = \frac{x^{(n)}(t)}{x^{(n-1)}(t)}u(t) - 1 < -1 \tag{7}$$

and $u(t_k^+) = \frac{a_k^{(n-2)}}{a_k^{(n-1)}}u(t_k), t_k \geq T_1$. By Lemma 1, we get

$$u(s) < u(t) \prod_{t \leq t_k < s} \frac{a_k^{(n-2)}}{a_k^{(n-1)}} - \int_t^s \prod_{u \leq t_k < s} \frac{a_k^{(n-2)}}{a_k^{(n-1)}} du, s > t \geq T_1. \tag{8}$$

Since $u(s) > 0$, we have

$$u(t) > \int_t^s \prod_{t \leq t_k < u} \frac{a_k^{(n-1)}}{a_k^{(n-2)}} du. \tag{9}$$

So

$$x^{(n-2)}(t) < -x^{(n-1)}(t) \int_t^s \prod_{t \leq t_k < u} \frac{a_k^{(n-1)}}{a_k^{(n-2)}} du. \tag{10}$$

It follows from (1) that

$$x^{(n)}(t) = -p(t)x(t - \tau) < 0, t \geq T_1 + \tau, \\ x^{(n-1)}(t_k^+) = a_k^{(n-1)}x^{(n-1)}(t_k), t_k \geq T_1 + \tau.$$

Set $v(t) = x^{(n-1)}(t)$. Then

$$v'(t) < 0, t \geq T_1 + \tau, v(t_k^+) = a_k^{(n-1)}v(t_k), t_k \geq T_1 + \tau.$$

Applying Lemma 1, we obtain

$$v(u) < v(t) \prod_{t \leq t_k < u} a_k^{(n-1)}, u > t \geq T_1 + \tau.$$

That is

$$x^{(n-1)}(u) < x^{(n-1)}(t) \prod_{t \leq t_k < u} a_k^{(n-1)}, u > t \geq T_1 + \tau. \tag{11}$$

From (10) and (11), we have

$$x^{(n-2)}(t) < - \int_t^s \prod_{t \leq t_k < u} \frac{1}{a_k^{(n-2)}} x^{(n-1)}(u) du. \tag{12}$$

Let $m(t) = x^{(n-3)}(t)$. Then:

$$m(t_k^+) = x^{(n-3)}(t_k^+) = a_k^{(n-3)}x^{(n-3)}(t_k) = a_k^{(n-3)}m(t_k),$$

$$m'(t) < - \int_t^s \prod_{t \leq t_k < u} \frac{1}{a_k^{(n-2)}} x^{(n-1)}(u) du.$$

By Lemma 1, we have

$$m(s) < m(t) \prod_{t \leq t_k < s} a_k^{(n-3)} - \int_t^s \prod_{v \leq t_k < s} a_k^{(n-3)} \left(\int_v^s \prod_{v \leq t_k < u} \frac{1}{a_k^{(n-2)}} x^{(n-1)}(u) du \right) dv. \tag{13}$$

From condition (ii) and $m(s) > 0$, we get

$$x^{(n-3)}(t) > \int_t^s \prod_{t \leq t_k < u} \frac{1}{c_k^{(n-3)}} (u-t)x^{(n-1)}(u) du. \tag{14}$$

By repeating the above argument repeatedly, we may show that

$$x'(t) \geq \int_t^s \prod_{t \leq t_k < u} \frac{1}{c_k^{(1)}} \frac{(u-t)^{n-3}}{(n-3)!} x^{(n-1)}(u) du, \quad s > t \geq T_1 + \tau. \tag{15}$$

Since $x(t_k^+) = a_k^{(0)}x(t_k)$, $t_k \geq T_1 + \tau$, by Lemma 1 and noting that

$$x^{(n-1)}(t) < \prod_{u \leq t_k < s} a_k^{(n-1)} x^{(n-1)}(u), \quad t > u,$$

and condition (ii), we have

$$\begin{aligned} x(t) &> x(T_1^+) \prod_{T_1 < t_k < t} a_k^{(0)} + \int_{T_1}^t \prod_{v \leq t_k < t} a_k^{(0)} \left(\int_v^s \prod_{v \leq t_k < u} \frac{1}{c_k^{(1)}} \frac{(u-v)^{n-3}}{(n-3)!} x^{(n-1)}(u) du \right) dv \\ &> \int_{T_1}^t \prod_{v \leq t_k < t} a_k^{(0)} \left(\int_v^s \prod_{v \leq t_k < u} \frac{1}{c_k^{(1)}} \frac{(u-v)^{n-3}}{(n-3)!} x^{(n-1)}(u) du \right) dv \\ &> \int_{T_1}^t \prod_{v \leq t_k < v} a_k^{(0)} \left(\int_v^t \prod_{v \leq t_k < u} \frac{1}{c_k^{(1)}} \frac{(u-v)^{n-3}}{(n-3)!} x^{(n-1)}(u) du \right) dv \\ &\geq x^{(n-1)}(t) \int_{T_1}^t \prod_{v \leq t_k < t} a_k^{(0)} \left(\int_v^t \prod_{v \leq t_k < u} \frac{1}{c_k^{(1)}} \prod_{u \leq t_k < t} \frac{1}{a_k^{(n-1)}} \frac{(u-v)^{n-3}}{(n-3)!} du \right) dv \\ &\geq x^{(n-1)}(t) \int_{T_1}^t \left(\int_v^t \frac{(u-v)^{n-3}}{(n-3)!} du \right) dv \geq x^{(n-1)}(t) \frac{(t-T_1)^{n-1}}{(n-1)!}. \end{aligned} \tag{16}$$

Thus, for any $0 < \theta < 1$, there exists some $T' \geq T_1 + \tau$ such that for $t \geq T'$,

$$x(t) \geq \frac{\theta}{(n-1)!} t^{n-1} x^{(n-1)}(t).$$

Case 2: $2 < l \leq n - 1$. From the proof of Lemma 3, we have:

$$x^{(l-1)}(t) > x^{(n-1)}(t) \frac{(t - T_1)^{n-l}}{(n-l)!}, \quad t \geq T_1 + \tau,$$

$$x^{(l-2)}(t_k^+) = a_k^{(l-2)} x^{(l-2)}(t_k), \quad t_k \geq T_1 + \tau.$$

Let $w(t) = x^{(l-2)}(t)$. Then

$$w'(t) > x^{(n-1)}(t) \frac{(t - T_1)^{n-l}}{(n-l)!} \quad \text{and} \quad w(t_k^+) = a_k^{(l-2)} w(t_k), \quad t_k \geq T_1 + \tau. \quad (17)$$

By Lemma 1 and noting that

$$x^{(n-1)}(t) < \prod_{u \leq t_k < t} a_k^{(n-1)} x^{(n-1)}(u), \quad t > u \geq T_1 + \tau$$

and condition (ii), we get

$$\begin{aligned} w(t) &> w(T_1^+) \prod_{T_1 < t_k < t} a_k^{(l-2)} + \int_{T_1}^t \prod_{u \leq t_k < t} a_k^{(l-2)} x^{(n-1)}(u) \frac{(u - T_1)^{n-l}}{(n-l)!} du \\ &> \int_{T_1}^t \prod_{u \leq t_k < t} a_k^{(l-2)} x^{(n-1)}(u) \frac{(u - T_1)^{n-l}}{(n-l)!} du \\ &> x^{(n-1)}(t) \int_{T_1}^t \prod_{u \leq t_k < t} \frac{a_k^{(l-2)}}{a_k^{(n-1)}} \frac{(u - T_1)^{n-l}}{(n-l)!} du \\ &> x^{(n-1)}(t) \int_{T_1}^t \frac{(u - T_1)^{n-l}}{(n-l)!} du = x^{(n-1)}(t) \frac{(t - T_1)^{n-l+1}}{(n-l+1)!}. \end{aligned} \quad (18)$$

That is

$$x^{(l-2)}(t) > x^{(n-1)}(t) \frac{(t - T_1)^{n-l+1}}{(n-l+1)!}. \quad (19)$$

By repeating the above argument, we have

$$x(t) > x^{(n-1)}(t) \frac{(t - T_1)^{n-1}}{(n-1)!}. \quad (20)$$

Thus, for any $0 < \theta < 1$, there exists some $T' \geq T_1 + \tau \geq T$ such that for $t \geq T'$, $x(t) \geq \frac{\theta}{(n-1)!} t^{n-1} x^{(n-1)}(t)$.

LEMMA 4. *Let $x(t)$ be an eventually positive solution of (1). If conditions (i), (ii), (iii) and (iv) hold, then for any sufficiently large t , there exists a $t^* \in [t - \tau, t]$ such that*

$$\frac{x^{(n-1)}(t^* - \tau)}{x^{(n-1)}(t^*)} < +\infty.$$

Proof. Without loss of generality, we may assume that $k_0 = 1$. Since $x(t)$ is a eventually positive solution of (1). We may assume that $x(t) > 0 (t \geq T_1 \geq t_0)$. By Lemma 2, there exist a $T_2 \geq T_1$ and an integer $l \in \{1, 2, \dots, n - 1\}$ such that for $t \geq T_2$,

$$x'(t) > 0 \cdots x^{(l)}(t) > 0, \quad (-1)^{(i-l)}x^{(i)}(t) > 0, \quad i = l + 1, \dots, n - 1.$$

From Lemma 3, there exists $\theta, 0 < \theta < 1$ and $T_3 \geq T_1$ such that for $t \geq T_3$,

$$x(t) \geq \frac{\theta}{(n - 1)!}t^{n-1}x^{(n-1)}(t).$$

Since condition (iv) holds, so choosing $0 < L \leq c$, then exists T' such that for $t \geq T'$

$$\int_{t-\tau}^t (s - \tau)^{n-1}p(s)ds > L. \tag{21}$$

Fixing a $t \geq T'$, then there exists some $t^* \in [t - \tau, t]$ such that

$$\int_{t-\tau}^{t^*} (s - \tau)^{n-1}p(s)ds > \frac{L}{2} \quad \text{and} \quad \int_{t^*}^t (s - \tau)^{n-1}p(s)ds > \frac{L}{2}. \tag{22}$$

We suppose that T is sufficiently large and (22) holds for $t \geq T$ and

$$x^{(n-1)}(t - \tau) > 0 \quad x(t - \tau) \geq \frac{\theta}{(n - 1)!}(t - \tau)^{n-1}x^{(n-1)}(t - \tau). \tag{23}$$

In view of (1) and (23), we have

$$x^{(n)}(t) + \frac{\theta}{(n - 1)!}(t - \tau)^{n-1}p(t)x^{(n-1)}(t - \tau) \leq 0, t \geq T \tag{24}$$

Integrating the both sides of (24) on $[t - \tau, t^*]$, we have

$$x^{(n-1)}(t^*) - x^{(n-1)}((t - \tau)^+) - \sum_{t-\tau < t_k < t^*} (a_k^{(n-1)} - 1)x^{(n-1)}(t_k) + \frac{\theta}{(n - 1)!} \int_{t-\tau}^{t^*} (s - \tau)^{n-1}p(s)x^{(n-1)}(s - \tau)ds \leq 0. \tag{25}$$

From (25), it follows that

$$x^{(n-1)}((t - \tau)^+) + \sum_{t-\tau < t_k < t^*} (a_k^{(n-1)} - 1)x^{(n-1)}(t_k) \geq \frac{\theta}{(n - 1)!} \int_{t-\tau}^{t^*} (s - \tau)^{n-1}p(s)x^{(n-1)}(s - \tau)ds. \tag{26}$$

Since

$$x^{(n)}(t) < 0, \quad t \geq T \quad \text{and} \quad x^{(n-1)}(t_k^+) = a_k^{(n-1)}x^{(n-1)}(t_k), \quad t_k > T,$$

by Lemma 1, it is easy to see that

$$x^{(n-1)}(s-\tau) > \prod_{s-\tau \leq t_k < t^* - \tau} \frac{1}{a_k^{(n-1)}} x^{(n-1)}(t^* - \tau), t - \tau \leq s \leq t^*. \quad (27)$$

We claim

$$\begin{aligned} x^{(n-1)}((t-\tau)^+) + \sum_{t-\tau < t_k < t^*} (a_k^{(n-1)} - 1)x^{(n-1)}(t_k) \\ \leq \prod_{t-\tau < t_k < t^*} a_k x^{(n-1)}((t-\tau)^+). \end{aligned} \quad (28)$$

Indeed, we assume that $t_{p+1} < t_{p+2} < \dots < t_{p+m}$ are m impulsive points on $(t-\tau, t^*)$. Note that for $t_k > t-\tau$,

$$x^{(n-1)}(t_k) < \prod_{t-\tau < t_j < t_k} a_k^{(n-1)} x^{(n-1)}((t-\tau)^+) \leq \prod_{t-\tau < t_j < t_k} a_k^{(n-1)} x^{(n-1)}((t-\tau)^+).$$

Thus

$$\begin{aligned} x^{(n-1)}((t-\tau)^+) + \sum_{t-\tau < t_k < t^*} (a_k^{(n-1)} - 1)x^{(n-1)}(t_k) \\ \leq x^{(n-1)}((t-\tau)^+) + [(a_{p+1}^{(n-1)} - 1)x^{(n-1)}(t_{p+1}) + (a_{p+2}^{(n-1)} - 1)x^{(n-1)}(t_{p+2}) + \dots \\ + (a_{p+m}^{(n-1)} - 1)x^{(n-1)}(t_{p+m})] \\ = x^{(n-1)}((t-\tau)^+) + (a_{p+1}^{(n-1)} x^{(n-1)}(t_{p+1}) + a_{p+2}^{(n-1)} x^{(n-1)}(t_{p+2}) + \dots \\ + a_{p+m}^{(n-1)} x^{(n-1)}(t_{p+m})) - (x^{(n-1)}(t_{p+1}) + x^{(n-1)}(t_{p+2}) + \dots + x^{(n-1)}(t_{p+m})) \\ \leq x^{(n-1)}((t-\tau)^+) + \sum_{t-\tau < t_k < t^*} (a_k - 1) \prod_{t-\tau < t_j < t_k} a_k x^{(n-1)}((t-\tau)^+) \\ = x^{(n-1)}((t-\tau)^+) [1 + \sum_{t-\tau < t_k < t^*} (a_k - 1) \prod_{t-\tau < t_j < t_k} a_k] \\ = x^{(n-1)}((t-\tau)^+) [1 + (a_{p+1} - 1) + (a_{p+2} - 1)a_{p+1} \\ + \dots + (a_{p+m} - 1)a_{p+1}a_{p+2} \dots a_{p+m-1}] \\ = x^{(n-1)}((t-\tau)^+) a_{p+1}a_{p+2} \dots a_{p+m} = \prod_{t-\tau < t_k < t^*} a_k x^{(n-1)}((t-\tau)^+). \end{aligned} \quad (29)$$

From (22), (25), (27) and (28), we have

$$\begin{aligned} \prod_{t-\tau < t_k < t^*} a_k x^{(n-1)}((t-\tau)^+) &\geq \frac{\theta}{(n-1)!} \int_{t-\tau}^{t^*} (s-\tau)^{n-1} p(s) x^{(n-1)}(s-\tau) ds \\ &> \frac{\theta}{(n-1)!} x^{(n-1)}(t^* - \tau) \int_{t-\tau}^{t^*} \prod_{s-\tau \leq t_k < t^* - \tau} \frac{1}{a_k^{(n-1)}} (s-\tau)^{n-1} p(s) ds \\ &\geq \frac{L\theta}{2(n-1)!} \prod_{t-2\tau < t_k < t^* - \tau} \frac{1}{a_k} x^{(n-1)}(t^* - \tau), t - \tau < s < t^*. \end{aligned} \quad (30)$$

So

$$x^{(n-1)}(t^* - \tau) < \frac{2(n-1)!}{L\theta} \prod_{t-\tau < t_k < t^*} a_k \prod_{t-2\tau < t_k < t^* - \tau} a_k x^{(n-1)}((t - \tau)^+). \tag{31}$$

Similar to (31), we get

$$x^{(n-1)}((t - \tau)^+) < \frac{2(n-1)!}{L\theta} \prod_{t^* \leq t_k < t} a_k \prod_{t^* - \tau \leq t_k < t - \tau} a_k x^{(n-1)}(t^*). \tag{32}$$

So

$$x^{(n-1)}(t^* - \tau) < \left(\frac{2(n-1)!}{L\theta}\right)^2 \prod_{t-\tau < t_k < t} a_k \prod_{t-2\tau < t_k < t - \tau} a_k x^{(n-1)}(t^*) \tag{33}$$

Thus, from condition (iii), we have $\frac{x^{(n-1)}(t^* - \tau)}{x^{(n-1)}(t^*)} < +\infty$.

REMARK 2. $x(t)$ is a eventually negative solution of (1), we also have conclusions similar to Lemmas 2-4.

LEMMA 5. [1] *Definite a sequence $\{h^n(1)\}_{n=1}^\infty$ satisfying*

$$h^1(1) = h(1), h^2(1) = e^{\theta L h^1(1)}, \dots, h^n(1) = e^{\theta L h^{n-1}(1)},$$

where $h(s) = e^{\theta L s}$. If $\theta L > \frac{1}{e}$, then sequence $\{h^n(1)\}_{n=1}^\infty$ is monotonically increasing and tends to $+\infty$.

We now turn to the proof of theorems 1 and 2.

Proof of Theorem 1. Without loss of generality, we may assume that $k_0 = 1$. If (1) has a nonoscillation $x(t)$, we may assume that $x(t) > 0 (t \geq T \geq t_0)$. By Lemma 2, there exist $T_1 \geq T$ and an integer $l \in \{1, 3, \dots, n-1\}$ such that for $t \geq T_1$,

$$x'(t) > 0 \dots x^{(l)}(t) > 0, (-1)^{(i-1)} x^{(i)}(t) > 0, i = l + 1, \dots, n - 1.$$

Let

$$c_1 = \limsup_{t \rightarrow +\infty} \frac{1}{(n-1)!} \int_t^{t+\tau} \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s - \tau)^{n-1} p(s) ds.$$

From (2) we have $c_1 > 1$. Taking $L, 1 < L \leq c_1$, by Lemma 3 there exist a constant $\theta, 2/(L+1) < \theta < 1$ and $t^* \geq T_1$ such that for $t \geq t^* + \tau$,

$$x(t - \tau) \geq \frac{\theta}{(n-1)!} (t - \tau)^{n-1} x^{(n-1)}(t - \tau). \tag{34}$$

From (1) and (34), we have

$$x^{(n)}(t) + \frac{\theta}{(n-1)!} (t - \tau)^{n-1} p(t) x^{(n-1)}(t - \tau) \leq 0, t \geq t^* + \tau, t \neq t_k, t_k \geq t^* + \tau. \tag{35}$$

Set $v(t) = x^{(n-1)}(t)$. From above formula, it follows that

$$\begin{cases} v'(t) \leq -\frac{\theta}{(n-1)!}(t-\tau)^{n-1}p(t)x^{(n-1)}(t-\tau), & t \geq t^* + \tau, t \neq t_k, \\ v(t_k^+) = a_k^{(n-1)}v(t_k), & t_k \geq t^* + \tau. \end{cases} \quad (36)$$

Applying Lemma 1, we obtain

$$\begin{aligned} v(t+\tau) &\leq v(t^+) \prod_{t < t_k < t+\tau} a_k^{(n-1)} \\ &\quad - \frac{\theta}{(n-1)!} \int_t^{t+\tau} \prod_{s \leq t_k < t+\tau} a_k^{(n-1)} (s-\tau)^{n-1} p(s) x^{(n-1)}(s-\tau) ds. \end{aligned}$$

That is

$$\begin{aligned} x^{(n-1)}(t+\tau) &\leq x^{(n-1)}(t^+) \prod_{t < t_k < t+\tau} a_k^{(n-1)} \\ &\quad - \frac{\theta}{(n-1)!} \int_t^{t+\tau} \prod_{s \leq t_k < t+\tau} a_k^{(n-1)} (s-\tau)^{n-1} p(s) x^{(n-1)}(s-\tau) ds. \end{aligned} \quad (37)$$

Since

$$x^{(n)}(t) < 0, t \geq t^* + \tau \text{ and } x^{(n-1)}(t_k^+) = a_k^{(n-1)} x^{(n-1)}(t_k), t_k > t^* + \tau,$$

by Lemma 1, we have:

$$x^{(n-1)}(s-\tau) \geq x^{(n-1)}(t^+) \prod_{s-\tau \leq t_k \leq t} a_k^{(n-1)}. \quad (38)$$

From the inequalities (37) and (38), it follows that

$$\begin{aligned} x^{(n-1)}(t+\tau) &\leq x^{(n-1)}(t^+) \prod_{t < t_k < t+\tau} a_k^{(n-1)} \left[1 - \right. \\ &\quad \left. \frac{\theta}{(n-1)!} \int_t^{t+\tau} \prod_{s-\tau \leq t_k < s} a_k^{(n-1)} \frac{1}{a_k^{(n-1)}} (s-\tau)^{n-1} p(s) ds \right], t \geq t^* + \tau. \end{aligned} \quad (39)$$

In view of (2). Then for $\varepsilon = \frac{L-1}{2} > 0$, there exists $T_\lambda > t^* + \tau$ such that

$$\frac{1}{(n-1)!} \int_{T_\lambda}^{T_\lambda+\tau} \prod_{s-\tau \leq t_k < s} a_k^{(n-1)} (s-\tau)^{n-1} p(s) ds > L - \varepsilon = \frac{L+1}{2}. \quad (40)$$

From (39), (40) and $\frac{2}{L+1} < \theta < 1$, we have

$$x^{(n-1)}(T_\lambda + \tau) \leq x^{(n-1)}(T_\lambda^+) \left[1 - \right.$$

$$\frac{\theta}{(n-1)!} \int_{T_\lambda}^{T_\lambda+\tau} \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s-\tau)^{n-1} p(s) ds \Big] < 0. \tag{41}$$

This is a contrary to the fact that $x^{(n-1)}(t) > 0, t \geq T_1$. Thus every solution of (1) is oscillatory.

Proof of Theorem 2. Without loss of generality, we may assume that $k_0 = 1$. If (1) has a nonoscillation $x(t)$, we may assume that $x(t) > 0(t \geq T \geq t_0)$. By Lemma 2, there exist $T_1 \geq T$ and an integer $l \in \{1, 3, \dots, n-1\}$ such that for $t \geq T_1$,

$$x'(t) > 0 \dots x^{(l)}(t) > 0, (-1)^{(i-1)} x^{(i)}(t) > 0, i = l+1, \dots, n-1.$$

Let

$$c_2 = \liminf_{t \rightarrow +\infty} \frac{1}{(n-1)!} \int_{t-\tau}^t \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s-\tau)^{n-1} p(s) ds,$$

(3) yields $c_2 > 1/e$. Taking L such that $1/e < L \leq c$, by Lemma 3 there exist a constant $\theta, 2/(Le+1) < \theta < 1$ and $t^* \geq T_1$ such that for $t \geq t^* + \tau$,

$$x(t-\tau) \geq \frac{\theta}{(n-1)!} (t-\tau)^{n-1} x^{(n-1)}(t-\tau). \tag{42}$$

From Lemma 4, for any sufficiently large t , there exists $T_2 \in [t-\tau, t]$ such that

$$\frac{x^{(n-1)}(T_2-\tau)}{x^{(n-1)}(T_2)} < +\infty \tag{43}$$

From (3), it follows that for sufficiently large t

$$\frac{1}{(n-1)!} \int_{t-\tau}^t \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s-\tau)^{n-1} p(s) ds > L. \tag{44}$$

We suppose that (42), (43) and (44) hold for $t \geq t^* + \tau$. On the other hand, noting that

$$x^{(n-1)}(t-\tau) \geq \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} x^{(n-1)}(t). \tag{45}$$

From (1), (42) and (45), we have

$$x^{(n)}(t) + \frac{\theta}{(n-1)!} \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} (t-\tau)^{n-1} p(t) x^{(n-1)}(t) \leq 0, t \geq t^* + \tau. \tag{46}$$

Set $v(t) = x^{(n-1)}(t)$, then $v(t_k^+) = a_k^{(n-1)} v(t_k)$ and

$$v'(t) \leq -\frac{\theta}{(n-1)!} \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} (t-\tau)^{n-1} p(t) v(t), t \geq t^* + \tau, t \neq t_k. \tag{47}$$

Applying Lemma 1, we obtain

$$x^{(n-1)}(t) \leq x^{(n-1)}(t - \tau) \prod_{t-\tau \leq t_k < t} a_k^{(n-1)} \exp \left[-\frac{\theta}{(n-1)!} \int_{t-\tau}^t \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s - \tau)^{n-1} p(s) ds \right]. \quad (48)$$

That is

$$\begin{aligned} \frac{x^{(n-1)}(t - \tau)}{x^{(n-1)}(t)} &\geq \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} \exp \left[\frac{\theta}{(n-1)!} \int_{t-\tau}^t \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s - \tau)^{n-1} p(s) ds \right] \\ &\geq \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} e^{\theta L} = \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} h^1(1) \end{aligned} \quad (49)$$

where $h(s) = e^{\theta Ls}$, $\theta L > 1/e$. Thus, from (1), (42), (45) and (49), we have

$$x^{(n)}(t) + \frac{\theta}{(n-1)!} h^1(1) \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} (t - \tau)^{n-1} p(t) x^{(n-1)}(t) \leq 0, \quad t \geq t^* + \tau. \quad (50)$$

Applying the above method, by Lemma 1 we have

$$\frac{x^{(n-1)}(t - \tau)}{x^{(n-1)}(t)} \geq \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} e^{\theta L h^1(1)} = \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} h^2(1), \quad t \geq t^* + 2\tau. \quad (51)$$

By repeating the above method repeatedly, we get

$$\frac{x^{(n-1)}(t - \tau)}{x^{(n-1)}(t)} \geq \prod_{t-\tau \leq t_k < t} \frac{1}{a_k^{(n-1)}} h^n(1), \quad t \geq t^* + n\tau, \quad n = 1, 2, \dots \quad (52)$$

From Lemma 5, we have $h^n(1) \rightarrow \infty$. That is $\frac{x^{(n-1)}(t - \tau)}{x^{(n-1)}(t)} \rightarrow \infty$. This is a contrary to (43). Thus every solution of (1) is oscillatory.

REMARK 3. It is easy to see that the above theorems 1 and 2 can be generalized to the equations of the form

$$\begin{cases} x^{(n)}(t) + \sum_{j=1}^m p_j(t)x(t - \tau_j) = 0, \quad t \geq t_0, \quad t \neq t_k, \\ x^{(i)}(t_k^+) = a_k^{(i)} x^{(i)}(t_k), \quad i = 0, 1, \dots, n-1, \quad k = 1, 2, \dots \end{cases}$$

where n is even, $\tau_j > 0$. Indeed, if we replace the conditions (iii), (iv), (2) and (3) with (iii'), (iv'), (2') and (3') as follows:

(iii') $\limsup_{t \rightarrow +\infty} \prod_{t-2\tau < t_k < t} a_k < +\infty$, where $a_k = \max\{a_k^{(n-1)}, 1\}$, $\tau = \max\{\tau_1, \dots, \tau_m\}$,

$$(iv') \quad c = \liminf_{t \rightarrow +\infty} \int_{t-\tau}^t \sum_{j=1}^m (s - \tau_j)^{n-1} p_j(s) ds > 0, \text{ where } \tau' = \min\{\tau_1, \dots, \tau_m\}$$

(2')

$$\limsup_{t \rightarrow +\infty} \int_t^{t+\tau} \prod_{j=1}^m \frac{1}{a_k^{(n-1)}} (s - \tau_j)^{n-1} p_j(s) ds > (n-1)!,$$

(3')

$$\liminf_{t \rightarrow +\infty} \int_{t-\tau}^t \sum_{j=1}^m \prod_{s-\tau_j \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s - \tau')^{n-1} p_j(s) ds > \frac{(n-1)!}{e}.$$

Then, we will obtain the similar Theorems of the above equations.

EXAMPLE 1. Consider the Equation

$$\begin{cases} x^{(n)}(t) + \frac{1 \times 3 \times \dots \times (2n-3)}{2^n} t^{-n+\frac{1}{2}} (t - \frac{1}{3})^{-\frac{1}{2}} x(t - \frac{1}{3}) = 0, & t \geq \frac{1}{2}, t \neq k, \\ x^{(i)}(k^+) = \frac{k^{i-2}}{(k+1)^i} x^{(i)}(k), & i = 0, 1, 2, \dots, n-1, k = 1, 2, \dots \end{cases} \quad (53)$$

where n is even, $t_k = k$,

$$a_k^{(i)} = \frac{k^{i-2}}{(k+1)^i}, \quad p(t) = \frac{1 \times 3 \times \dots \times (2n-3)}{2^n} t^{-n+\frac{1}{2}} (t - \frac{1}{3})^{-\frac{1}{2}} \text{ and } \tau = \frac{1}{3}.$$

It is easy to see that conditions (i) and (ii) satisfy:

$$\begin{aligned} \limsup_{t \rightarrow +\infty} \int_t^{t+\tau} \prod_{s-\tau \leq t_k < s} \frac{1}{a_k^{(n-1)}} (s - \tau)^{n-1} p(s) ds \\ = \lim_{k \rightarrow +\infty} \sup \frac{1}{a_k^{(n-1)}} \int_{t_k}^{t_k+\tau} (s - \tau)^{n-1} p(s) ds = +\infty > (n-1)!. \end{aligned}$$

By Theorem 1, every solution of (53) is oscillatory. But the delay differential equation

$$x^{(n)}(t) + \frac{1 \times 3 \times \dots \times (2n-3)}{2^n} t^{-n+\frac{1}{2}} (t - \frac{1}{3})^{-\frac{1}{2}} x(t - \frac{1}{3}) = 0, \quad t \geq \frac{1}{2}$$

has a nonnegative solution $x = \sqrt{t}$. This example shows that impulses play an important role in the oscillatory behavior of equations under perturbing impulses.

REFERENCES

[1] T. H. BARR, *Oscillations in linear delay differential equations*, J. Math. Anal. Appl., **195** (1995), 261–277.
 [2] L. BEREZANSKY AND E. BRAVERMAN, *Some oscillation problems for a second order linear delay differential equation*, J. Math. Anal. Appl., **220** (1998), 719–740.
 [3] L. BEREZANSKY AND E. BRAVERMAN, *On oscillation of a second order impulsive linear delay differential equation*, J. Math. Anal. Appl., **233** (1999), 276–300.
 [4] Y. S. CHEN, W. Z. FENG, *Oscillations of second order nonlinear ODE with impulses*, J. Math. Anal. Appl., **210** (1997), 150–169.

- [5] K. GOPALSAMY AND B. G. ZHANG, *On delay differential equations with impulses*, J. Math. Anal. Appl., **139** (1989), 110–122.
- [6] Z. M. HE AND W. G. GE, *Oscillations of second-order nonlinear impulsive ordinary differential equations*, Comput. Appl. Math., **158** (2003), 397–406.
- [7] I. T. KIGURADZE, *The capability of certain solutions of ordinary differential equations to oscillate*, Dokl. Akad. Nauk. SSSR, **144** (1962), 33–36.
- [8] V. LAKSHMIKANTHAM, D. D. BAINOV, P. S. SIMEONOV, *Theory of impulsive differential equations*, World Scientific, Singapore, 1989.
- [9] J. W. LUO AND L. DEBNATH, *Oscillations of second order nonlinear ordinary differential equations with impulses*, J. Math. Anal. Appl., **240** (1999), 105–114.
- [10] L. J. PAN, G. Q. WANG AND S. S. CHEN, *Oscillation of even order nonlinear differential equations with impulses*, Funkcialaj Ekvacioj, **50** (2007), 117–131.
- [11] G. H. RYDER AND V. V. WEND, *Oscillation of solutions certain ordinary differential equations of n -th order*, Proc. Amer. Math. Soc., **21** (1970), 463–469.
- [12] W. TRENCH, *An oscillation condition for differential equations of arbitrary order*, Proc. Amer. Math. Soc., **82** (1981), 548–552.
- [13] J. R. YAN, *Oscillation of nonlinear delay impulsive differential equations and inequalities*, J. Math. Anal. Appl., **265** (2002), 332–342.
- [14] Y. Z. ZHANG, A. M. ZHAO, J. R. YAN, *Oscillation criteria for impulsive delay differential equation*, J. Math. Anal. Appl., **205** (1997), 461–470.

(Received July 1, 2009)

(Revised December 21, 2009)

Lijun Pan and JinDe Cao
Department of Mathematics
Southeast University
Nanjing 210096
P. R. China

Lijun Pan
Department of Mathematics
Jia Ying University
Meizhou Guangdong 514015
P. R. China
e-mail: plj1977@126.com