

ORLICZ–QUASI–CAUCHY DOUBLE SEQUENCE SPACES FOR RH –REGULAR MATRIX

KULDIP RAJ, AYHAN ESI AND SEEMA JAMWAL

Abstract. In this research paper we pioneer, a new double sequence space called Orlicz-Quasi-Cauchy double sequence space, $(M, A, \Delta)_{OQC}$. We investigate continuity type properties of Orlicz double function defined on a double subset $A \times A$ of R^2 into R and study some vital results related to uniform continuity. In the last we also make an attempt to prove some topological properties of $(M, A, \Delta)_{OQC}$.

1. Introduction

An Orlicz function M is a function, which is continuous, non-decreasing and convex with $M(0) = 0$, $M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

Lindenstrauss and Tzafriri [13] used the idea of Orlicz function to define the following sequence space. Let w be the space of all real or complex sequences $x = (x_k)$, then

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0 \right\}$$

which is called as an Orlicz sequence space. The space ℓ_M is a Banach space with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}.$$

It is shown in [13] that every Orlicz sequence space ℓ_M contains a subspace isomorphic to ℓ_p ($p \geq 1$). The Δ_2 -condition is equivalent to $M(Lx) \leq kLM(x)$ for all values of $x \geq 0$ and for $L > 1$.

The study of single Quasi-Cauchy sequences have been discussed in ([9], [10], [17]). In ([9], [10]) Çakallı and Çakallı and Hazarika have introduced and studied the statistical Quasi-Cauchy sequences and Ideal Quasi-Cauchy sequences respectively. Also in [17], Hazarika studied ϕ -statistical Quasi-Cauchy sequences in details.

The notion of difference sequence spaces was introduced by Kızmaz [12], who studied the difference sequence spaces $l_{\infty}(\Delta)$, $c(\Delta)$ and $c_0(\Delta)$. For Z a given sequence space, we have

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\}$$

Mathematics subject classification (2010): 40A05, 46E30.

Keywords and phrases: Orlicz function, difference sequence spaces, RH -regular matrix, uniformity continuity, quasi-Cauchy sequences.

for $Z = c, c_0$ and l_∞ where $\Delta x = (\Delta x_k) = (x_k - x_{k+1})$. The difference sequence spaces was also discussed in (Altınok et. al. [3], Isik [18], Tripathy et al. [34]). Similarly, we can define difference operators on double sequence spaces as:

$$\begin{aligned} \Delta x_{k,l} &= (x_{k,l} - x_{k,l+1}) - (x_{k+1,l} - x_{k+1,l+1}) \\ &= x_{k,l} - x_{k,l+1} - x_{k+1,l} + x_{k+1,l+1} \end{aligned}$$

In 1900, Pringsheim [24] introduced the concept of convergence of real double sequences. Four year later, Hardy [15] introduced the notion of regular convergence for double sequences.

The initial work on double sequences is found in Bromwich [5]. Later on Móricz [19], Móricz and Rhoades [20], Tripathy ([32], [33]), Başarır and Sonalcan [4] and many others. Quite recently, Zeltser [35] in her Ph.D thesis has essentially studied both the theory of topological double sequence spaces and the theory of summability of double sequences. Mursaleen and Edely [23] have recently introduced the statistical convergence and Cauchy convergence for double sequences and given the relation between statistical convergent and strongly Cesàro summable double sequences. Nextly, Mursaleen [21] and Mursaleen and Edely [22] have defined the almost strong regularity of matrices for double sequences and applied these matrices to establish a core theorem and introduced the M -core for double sequences and determined those four dimensional matrices transforming every bounded double sequences $x = (x_{jk})$ into one whose core is a subset of the M -core of x . Recently Quasi Cauchy double sequence spaces was studied by Patterson and Cakalli [28]. A considerable number of papers which appeared in recent years study double sequences from various point of view (see [1], [2], [6], [7], [8], [11], [16], [26], [27], [29], [30]).

DEFINITION 1. A double sequence $x = \{x_{k,l}\}$ is Cauchy provided that, given an $\varepsilon > 0$ there exists an $N \in \mathbb{N}$ such that $|x_{k,l} - x_{s,t}| < \varepsilon$ whenever $k, l, s, t > N$.

DEFINITION 2. By the convergence of a double sequence we mean the convergence in the Pringsheim sense i.e. a double sequence $x = (x_{k,l})$ has *Pringsheim limit* L (denoted by $P - \lim x = L$) provided that given $\varepsilon > 0$ there exists $n \in \mathbb{N}$ such that $|x_{k,l} - L| < \varepsilon$ whenever $k, l > n$. We shall write more briefly as P -convergent.

If $\lim x = \infty$, (equivalently, for every $\varepsilon > 0$ there are $n_1, n_2 \in \mathbb{N}$ such that $|x_{m,n}| > M$ whenever $m > n_1, n > n_2$, then $x = \{x_{m,n}\}$ is said to be definitely divergent. A double sequence $x = \{x_{m,n}\}$ is bounded if there is an $M > 0$ such that $|x_{m,n}| < M$ for all $m, n \in \mathbb{N}$. Notice that a P -convergent double sequence need not be bounded.

DEFINITION 3. [25] A double sequence y is a double subsequence of x provided that there exist increasing index sequences $\{n_j\}$ and $\{k_j\}$ such that if $\{x_j\} = \{x_{n_j, k_j}\}$, then y is formed by

x_1	x_2	x_5	x_{10}
x_4	x_3	x_6	—
x_9	x_8	x_7	—
—	—	—	—

DEFINITION 4. Let $A = (a_{mnkl})$ denote a four dimensional summability method that maps the complex double sequences x into the double sequence Ax where the mn^{th} term to Ax is as follows:

$$(Ax)_{mn} = \sum_{k,l=1,1}^{\infty,\infty} a_{mnkl}x_{k,l}.$$

Such a transformation is said to be non-negative if a_{mnkl} is nonnegative for all n, m, k and l .

DEFINITION 5. A 4-dimensional matrix A is said to be *RH*-regular if it maps every bounded P -convergent sequence into a P -convergent sequence with the same P -limit.

An *RH*-regular matrix is characterized as follows:

LEMMA 1. (Robison [31] and Hamilton [14]) *A four dimensional matrix A is *RH*-regular if and only if*

$$RH_1 : P - \lim_{mn} a_{mnkl} = 0 \text{ for each } (k, l) \in \mathbb{N}^2;$$

$$RH_2 : P - \lim_{mn} \sum_{k,l} |a_{mnkl}| = 1;$$

$$RH_3 : P - \lim_{mn} \sum_l |a_{mnkl}| = 0 \text{ for each } k;$$

$$RH_4 : P - \lim_{mn} \sum_k |a_{mnkl}| = 0 \text{ for each } l;$$

$$RH_5 : \sum_{k,l} |a_{mnkl}| \text{ is } P\text{-convergent};$$

$RH_6 : \text{there exists finite positive integers } A \text{ and } B \text{ such that } \sum_{j,k>B} a_{mnkl} < A \text{ for each } (m, n) \in \mathbb{N}^2.$

Let M be an Orlicz function and $A = (a_{mnkl})$ is a non-negative four dimensional *RH*-regular matrix. A double sequence $x = (x_{k,l})$ is called Orlicz-Quasi-Cauchy if for given $\varepsilon > 0$ there exists an $N \in \mathbb{N}$ such that

$$\frac{1}{nm} \sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}|] < \varepsilon$$

where $\Delta x_{k,l} = \max_{r,s=1 \text{ and/or } 0} \{x_{k,l} - x_{k+r,l+s}\}$, whenever $k, l > N$. This Orlicz-Quasi-Cauchy double sequence space is denoted by $(M, A, \Delta)_{OQC}$ and is defined as

$$(M, A, \Delta)_{OQC} = \left\{ x = (x_{k,l}) : \frac{1}{nm} \sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}|] < \varepsilon \right\}$$

Any P -convergent double Orlicz sequence is Orlicz-Quasi-Cauchy, so any regularly convergent double Orlicz sequence is Orlicz-Quasi-Cauchy. Any Cauchy double Orlicz sequence is Orlicz-Quasi-Cauchy. Any subsequence of a P -convergent double sequence is P -convergent. Any subsequence of a Cauchy double Orlicz sequence is

Cauchy. But for Orlicz-Quasi-Cauchy double sequence, the situation is different. There are subsequence of Orlicz-Quasi-Cauchy double sequence which are not Orlicz-Quasi-Cauchy.

DEFINITION 6. A double Orlicz function M defined on a double subset $A \times A$ of R^2 into R is called double sequentially continuous at a point L of $A \times A$ if $M(\Delta x)$ is P -convergent to $M(L)$, whenever $\Delta x = (\Delta x_{k,l})$ is a P -convergent double Orlicz sequence of points in $A \times A$ with P -limit L . If M is double Orlicz sequentially continuous at every point of $A \times A$, we say M is double Orlicz sequentially continuous on $A \times A$.

We note that any continuous function at a point L of $A \times A$ is also double Orlicz sequentially continuous on $A \times A$. The converse is also true.

2. Continuity and uniformly continuity properties

In this section we made an effort to obtain interesting results related to continuity, sequentially continuity and uniform continuity of double Orlicz-Quasi-Cauchy sequences.

THEOREM 1. *If double Orlicz function M defined on a double subset $A \times A$ of R^2 is double Orlicz sequentially continuous at L , then it is continuous.*

Proof. Suppose M is not continuous at L . Then there is an $\varepsilon_0 > 0$ such that for any $\delta > 0$ there exists an x_δ so that $|\Delta x_1(\delta) - L_1| < \delta$ and $|\Delta x_2(\delta) - L_2| < \delta$ but $|M(\Delta x_1(\delta) - \Delta x_2(\delta)) - M(L)| \geq \varepsilon_0$. It is easy to construct a convergent double Orlicz sequence with limit L whose transformed sequence is not convergent to $M(L)$. Thus M is not double Orlicz sequentially continuous at L . This contradiction completes the proof of the theorem. \square

From the above theorem we conclude that a double-Orlicz function M defined on a double subset $A \times A$ of R^2 is double Orlicz sequentially continuous at a point L if and only if it is continuous.

THEOREM 2. *If double-Orlicz function M defined on a double subset $A \times A$ of R^2 preserves double Orlicz-Quasi-Cauchy sequences from $A \times A$, then it is continuous.*

Proof. Suppose that M preserves double Orlicz-Quasi-Cauchy sequences from $A \times A$. Let $x = (x_{i,j})$ be a double sequence define by

$$\begin{array}{cccc} x_{1,1} & x_{1,2} & x_{1,3} & \cdots \\ x_{2,1} & x_{2,2} & x_{2,3} & \cdots \\ x_{3,1} & x_{3,2} & x_{3,3} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array}$$

be any P -convergent double sequence with P -limit L . Applying the double difference operator, the above double sequence reduced to

$$\begin{array}{ccccccc}
 \Delta x_{1,1}L & \Delta x_{1,2}L & \Delta x_{1,3}L & \cdots & & & \\
 L & L & L & L & L & L & \cdots \\
 \Delta x_{2,1}L & \Delta x_{2,2}L & \Delta x_{3,3}L & \cdots & & & \\
 L & L & L & L & L & L & \cdots \\
 \Delta x_{3,1}L & \Delta x_{3,2}L & \Delta x_{3,3}L & \cdots & & & \\
 L & L & L & L & L & L & \cdots \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots
 \end{array}$$

is also P -convergent with P -limit L . Since any convergent double-Orlicz sequence space is Orlicz-Quasi-Cauchy sequence, so the transformed sequence $M(\Delta x) = M|\Delta x_{k,l}|$ of the sequence Δx is Orlicz-Quasi-Cauchy. Thus it follows that

$$\begin{array}{ccccccc}
 M|\Delta x_{1,1}|M(L) & M|\Delta x_{1,2}|M(L) & M|\Delta x_{1,3}|M(L) & \cdots & & & \\
 M(L) & M(L) & M(L) & M(L) & M(L) & M(L) & \cdots \\
 M|\Delta x_{2,1}|M(L) & M|\Delta x_{2,2}|M(L) & M|\Delta x_{3,3}|M(L) & \cdots & & & \\
 M(L) & M(L) & M(L) & M(L) & M(L) & M(L) & \cdots \\
 M|\Delta x_{3,1}|M(L) & M|\Delta x_{3,2}|M(L) & M|\Delta x_{3,3}|M(L) & \cdots & & & \\
 M(L) & M(L) & M(L) & M(L) & M(L) & M(L) & \cdots \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \cdots
 \end{array}$$

Using the non-negative four dimensional RH -regular matrix on this double sequence space, we get Orlicz-Quasi-Cauchy double sequence. From this it follows that $\{M|\Delta x_{i,j}|\}$ is a P -convergent Orlicz double sequence with P -limit $M(L)$. By combining the result of Theorem 1 we get that an Orlicz function M is Continuous. This completes the proof. \square

COROLLARY 1. *Suppose that $I \times I$ is a two dimensional interval and*

$$\begin{array}{ccccccc}
 a_{1,1} & b_{1,1} & a_{1,2} & b_{1,2} & a_{1,3} & b_{1,3} & \cdots \\
 d_{1,1} & c_{1,1} & d_{1,2} & c_{1,2} & d_{1,3} & c_{1,3} & \cdots \\
 a_{2,1} & b_{2,1} & a_{2,2} & b_{2,2} & a_{2,3} & b_{2,3} & \cdots \\
 d_{2,1} & c_{2,1} & d_{2,2} & c_{2,2} & d_{2,3} & c_{2,3} & \cdots \\
 a_{3,1} & b_{3,1} & a_{3,2} & b_{3,2} & a_{3,3} & b_{3,3} & \cdots \\
 d_{3,1} & c_{3,1} & d_{3,2} & c_{3,2} & d_{3,3} & c_{3,3} & \cdots \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots
 \end{array}$$

is a double sequence of ordered pairs in $I \times I$ with

$$\lim_i |a_{i,j} - b_{i,j}| = \lim_i |a_{i,j} - c_{i,j}| = \lim_i |a_{i,j} - d_{i,j}| = 0$$

then there exists a double Orlicz quasi Cauchy sequence $\{M|\Delta x_{i,j}|\}$ with the property that for any order pair of integers $(i, j) : i, j > 1$, then there exists an ordered pair $(\bar{i}, \bar{j}) ; \bar{i}, \bar{j} > 1$ such that

$$(a_{i,j}, b_{i,j}) = (\Delta x_{\bar{i}, \bar{j}}, \Delta x_{\bar{i}, \bar{j}+1})$$

$$(a_{i,j}, c_{i,j}) = (\Delta x_{\bar{i}, \bar{j}}, \Delta x_{\bar{i}+1, \bar{j}+1})$$

$$(a_{i,j}, d_{i,j}) = (\Delta x_{\bar{i}, \bar{j}}, \Delta x_{\bar{i}+1, \bar{j}}).$$

Proof. See [28]. \square

THEOREM 3. *Suppose that $I \times I$ is a two dimensional interval. Then a two dimensional Orlicz real-valued function is uniformly continuous on $I \times I$ if and only if it is defined on $I \times I$ and preserves double Orlicz-Quasi-Cauchy sequences from $I \times I$.*

Proof. Clearly, the two dimensional uniformly continuous function preserves double Orlicz-Quasi-Cauchy sequence.

Conversely, Suppose that M defined on $I \times I$ is not uniformly continuous. Then there exists an $\varepsilon > 0$ such that for any $\delta > 0$ there exist $(a, b), (\bar{a}, \bar{b}) \in I \times I$, with $\sqrt{(a - \bar{a})^2 + (b - \bar{b})^2} < \delta$ but $|M(a, b) - M(\bar{a}, b)| \geq \varepsilon$, $|M(a, b) - M(a, \bar{b})| \geq \varepsilon$, and $|M(a, b) - M(\bar{a}, \bar{b})| \geq \varepsilon$, respectively. Then by the corollary here exists a double Orlicz-Quasi-Cauchy sequence such that for any ordered pair $(i, j) : i \geq 1, j \geq 1$, then there exists an ordered pairs integers (\bar{i}, \bar{j}) with $a_{i,j} = \Delta x_{\bar{i}, \bar{j}}$ and $b_{i,j} = \Delta x_{\bar{i}+1, \bar{j}+1}$. This implies that

$$|M(\Delta x_{\bar{i}}, \Delta x_{\bar{j}}) - M(\Delta x_{\bar{i}+1}, \Delta x_{\bar{j}})| \geq \varepsilon,$$

$$|M(\Delta x_{\bar{i}}, \Delta x_{\bar{j}}) - M(\Delta x_{\bar{i}}, \Delta x_{\bar{j}+1})| \geq \varepsilon,$$

and

$$|M(\Delta x_{\bar{i}}, \Delta x_{\bar{j}}) - M(\Delta x_{\bar{i}+1}, \Delta x_{\bar{j}+1})| \geq \varepsilon,$$

Thus, $\{M(\Delta x_i, \Delta x_j)\}$ is not Orlicz-Quasi-Cauchy. Therefore, M does not preserves double Orlicz Quasi Cauchy sequence. \square

THEOREM 4. *Suppose that M is Orlicz double function defined on bounded double interval $I \times I$. Then M , an Orlicz function is uniformly continuous on $I \times I$ if and only if the image under M of any Cauchy double sequence in $I \times I$ is Orlicz-Quasi-Cauchy.*

Proof. By using the concept of Theorem 4, clearly the image of any double Cauchy under factorable function is Orlicz-Quasi-Cauchy. Conversely, suppose the image of every Cauchy double sequence is Orlicz-Quasi-Cauchy but the Orlicz function to be uniformly continuous. Then there exists an $\varepsilon > 0$ such that for any $\delta > 0$ there exist $(x, y), (\bar{x}, \bar{y}) \in I \times I$, with $\sqrt{(x - \bar{x})^2 + (y - \bar{y})^2} < \delta$ but $|M(x, y) - M(\bar{x}, y)| \geq \varepsilon$, $|M(x, y) - M(x, \bar{y})| \geq \varepsilon$, and $|M(x, y) - M(\bar{x}, \bar{y})| \geq \varepsilon$, respectively.

For each $(m, n) : m, n \geq 1$, for fix double sequence $(\Delta x_m, \Delta y_n)$ and $(\Delta x_m^-, \Delta y_n^-)$ in $I \times I$ with $\sqrt{(\Delta x_m - \Delta x_m^-)^2 + (\Delta y_n - \Delta y_n^-)^2} < \delta$

$$|M(\Delta x_m, \Delta y_n) - M(\Delta x_m^-, \Delta y_n)| \geq \varepsilon,$$

$$|M(\Delta x_m, \Delta y_n) - M(\Delta x_m, \Delta y_n^-)| \geq \varepsilon,$$

and

$$|M(\Delta x_m, \Delta y_n) - M(\Delta x_m^-, \Delta y_n^-)| \geq \varepsilon,$$

respectively. Since $I \times I$ is bounded there exists a P -convergent subsequence by a simple extension of Bolzano-Weierstrass theorem, say $\{\Delta x_{k,l}\}$. The following double sequence

$$\begin{matrix} \Delta x_{1,1} & \Delta y_{1,2} & \Delta x_{1,3} & \Delta y_{1,4} & \Delta x_{1,5} & \Delta y_{1,6} & \cdots \\ \Delta y_{2,1} & \Delta x_{2,2} & \Delta y_{2,3} & \Delta x_{2,4} & \Delta y_{2,5} & \Delta x_{2,6} & \cdots \\ \Delta x_{3,1} & \Delta y_{3,2} & \Delta x_{3,3} & \Delta y_{3,4} & \Delta x_{3,5} & \Delta y_{3,6} & \cdots \\ \Delta y_{4,1} & \Delta x_{4,2} & \Delta y_{4,3} & \Delta x_{4,4} & \Delta y_{4,5} & \Delta x_{4,6} & \cdots \\ \Delta x_{5,1} & \Delta y_{5,2} & \Delta x_{5,3} & \Delta y_{5,4} & \Delta x_{5,5} & \Delta y_{5,6} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{matrix}$$

is P -convergent. However the following image of Cauchy sequence

$$\begin{matrix} M(\Delta x_1, \Delta x_1) & M(\Delta y_1, \Delta y_2) & M(\Delta x_1, \Delta x_3) & M(\Delta y_1, \Delta y_4) & M(\Delta x_1, \Delta x_5) & M(\Delta y_1, \Delta y_6) & \cdots \\ M(\Delta y_2, \Delta y_1) & M(\Delta x_2, \Delta x_2) & M(\Delta y_2, \Delta y_3) & M(\Delta x_2, \Delta x_4) & M(\Delta y_2, \Delta y_5) & M(\Delta x_2, \Delta x_6) & \cdots \\ M(\Delta x_3, \Delta x_1) & M(\Delta y_3, \Delta y_2) & M(\Delta x_3, \Delta x_3) & M(\Delta y_3, \Delta y_4) & M(\Delta x_3, \Delta x_5) & M(\Delta y_3, \Delta y_6) & \cdots \\ M(\Delta y_4, \Delta y_1) & M(\Delta x_4, \Delta x_2) & M(\Delta y_4, \Delta y_3) & M(\Delta x_4, \Delta x_4) & M(\Delta y_4, \Delta y_5) & M(\Delta x_4, \Delta x_6) & \cdots \\ M(\Delta x_5, \Delta x_1) & M(\Delta y_5, \Delta y_2) & M(\Delta x_5, \Delta x_3) & M(\Delta y_5, \Delta y_4) & M(\Delta x_5, \Delta x_5) & M(\Delta y_5, \Delta y_6) & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{matrix}$$

is not a Orlicz Quasi Cauchy. Thus we have a contradiction. \square

3. Topological properties of Orlicz-Quasi-Cauchy double sequence space

This sections is devoted to the study of some topological properties like linearity and completeness of $(M, A, \Delta)_{OQC}$.

THEOREM 5. *Let $A = (a_{mnkl})$ be RH-regular matrix, M be an Orlicz function. Then an Orlicz-Quasi-Cauchy double sequence space, $(M, A, \Delta)_{OQC}$ is a linear space over the field of complex numbers \mathbb{C} .*

Proof. Let $x, y \in (M, A, \Delta)_{OQC}$ and for $\alpha, \beta \in \mathbb{C}$ there exist integers M_α and N_β such that $|\alpha| < M_\alpha$ and $|\beta| < N_\beta$. Since M is an Orlicz functions, so we have

$$\begin{aligned} \frac{1}{mn} \sum_{k,l=0, \infty}^{\infty, \infty} a_{mnkl} [M|\alpha \Delta x_{k,l} + \beta \Delta y_{k,l}|] &\leq M_\alpha \frac{1}{mn} \sum_{k,l=0, \infty}^{\infty, \infty} a_{mnkl} [M|\Delta x_{k,l}|] \\ &+ N_\beta \frac{1}{nm} \sum_{k,l=0, \infty}^{\infty, \infty} a_{mnkl} [M|\Delta y_{k,l}|]. \end{aligned}$$

Thus, $\alpha x + \beta y \in (M, A, \Delta)_{OQC}$ for all k, l . Hence $(M, A, \Delta)_{OQC}$ is a linear space. This completes the proof of the theorem. \square

THEOREM 6. *Let M be an Orlicz function, $A = (a_{mnkl})$ be a RH-regular 4-dimensional matrix summability method such that $\sup_{mn} \sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} < \infty$, Then $(M, A, \Delta)_{OQC}$ is a complete topological linear space with the paranorm defined by*

$$g(x) = \sup_{mn} \sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}|].$$

Proof. Let (x_{kl}^q) be a Cauchy sequence in $(M, A, \Delta)_{OQC}$. Then, we write $g(x^q - x^t) \rightarrow 0$ as $q, t \rightarrow \infty$ for all m, n , we have

$$\sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}^q - \Delta x_{k,l}^t|] \rightarrow 0. \quad (1)$$

Thus for each fixed k and l as $q, t \rightarrow \infty$, since $A = (a_{mnkl})$ is nonnegative, we are granted that

$$M|\Delta x_{k,l}^q - \Delta x_{k,l}^t| \rightarrow 0$$

and by continuity of M , $(x_{k,l}^q)$ is a Cauchy sequence in \mathbb{C} for each fixed k and l .

Since \mathbb{C} is complete as $t \rightarrow \infty$, we have $x_{k,l}^q \rightarrow x_{k,l}$ for each (k, l) Now from equation (1), we have for $\varepsilon > 0$, there exists a natural number N such that

$$\sum_{k,l=0,0}^{\infty,\infty} \sum_{q,t>N} a_{mnkl} [M|\Delta x_{k,l}^q - \Delta x_{k,l}^t|] < \varepsilon \text{ for all } m, n. \quad (2)$$

Since for any fixed natural number R , from equation (2) we have

$$\sum_{k,l \leq R} \sum_{q,t>N}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}^q - \Delta x_{k,l}^t|] < \varepsilon \text{ for all } m, n$$

by letting $t \rightarrow \infty$ in the above expression we obtain

$$\sum_{k,l \leq R} \sum_{q>N}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}^q - \Delta x_{k,l}|] < \varepsilon.$$

Since R is arbitrary, by letting $R \rightarrow \infty$ we obtain

$$\sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} [M|\Delta x_{k,l}^q - \Delta x_{k,l}|]^{p_{k,l}} < \varepsilon \text{ for all } m, n.$$

Thus $g(x^q - x) \rightarrow 0$ as $q \rightarrow \infty$. This proves that $(M, A, \Delta)_{OQC}$ is a complete topological linear space. \square

THEOREM 7. *Let M be an Orlicz function, $A = (a_{mnkl})$ be a RH-regular matrix such that $\sup_{mn} \sum_{k,l=0,0}^{\infty,\infty} a_{mnkl} < \infty$ and $\beta = \lim_{t \rightarrow \infty} \frac{M(t)}{t} > \infty$. Then*

$$(A, \Delta)_{OQC} = (M, A, \Delta)_{OQC}.$$

Proof. In order to prove that $(A, \Delta)_{OQC} = (M, A, \Delta)_{OQC}$. It is sufficient to show that $(M, A, \Delta)_{OQC} \subseteq (A, \Delta)_{OQC}$. Now, let $\beta > 0$. By definition of β , we have $M(t) \geq \beta t$ for all $t \geq 0$. Since $\beta > 0$, we have $t \leq \frac{1}{\beta} M(t)$ for all $t \geq 0$.

Let $x = (x_{kl}) \in (M, A, \Delta)_{OQC}$. Thus, we have

$$\sum_{k,l=0,0}^{\infty,\infty} a_{mkl} \left[|\Delta x_{k,l}| \right] \leq \frac{1}{\beta} \sum_{k,l=0,0}^{\infty,\infty} a_{mkl} \left[M |\Delta x_{k,l}| \right].$$

This completes the proof. \square

REFERENCES

- [1] A. ALOTAIBI, M. MURSALEEN AND M. A. ALGHAMDI, *Invariant and absolute invariant means of double sequences*, J. Funct. Spaces Appl., Art. ID 465364, 9 pp., 2012. MR2923802.
- [2] A. ALOTAIBI, M. MURSALEEN AND K. RAJ, *Double sequence spaces by means of Orlicz functions*, Abstr. Appl. Anal., **2014**, Art. ID 260326, 8 pages.
- [3] H. ALTINOK, M. ET AND Y. ALTIN, *Strongly almost summable difference sequences*, Vietnam J. Math., **34** (2006), 331–339.
- [4] M. BAŞARIR AND O. SONALCAN, *On some double sequence spaces*, J. Indian Acad. Math., **21** (1999), 193–200.
- [5] T. J. BROMWICH, *An introduction to the theory of infinite series*, Macmillan and co. Ltd., New York (1965).
- [6] D. BURTON AND J. COLEMAN, *Quasi-Cauchy sequences*, Amer. Math. Monthly, **117** (2010), 328–333.
- [7] H. ÇAKALLI, *Forward continuity*, J. Comput. Anal. Appl., **13** (2011), 225–230.
- [8] H. ÇAKALLI, *Statistical ward continuity*, Appl. Math. Lett., **24** (2011), 1724–1728.
- [9] H. ÇAKALLI, *Statistical-quasi Cauchy sequences*, Math. Comput. Modelling, **54** (2011), 1620–1624.
- [10] H. ÇAKALLI AND B. HAZARIKA, *Ideal quasi-Cauchy sequences*, J. Inequal. Appl., **2012** (2012), pages 11.
- [11] H. DUTTA, *A characterization of the class of statistically pre-Cauchy double sequences of fuzzy numbers*, Appl. Math. Inf. Sci., **7** (2013), 1437–1440.
- [12] H. KIZMAZ, *On certain sequence spaces*, Canad. Math. Bull., **24** (1981), 169–176.
- [13] J. LINDENSTRAUSS AND L. TZAFRIRI, *On Orlicz sequence spaces*, Israel J. Math., **10** (1971), 379–390.
- [14] H. J. HAMILTON, *Transformation of multiple sequences*, Duke Math. J., **2** (1936), 29–60.
- [15] G. H. HARDY, *On the convergence of certain multiple series*, Proc. Camb. Phil., Soc., **19** (1904), 86–95.
- [16] B. HAZARIKA, *Second order ideal-ward continuity*, Int. J. Anal., Vol. 2014, Article ID 480918, 4 pages.
- [17] B. HAZARIKA, *ϕ -statistically quasi Cauchy sequences*, J. Egyptian Math. Soc., **24** (2) (2016), 239–243.
- [18] M. ISIK, *On statistical convergence of generalized difference sequences*, Soochow J. Math., **30** (2004), 197–205.
- [19] F. MÓRICZ, *Extension of the spaces c and c_0 from single to double sequences*, Acta Math. Hungar., **57** (1991), 129–136.
- [20] F. MÓRICZ AND B. E. RHOADES, *Almost convergence of double sequences and strong regularity of summability matrices*, Math. Proc. Camb. Phil. Soc., **104** (1988), 283–294.
- [21] M. MURSALEEN, *Almost strongly regular matrices and a core theorem for double sequences*, J. Math. Anal. Appl., **293** (2004), 523–531.
- [22] M. MURSALEEN AND O. H. H. EDELY, *Statistical convergence of double sequences*, J. Math. Anal. Appl., **288** (2003), 223–231.
- [23] M. MURSALEEN AND O. H. H. EDELY, *Almost convergence and a core theorem for double sequences*, J. Math. Anal. Appl., **293** (2004), 532–540.

- [24] A. PRINGSHEIM, *Zur theorie der zweifach unendlichen zahlenfolgen*, Math. Ann., **53** (1900), 289–321.
- [25] R. F. PATTERSON, *Analogues of some fundamental theorems of summability theory*, Internat. J. Math. Math. Sci., **23** (2000), 1–9.
- [26] R. F. PATTERSON, *Four dimensional matrix characterization P -convergence fields of summability methods*, Appl. Math. Comput., **219** (2013), 6783–6791.
- [27] R. F. PATTERSON, *A theorem on entire four dimensional summability methods*, Appl. Math. Comput., **219** (2013), 7777–7782.
- [28] R. F. PATTERSON AND H. ÇAKALLI, *Quasi Cauchy double sequences*, DOI 10.1515/tmj-2015-0023.
- [29] K. RAJ AND A. KILICMAN, *On certain generalized paranormed spaces*, J. Inequal. Appl., (2015), 2015: 37.
- [30] K. RAJ, S. JAMWAL AND S. K. SHARMA, *New classes of generalized sequence spaces defined by an Orlicz function*, J. Comput. Anal. Appl., **15** (2013), 730–737.
- [31] G. M. ROBISON, *Divergent double sequences and series*, Trans. Amer. Math. Soc., **28** (1926), 50–73.
- [32] B. C. TRIPATHY, *Generalized difference paranormed statistically convergent sequences defined by Orlicz function in a locally convex spaces*, Soochow J. Math., **30** (2004), 431–446.
- [33] B. C. TRIPATHY, *Statistically convergent double sequences*, Tamkang J. Math., **34** (2003), 231–237.
- [34] B. C. TRIPATHY, M. ET, Y. ALTIN AND S. MAHANTA, *Generalized difference paranormed sequence spaces defined by Orlicz function in a locally convex space*, Indian J. Math., **48** (2006), 187–199.
- [35] M. ZELTSER, *Investigation of double sequence spaces by soft and hard analytical methods*, Diss. Math. Univ. Tartu., **25**, Tartu University Press, Univ. of Tartu, Faculty of Mathematics and Computer Science, Tartu (2001).

(Received December 21, 2016)

Kuldip Raj
 Department of Mathematics
 Shri Mata Vaishno Devi University
 Katra-182320, J&K, India
 e-mail: kuldipraj68@gmail.com

Ayhan Esi
 Department of Mathematics
 Adiyaman University
 Adiyaman-02040, Turkey
 e-mail: aesi23@hotmail.com

Seema Jamwal
 Department of Applied Sciences
 Yogananda College of Engineering and Technology
 Gurha Brahmana (Patoli), Jammu-181205, J&K, India
 e-mail: seemajamwal18@gmail.com