

A p -FREE ℓ^p -INEQUALITY

GRAHAME BENNETT

(Communicated by P. Mercer)

Abstract. We show how certain simple ℓ^p -inequalities may be proved by “ignoring the p .”

We prove the following result, which confirms a conjecture made in [3].

THEOREM. *Suppose that a, b, c, d, w, x, y, z are positive numbers. Then the inequality*

$$a^p + b^p + c^p + d^p \leq w^p + x^p + y^p + z^p \quad (1)$$

is valid whenever $|p| \geq 1$, and it reverses direction whenever $|p| \leq 1$, if and only if the following five conditions are satisfied:

$$a + b + c + d = w + x + y + z \quad (2)$$

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} = \frac{1}{w} + \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \quad (3)$$

$$abcd = wxyz \quad (4)$$

$$\max\{a, b, c, d\} \leq \max\{w, x, y, z\} \quad (5)$$

and

$$\min\{a, b, c, d\} \geq \min\{w, x, y, z\}. \quad (6)$$

Moreover, inequality (1) is then strict, except when $p = -1, 0$ or 1 , or the sets $\{a, b, c, d\}$ and $\{w, x, y, z\}$ coincide.

The theorem is an example of a p -free ℓ^p -inequality. We have here an “ ℓ^p -inequality”, (1), valid for all real values of p (in the directions indicated), yet the inequality is equivalent to certain hypotheses, (2)–(6), that make no explicit mention of p ; that are, in effect, “ p -free.”

Mathematics subject classification (2000): 26D15.

Keywords and phrases: ℓ^p -inequality, majorization, higher order convexity.

COROLLARY. (cf. Lemma 1 and Theorem 4 of [1]). *Suppose that p is fixed. Then the sequence*

$$\frac{(n+1)^p}{(n+1)^p - n^p} \quad (n = 1, 2, \dots) \tag{7}$$

is convex if $p \geq 1$ or $-1 \leq p < 0$ and concave if $0 < p \leq 1$ or $p \leq -1$.

Proof. The assertion of the corollary is equivalent to the inequality

$$2[(n+1)^2(n+3)]^p + 2[n(n+2)^2]^p \leq [(n+1)(n+2)^2]^p + [n(n+2)(n+3)]^p + [(n+1)^2(n+2)]^p + [n(n+1)(n+3)]^p \tag{8}$$

being valid whenever $|p| \geq 1$, and reversing direction whenever $|p| \leq 1$. \square

Proof of theorem. (Necessity). If inequality (1) holds as stated, there must be equality when $p = \pm 1$, so (2) and (3) are guaranteed. To deduce (4), (5) and (6), we first rephrase (1) in terms of L^p -means,

$$\left(\frac{a^p + b^p + c^p + d^p}{4}\right)^{\frac{1}{p}} \leq \left(\frac{w^p + x^p + y^p + z^p}{4}\right)^{\frac{1}{p}}. \tag{9}$$

It is clear that (9) is valid whenever $p \geq 1$ or $-1 \leq p < 0$ and that the inequality reverses direction whenever $p \leq -1$ or $0 < p \leq 1$. Making $p \rightarrow \infty$ in (9), the means approach the corresponding maxima ([6], §2.3.4) forcing (5) to hold. (6) follows similarly by making $p \rightarrow -\infty$. To prove (4), we make $p \rightarrow 0$ in (9), whereupon the L^p -means are replaced by the corresponding geometric means ([6], §2.3.3). When $p \rightarrow 0^-$ we deduce that

$$(abcd)^{\frac{1}{4}} \leq (wxyz)^{\frac{1}{4}}, \tag{10}$$

and, when $p \rightarrow 0^+$, that (10) is reversed.

(Sufficiency). This, of course, is the gist of the theorem: inequality (1) holds for all real p if it holds at just five “points”, $p = 0, \pm 1$ and $\pm\infty$.

We assume throughout that the sets $\{a, b, c, d\}$ and $\{w, x, y, z\}$ are disjoint; otherwise they coincide and the theorem is then trivial. (If they have a point in common, say $d = z$, then the polynomial

$$(t-a)(t-b)(t-c) = t^3 - (a+b+c)t^2 + \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) abct - abc$$

is the same as $(t-w)(t-x)(t-y)$ by hypotheses (2), (3) and (4).)

For the remainder of the proof we assume, as we may, that the sets $\{a, b, c, d\}$ and $\{w, x, y, z\}$ are each arranged in descending order:

$$a \geq b \geq c \geq d \quad \text{and} \quad w \geq x \geq y \geq z. \tag{11}$$

It then follows, in fact, that

$$w > a \geq b > x \geq y > c \geq d > z. \tag{12}$$

The first and last inequalities are consequences of hypothesis (5) and (6); strictness therein follows from the discussion above.

We justify the third inequality by assuming the opposite,

$$b \leq x, \tag{13}$$

and showing that this leads to a contradiction. It follows that

$$a \leq w \tag{14} \quad \text{by (5)}$$

$$a + b \leq w + x \tag{15} \quad \text{by (13) and (14)}$$

$$a + b + c \leq w + x + y \tag{16} \quad \text{by (2) and (6)}$$

$$a + b + c + d = w + x + y + z \tag{17} \quad \text{by (2).}$$

These inequalities, in conjunction with (11), are precisely the ones stipulated by Hardy, Littlewood and Pólya ([6], page 45) in order that the majorization

$$(a, b, c, d) \prec (w, x, y, z) \tag{18}$$

be valid. We deduce from Theorem 108 of [6] that the inequality

$$\varphi(a) + \varphi(b) + \varphi(c) + \varphi(d) < \varphi(w) + \varphi(x) + \varphi(y) + \varphi(z) \tag{19}$$

holds whenever $\varphi : [z, w] \rightarrow \mathbb{R}$ is strictly convex. But this assertion is contradicted by hypothesis (3) when we take $\varphi(t) = 1/t$ in (19).

A similar argument, via the majorization

$$\left(\frac{1}{d}, \frac{1}{c}, \frac{1}{b}, \frac{1}{a}\right) \prec \left(\frac{1}{z}, \frac{1}{y}, \frac{1}{x}, \frac{1}{w}\right), \tag{20}$$

justifies the fifth inequality, $y > c$, in (12).

We deduce from (12) that

$$\int_z^d \frac{g(t)}{t^2} dt + \int_x^b \frac{g(t)}{t^2} dt \leq \int_c^y \frac{g(t)}{t^2} dt + \int_a^w \frac{g(t)}{t^2} dt \tag{21}$$

is valid whenever $g : [z, w] \rightarrow \mathbb{R}$ is a 3-convex function (i.e. the third-order divided differences, $g[\alpha, \beta, \gamma, \delta]$, are all non-negative).

To see this, we consider the quadratic function, Q , that agrees with g at d, y and b . It is known that Q alternates successively **above** and below g on the intervals $[z, d]$, $[d, y]$, $[y, b]$ and $[b, w]$. (This striking observation, due to Bullen ([5], Theorem 5), is the analogue, for 3-convex functions, of the familiar fact that *the graph of a convex function lies always beneath its chords*.) In particular, we have

$$g(t) \leq Q(t) \quad \text{if } t \in [z, d] \text{ or } t \in [x, b]$$

and

$$g(t) \geq Q(t) \quad \text{if } t \in [c, y] \text{ or } t \in [a, w].$$

It suffices, therefore, to prove inequality (21) with g replaced by Q . But (21) is trivially satisfied by *any* quadratic function (being then an identity), courtesy of hypotheses (2), (3) and (4).

Applying (21) to the function

$$g(t) = \begin{cases} pt^{p+1} & \text{if } |p| \geq 1 \\ -pt^{p+1} & \text{if } |p| \leq 1 \end{cases} \tag{22}$$

(which is 3-convex since $g'''(t) \geq 0$), we obtain the inequality

$$d^p - z^p + b^p - x^p \leq y^p - c^p + w^p - a^p \tag{23}$$

when $|p| \geq 1$ and its reversal when $|p| \leq 1$. This completes our proof of sufficiency.

The last sentence of the theorem, concerning cases of equality in (1), is justified by observing that the function (22) is strictly 3-convex except when $p = -1$, or 0 or 1. [A simple modification of Bullen’s proof shows that the graph of Q lies strictly above that of g on the intervals $[z, d)$ and (y, b) , and strictly below on (d, y) and $(b, w]$, whenever g is strictly 3-convex. It follows that inequality (21), and hence also (23), is then strict.] \square

Our proof shows rather more than has been stated. Indeed, (21) suggests a new kind of Majorization, one that is not considered in [3]. *The inequality*

$$f(a) + f(b) + f(c) + f(d) \leq f(w) + f(x) + f(y) + f(z) \tag{24}$$

is valid for all functions f such that

$$(t^2 f'(t))' \quad \text{is convex} \tag{25}$$

if and only if hypotheses (2)—(6) are in effect.

This is seen by replacing g in (21) by $t^2 f'(t)$, and using the characterization of 3-convex functions given in Proposition 1 of [3].

There are other p -free ℓ^p -inequalities in the literature. The simplest example, via the Theory of Majorization, is described on page 820 of [1]. A second example, with applications to Moment Theory, is given in [2], while a third, solving a problem of Knuth, appears in [4]. These all deal with 3-tuples, however, and they do not extend to higher dimensions. Some interest attaches, therefore, to the fact that our theorem applies to 4-tuples.

It was natural, in seeking such a theorem, to focus attention on the main result of [1]: *the sequence*

$$\frac{1^p + 2^p + \dots + n^p}{n^p} \quad (n = 1, 2, \dots) \tag{26}$$

is convex if $p \geq 1$ or $p \leq 0$, concave otherwise. (The inequalities implicit in (26), after all, show that the Theory of Majorization produces p -free ℓ^p -inequalities *only* for 1-, 2- and 3-tuples. See section 5 of [1].)

This led us back to our corollary, which is an important component in the proof of (26). (See sections 2 and 3 of [1].) It is easy to check that hypotheses (2)–(6) are satisfied by the variables in (8); indeed, it was this observation that led us to the formulation of our conjecture. The corollary, of course, may be proved independently (see Lemma 1 of [1], where it appears as a calculus exercise). But our theorem is a vastly more general result, and the proof given here extends readily to provide p -free ℓ^p -inequalities in any dimension.

REFERENCES

- [1] G. BENNETT, *Sums of powers and the meaning of ℓ^p* , Houston J. Math., **32**, 3 (2006), 801–831.
- [2] G. BENNETT, *An inequality from Moment Theory*, Positivity, **11**, 2 (2007), 231–238.
- [3] G. BENNETT, *Some forms of Majorization*, to appear in Houston J. Math.
- [4] G. BENNETT, *A problem of Knuth*, submitted to Amer. Math. Monthly.
- [5] P.S. BULLEN, *A criterion for n -convexity*, Pacific J. Math., **36** (1971), 81–98.
- [6] G.H. HARDY, J.E. LITTLEWOOD AND G. PÓLYA, *Inequalities. 2nd edition*, Cambridge University Press, Cambridge, 1967.

(Received October 20, 2008)

Grahame Bennett
Department of Mathematics
Indiana University
Bloomington, IN 47405
U. S. A.
e-mail: bennettg@indiana.edu