

THE PROBABILISTIC STABILITY FOR A FUNCTIONAL NONLINEAR EQUATION IN A SINGLE VARIABLE

DOREL MIHEȚ

*Dedicated to Professor Josip Pečarić
on the occasion of his 60th birthday*

Abstract. We use the fixed point method to prove the probabilistic Hyers–Ulam and generalized Hyers–Ulam–Rassias stability for the nonlinear equation $f(x) = \Phi(x, f(\eta(x)))$ where the unknown is a mapping f from a nonempty set S to a probabilistic metric space (X, F, T_M) and $\Phi : S \times X \rightarrow X$, $\eta : S \rightarrow X$ are two given functions.

1. Introduction and Preliminaries

The first stability theorem was proved by Hyers in [14], who affirmatively solved a problem of Ulam and proved that the additive Cauchy functional equation is “stable in the sense of Hyers-Ulam”. We quote from the paper by Hyers [14]:

THEOREM 1.1. (Hyers [14], Theorem 1) *Let E and E' be Banach spaces and let $f(x)$ be a δ -linear transformation of E into E' , that is, a mapping $f : E \rightarrow E'$ such that $\|f(x+y) - f(x) - f(y)\| < \delta$, $\forall x, y \in E$. Then the limit $l(x) = \lim_{n \rightarrow \infty} f(2^n x)/2^n$ exists for each x in E , $l(x)$ is a linear transformation, and the inequality $\|f(x) - l(x)\| \leq \delta$ is true for all x in E . Moreover $l(x)$ is the only linear transformation satisfying this inequality.*

Subsequently the result of Hyers has been generalized by considering unbounded control functions. Significant results have been obtained by Aoki, Bourgin, Rassias and Găvruta ([2], [5], [32], [11]). The stability problems of various functional equations have been investigated in last decades by a number of authors, see e.g., [10], [15], [16], [9], [4], [12], [1], [7], [27], [30], [36], [19]. The generalized Hyers-Ulam-Rassias stability of the additive Cauchy functional equation and Jensen functional in probabilistic and fuzzy normed spaces has recently been studied in [28], [23] and [26].

Baker ([3]) used the fixed point method to study the stability of the nonlinear functional equation $f(x) = F(x, f(\eta(x)))$.

Mathematics subject classification (2000): 39B52, 39B82, 47H10, 54E70.

Keywords and phrases: Functional equation, fixed points, Hyers-Ulam stability, probabilistic metric space.

THEOREM 1.2. (Baker [3], Theorem 2) *Let S be a nonempty set and (X, d) be a complete metric space. Let $\eta : S \rightarrow S, F : S \times X \rightarrow X$. Suppose that there is $k \in (0, 1)$ such that*

$$d(F(x, u), F(x, v)) \leq kd(u, v), \forall x \in S, \forall u, v \in X,$$

and let $f : S \rightarrow X, \delta > 0$ be such that

$$d(f(x), F(x, f(\eta(x)))) \leq \delta, \forall x \in S.$$

Then there is a unique mapping $f_s : S \rightarrow X$ with the properties

$$f_s(x) = F(x, f_s(\eta(x))), \forall x \in S$$

and

$$d(f(x), f_s(x)) \leq \frac{\delta}{1-k}, \forall x \in S.$$

In this paper we provide some probabilistic versions of the stability result of Baker.

2. Fixed points in generalized metric spaces and in probabilistic metric spaces

The notion of generalized complete metric space has been introduced by Luxemburg in [20].

DEFINITION 2.1. A *generalized metric* on a nonempty set X is a mapping $d : X \times X \rightarrow \mathbb{R} \cup \{+\infty\}$ satisfying the following conditions:

- i) $d(x, y) \geq 0, \forall x, y \in X; d(x, y) = 0$ iff $x = y$
- ii) $d(x, y) = d(y, x), \forall x, y \in X$
- iii) If $d(x, z)$ and $d(y, z)$ are finite, then $d(x, y)$ is finite and

$$d(x, y) \leq d(x, z) + d(y, z).$$

If (X, d) is a generalized metric space, then the relation $\sim, x \sim y$ if and only if $d(x, y) < +\infty$, is an equivalence relation on X which determines a unique decomposition $X = \cup\{X_\alpha, \alpha \in A\}$ into disjoint equivalence classes, called the canonical decomposition. If $d_\alpha = d|_{X_\alpha \times X_\alpha}$, then (X, d) is a complete generalized metric space if and only if (X_α, d_α) is a complete metric space for each $\alpha \in A$.

Recently, Radu [31] pointed out that many theorems concerning the stability of functional equations are consequences of the fixed point alternative of Margolis and Diaz [21]. On the other hand, as it is shown in [18], the fixed point theorems of the alternative on generalized metric spaces can be obtained from the corresponding fixed point theorems on appropriate metric spaces: if (X, d) is a generalized metric space, $X = \cup\{X_\alpha, \alpha \in A\}$ is its canonical decomposition and $T : X \rightarrow X$ is a mapping such that

$$d(T(x), T(y)) < +\infty \text{ whenever } d(x, y) < +\infty,$$

then T has a fixed point if and only if $T_\alpha = T|_{X_\alpha} : X_\alpha \rightarrow X_\alpha$ has a fixed point for some $\alpha \in A$ ([18], Theorem 3.1). Accordingly, we obtain:

THEOREM 2.1. [18] *Let (X, d) be a complete generalized metric space and $T : X \rightarrow X$ be a strict contraction with the Lipschitz constant k , such that $d(x_0, T(x_0)) < +\infty$ for some $x_0 \in X$. Then T has a unique fixed point in the set $Y := \{y \in X, d(x_0, y) < \infty\}$ and the sequence $(T^n(x))_{n \in \mathbb{N}}$ converges to the fixed point x^* for every $x \in Y$. Moreover, $d(x_0, T(x_0)) \leq \delta$ implies $d(x^*, x_0) \leq \frac{\delta}{1-k}$.*

The above theorem can immediately be extended to uniform spaces with the uniformity generated by the separated family $D = \{d_\alpha\}_{\alpha \in I}$ of generalized pseudometrics on X (see e.g., [29], [6]):

THEOREM 2.2. *Let X be a sequentially complete generalized uniform space and f be a mapping from X into X , with the property that for every $\alpha \in I$ there is $k_\alpha \in (0, 1)$ such that $d_\alpha(f(x), f(y)) \leq k_\alpha d_\alpha(x, y)$ for all $x, y \in X$. Suppose that there exist $x \in X$ and $\delta_\alpha > 0$ such that $d_\alpha(x, f(x)) < \delta_\alpha \forall \alpha \in I$. Then the sequence $\{x_n\}_{n>0}$, $x_n = f^n(x)$ is convergent in X and $u := \lim_{n \rightarrow \infty} f^n(x)$ is the fixed point of f . Moreover, $d_\alpha(u, x) \leq \frac{\delta_\alpha}{1-k_\alpha} \forall \alpha \in I$.*

Probabilistic contractions were first defined and studied by V.M. Sehgal in his doctoral dissertation at Wayne State University. In the following we recall some useful facts from fixed point theory of probabilistic metric spaces. For more details the reader is referred to the books [13] and [33].

A *triangular norm* (shorter *t-norm*) is a binary operation $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$ which is commutative, associative, nondecreasing in each variable and has 1 as the unit element. Basic examples are: $T_L(a, b) = \text{Max}(a + b - 1, 0)$ (*Łukasiewicz t-norm*), $T_P(a, b) = ab$ and $T_M(a, b) = \text{Min}\{a, b\}$.

Let (b_n) be a strictly increasing sequence of positive numbers with $\lim_{n \rightarrow \infty} b_n = 1$. By a *(b_n) t-norm of Hadžić type* we understand a *t-norm* T with $T(b_n, b_n) = b_n \forall n$.

A *distribution function* is a mapping $F : [0, \infty) \rightarrow [0, 1]$ with the properties: $F(0) = 0$, F is increasing and F is left continuous on $(0, \infty)$. The class of all distribution functions is denoted by Δ_+ and D_+ is the subset of Δ_+ containing the functions F which also satisfy the condition $\lim_{x \rightarrow \infty} F(x) = 1$.

A special element in D_+ is ϵ_0 ,

$$\epsilon_0(t) = \begin{cases} 0, & \text{if } t = 0 \\ 1, & \text{if } t > 0 \end{cases}.$$

A *probabilistic distance* on a nonempty set X is a mapping F from $X \times X$ to Δ_+ ($F(x, y)$ will be denoted by F_{xy}). A triple (X, F, T) , where X is a nonempty set, F is a probabilistic distance on X and T is a *t-norm*, is called a *generalized Menger space* or a *probabilistic metric space in the sense of Schweizer and Sklar* if the following conditions are satisfied:

- (PM0) $F_{xy} = \epsilon_0$ if and only if $x = y$
- (PM1) $F_{xy} = F_{yx} \forall x, y \in X$
- (PM2_M) $F_{xy}(t+s) \geq T(F_{xz}(t), F_{zy}(s)), \forall x, y, z \in X, \forall t, s > 0$.

If (X, F, T) is a generalized Menger space with $\sup T(t, t) = 1$, then the family $\{U_{\varepsilon, \lambda}\}_{\varepsilon > 0, \lambda \in (0, 1)}$, where $U_{\varepsilon, \lambda} := \{(x, y) \in X \times X : F_{xy}(\varepsilon) > 1 - \lambda\}$ is a base for a metrizable uniformity on X , named the F -uniformity and denoted by \mathcal{U}_F ([33]).

DEFINITION 2.2. ([22]) Let b be the collection of all strictly increasing sequences $(b_n)_{n \in \mathbb{N}}$, with $\lim_{n \rightarrow \infty} b_n = 1$ and let $(b_n) \in b$. A pair (S, F) , where F is a probabilistic distance on S satisfying (PM0) and (PM1) is called a *probabilistic (b_n) -structure* if the following triangle inequality takes place:

$$(PM3_b) \quad (F_{pq}(s) \geq b_n, F_{qr}(t) \geq b_n) \Rightarrow F_{pr}(s+t) \geq b_n.$$

It is easy to see that every generalized Menger space under a $(b_n)t$ -norm T of Hadžić-type is a probabilistic (b_n) -structure.

DEFINITION 2.3. ([34]) Let S be a nonempty set and F be a probabilistic distance on S . A mapping $f : S \rightarrow S$ is called a *probabilistic contraction* (or *B-contraction*) if there exists $k \in (0, 1)$ such that

$$F_{f(p)f(q)}(kt) \geq F_{pq}(t), \forall p, q \in S, \forall t > 0.$$

The following theorem is a probabilistic analogous of the fixed point theorem of Luxemburg-Yung.

THEOREM 2.3. (see [17], [35]) *Let (S, F, T) be a complete generalized Menger space with T of Hadžić-type and $f : S \rightarrow S$ be a B-contraction. Then f has a fixed point iff there is $p \in S$ such that $F_{pf(p)} \in D_+$. If $F_{pf(p)} \in D_+$, then $u := \lim_{n \rightarrow \infty} f^n(p)$ is the unique fixed point of f in the set $Y = \{q \in S : F_{pq} \in D_+\}$.*

3. Probabilistic stability of the equation $f(x) = \Phi(x, f(\eta(x)))$

In this section we study the probabilistic Hyers–Ulam and generalized Hyers–Ulam–Rassias stability for the nonlinear equation $f(x) = \Phi(x, f(\eta(x)))$ where the unknown is a mapping f from a nonempty set S to a probabilistic metric space (X, F, T_M) and $\Phi : S \times X \rightarrow X$, $\eta : S \rightarrow X$ are two given functions.

First we define the notion of probabilistic approximate solution of the given equation.

DEFINITION 3.1. A probabilistic uniform approximate solution of the equation $f(x) = \Phi(x, f(\eta(x)))$ is any function $f : S \rightarrow X$ such that

$$\lim_{t \rightarrow \infty} F_{f(u)\Phi(u, f(\eta(u)))}(t) = 1$$

uniformly on S .

The proof of our first theorem makes essential use of the following lemma, which completes Theorem 2.3 with an estimation relation in the case $T = T_M$:

LEMMA 3.1. ([24]) *Let (S, F, T_M) be a complete generalized Menger space and $f : S \rightarrow S$ be a k - B contraction. Suppose that $F_{pf(p)} \in D_+$ and let $u = \lim_{n \rightarrow \infty} f^n(p)$. Then*

$$F_{up}(t + 0) \geq F_{pf(p)}((1 - k)t), \forall t > 0.$$

THEOREM 3.1. *Let S be a nonempty set and (X, F, T_M) be a complete generalized Menger space. Suppose that the mapping $\Phi : S \times X \rightarrow X$ satisfies the condition: there exists $k \in (0, 1)$ such that*

$$F_{\Phi(u,x)\Phi(u,y)}(kt) \geq F_{xy}(t)$$

for all $u \in S, x \in X, t > 0$ and let $f : S \rightarrow X$ be a probabilistic uniform approximate solution of the given equation. Then there is a function $f_s : S \rightarrow X$ with the properties:

$$f_s(u) = \Phi(u, f_s(\eta(u))), \forall u \in S$$

and, if for some $\varepsilon \in (0, 1)$ and $\delta > 0$

$$F_{f(u)\Phi(u,f(\eta(u)))}(\delta) > 1 - \varepsilon, \forall u \in S$$

then

$$F_{f(u)f_s(u)}\left(\frac{\delta}{1 - k} + 0\right) \geq 1 - \varepsilon, \forall u \in S.$$

Proof. Let Y be the set of all mappings a from S to X and the operator of Baker $T : Y \rightarrow Y, T(a)(x) = \Phi(x, a(\eta(x)))$ for all $a \in Y$ and $x \in S$. For every $a, b \in Y$ define the distribution function \mathbf{F}_{ab} by the formula

$$\mathbf{F}_{ab}(t) = \sup_{s < t} \inf_{x \in S} F_{a(x)b(x)}(s).$$

It can immediately be proved that (Y, \mathbf{F}, T_M) is a complete generalized probabilistic metric space. We also have

$$\begin{aligned} \mathbf{F}_{T(a)T(b)}(kt) &= \sup_{s < kt} \inf_{y \in S} F_{T(a)(y)T(b)(y)}(s) \\ &= \sup_{s < t} \inf_{y \in S} F_{T(a)(y)T(b)(y)}(ks) \geq \sup_{s < t} \inf_{y \in S} F_{a(\eta(y))b(\eta(y))}(s) \geq \mathbf{F}_{ab}(t), \end{aligned}$$

that is, T is a B - contraction on (Y, \mathbf{F}, T_M) .

As the condition $\lim_{t \rightarrow \infty} F_{f(u)\Phi(u,f(\eta(u)))}(t) = 1$, uniformly on X means that $\mathbf{F}_{fT(f)} \in D_+$, from Theorem 2.3 it follows that T has a fixed point, that is, there is a selfmapping f_s of Y such that $f_s(u) = \Phi(u, f_s(\eta(u)))$, for all $u \in S$.

If $F_{f(u)\Phi(u,f(\eta(u)))}(\delta) > 1 - \varepsilon, \forall u \in S$ then $\mathbf{F}_{fT(f)}(\delta) \geq 1 - \varepsilon$. It follows that $\mathbf{F}_{ff_s}(\frac{\delta}{1 - k} + 0) \geq 1 - \varepsilon$, that is, $\inf_{u \in S} F_{f(u)f_s(u)}(\frac{\delta}{1 - k} + 0) \geq 1 - \varepsilon$, concluding that $F_{f(u)f_s(u)}(\frac{\delta}{1 - k} + 0) \geq 1 - \varepsilon, \forall u \in S. \quad \square$

The next stability result refers to mappings on probabilistic (b_n) -structures. In the proof of Theorem 3.2 we use the fixed point theorem of Monna.

LEMMA 3.1. ([25]) *If (S, F) is a (b_n) -probabilistic structure then, for every $n \in N$, the mapping*

$$r_n : S \times S \rightarrow [0, \infty], r_n(p, q) = \inf\{t > 0 \mid F_{pq}(t) \geq b_n\}$$

is a generalized pseudometric on S and the family $\{r_n\}_{n \in N}$ generates the uniformity \mathcal{U}_F . Moreover, $r_n(p, q) < \varepsilon \Rightarrow F_{pq}(\varepsilon) \geq b_n$ and $F_{pq}(\varepsilon) \geq b_n \Rightarrow r_n(p, q) \leq \varepsilon$.

THEOREM 3.2. *Let S be a nonempty set and (X, F) be a complete probabilistic (b_n) -structure. Suppose that the mapping $\Phi : S \times X \rightarrow X$ satisfies the condition:*

$$\forall n \in N \exists k_n \in (0, 1) : F_{xy}(t) \geq b_n \Rightarrow F_{\Phi(u,x)\Phi(u,y)}(knt) \geq b_n.$$

Let $f : S \rightarrow X$ be a mapping with the property:

$$\lim_{t \rightarrow \infty} F_{f(u)\Phi(u,f(u))}(t) = 1,$$

uniformly on S .

Then there is a function $f_s : S \rightarrow X$ such that

$$f_s(u) = \Phi(u, f_s(u)), \forall u \in S$$

and if

$$F_{f(u)\Phi(u,f(u))}(\delta_n) \geq b_n, \forall u \in S$$

then

$$F_{f(u)f_s(u)}\left(\frac{\delta_n}{1 - k_n}\right) \geq b_n, \forall u \in S.$$

Proof. Let again Y be the set of all mappings a from S to X . Consider the family of pseudometrics $\{r_n\}_{n \in N}$ defined in the above lemma and, for every $n \in N$ and all $a, b \in Y$,

$$d_n(a, b) = \sup\{r_n(a(x), b(x)), x \in S\}.$$

Then $\{d_n\}_{n \in N}$ is a complete family of generalized pseudometrics on Y . Consider the mapping $T : Y \rightarrow Y$, $T(a)(x) = \Phi(x, a(x))$ for all $a, b \in Y$ and $x \in S$.

Let $a, b \in Y$, $\varepsilon > 0$ and $x \in S$ be given. Then, for all $a, b \in Y$,

$$\begin{aligned} k_n r_n(a(x), b(x)) < \varepsilon &\Rightarrow r_n(a(x), b(x)) < \frac{\varepsilon}{k_n} \\ &\Rightarrow F_{a(x)b(x)}\left(\frac{\varepsilon}{k_n}\right) \geq b_n \Rightarrow F_{T(a)(x)T(b)(x)}(\varepsilon) \geq b_n \\ &\Rightarrow r_n(T(a)(x), T(b)(x)) \leq \varepsilon. \end{aligned}$$

Therefore, $r_n(T(a)(x), T(b)(x)) \leq k_n r_n(a(x), b(x)) \leq k_n d_n(a, b)$. This means that $d_n(T(a), T(b)) \leq k_n d_n(a, b)$ for all $a, b \in Y$, and now we can apply Theorem 2.2 to conclude the proof. \square

The generalized Hyers-Ulam-Rassias stability of the equation $f(x) = \Phi(x, f(\eta(x)))$ has been studied in [8]. In order to establish a similar result in probabilistic setting, we need the following lemma (cf. [26]):

LEMMA 3.2. Let X be a nonempty set, (Y, F, T_M) be a complete probabilistic metric space and G be a mapping from $X \times \mathbb{R}$ into $[0, 1]$, such that $G(x, \cdot) \in D_+$ for all x . Consider the set $E := \{g : X \rightarrow Y, g(0) = 0\}$ and the mapping d_G defined on $E \times E$ by

$$d_G(g, h) = \inf\{a \in \mathbb{R}_+, F_{g(x)h(x)}(at) \geq G(x, t) \text{ for all } x \in X \text{ and } t > 0\}$$

($\inf \emptyset = +\infty$). Then d_G is a complete generalized metric on E .

THEOREM 3.3. Let S be a nonempty set, (X, F, T_M) be a complete generalized Menger space and $\Phi : S \times X \rightarrow X$ be a mapping with the property that there is $k > 0$ such that

$$F_{\Phi(u,x)\Phi(u,y)}(kt) \geq F_{xy}(t)$$

for all $u \in S, x \in X, t > 0$. If

$$F_{f(u)\Phi(u,f(\eta(u)))} \geq \Psi_u, \forall u \in S$$

where $\Psi : S \rightarrow D_+$ ($\Psi(u)$ is denoted by Ψ_u) is a mapping with the property:

$$\exists L \in (0, 1) : \Psi_{\eta(u)}(Lt) \geq \Psi_u(kt), \forall u \in S, \forall t > 0$$

then there is a unique function $f_s : S \rightarrow X$ such that

$$f_s(u) = \Phi(u, f_s(\eta(u))), \forall u \in S$$

and $F_{f(u)f_s(u)}(t) \geq \Psi_u((1-L)t), \forall u \in S, \forall t > 0$.

Proof. Consider the set $E := \{g : S \rightarrow X\}$ equipped with the complete generalized metric D defined by

$$D(g, h) = \inf\{K \in \mathbb{R}_+, F_{g(u)h(u)}(Kt) \geq \Psi_u(t), \forall u \in S, \forall t > 0\}.$$

Let $T : E \rightarrow E, T(a)(u) = \Phi(u, a(\eta(u)))$ for all $a, b \in E$ and $u \in S$ be the Baker operator.

If we suppose that $a, b \in E$ are such that $D(a, b) < \varepsilon$ then

$$F_{a(u)b(u)}(\varepsilon t) \geq \Psi_u(t), \forall u \in S, \forall t > 0$$

and

$$\begin{aligned} F_{T(a)T(b)}(L\varepsilon t) &= F_{\Phi(u,a(\eta(u)))\Phi(u,b(\eta(u)))}(L\varepsilon t) \\ &\geq F_{a(\eta(u))b(\eta(u))}\left(\frac{L}{k}\varepsilon t\right) \geq \Psi_{\eta(u)}\left(\frac{L}{k}t\right) \geq \Psi_u(t) \end{aligned}$$

for every $u \in S$ and $t > 0$. Therefore, $D(T(a), T(b)) \leq L\varepsilon$, which implies

$$D(T(a), T(b)) \leq LD(a, b), \forall a, b \in E.$$

Next, from $F_{f(u)\Phi(u,f(\eta(u)))} \geq \Psi_u, \forall u \in S$ it follows $D(f, Tf) \leq 1$. From Theorem 2.1 we obtain a fixed point of T , that is the existence of a mapping $f_s : S \rightarrow X$ such that $f_s(u) = \Phi(u, f_s(\eta(u))), \forall u \in S$. Moreover, $D(f, f_s) \leq \frac{1}{1-L}$ that is, the estimation $F_{f(u)f_s(u)}\left(\frac{1}{1-L}t\right) \geq \Psi_u(t), \forall u \in S, \forall t > 0$ holds and, as f_s is the unique fixed point of T in the set $\{g \in E, D(f, g) < \infty\}$, f_s is the unique mapping verifying both $f_s(u) = \Phi(u, f_s(\eta(u))), \forall u \in S$ and the above estimation. \square

REFERENCES

- [1] R. P. AGARWAL, B. XU, W. ZHANG, *Stability of functional equations in single variable*, J. Math. Anal. Appl., **288** (2003), 852–869.
- [2] T. AOKI, *On the stability of the linear transformation in Banach spaces*, J. M. Soc. Japan, **2** (1950), 64–66.
- [3] J. A. BAKER, *The stability of certain functional equations*, Proc. AMS, **112** (1991), 729–732.
- [4] B. BATKO, J. TABOR, *Stability of an alternative Cauchy equation on a restricted domain*, Aequationes Mathematicae, **57** (1999), 221–232.
- [5] D. G. BOURGIN, *Classes of transformations and bordering transformations*, Bull. AMS, **57** (1951), 223–237.
- [6] G. L. CAIN JR., R. H. KASRIEL, *Fixed and Periodic Points of Local Contraction Mappings on Probabilistic Metric Spaces*, Mathematical Systems Theory, **9**, 4 (1976), 289–297.
- [7] L. CĂDARIU AND V. RADU, *Fixed points and the stability of Jensen's functional equation*, Journal Ineq. Pure Appl. Math., **4**, 1 (2003), Article 4, 1–7.
- [8] L. CĂDARIU, V. RADU, *Fixed point method for the generalized stability of functional equations in single variable*, Fixed Point Theory and Applications, volume 2008.
- [9] S. CZERWIK, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, 2002.
- [10] G. L. FORTI, *Hyers-Ulam stability of functional equations in several variables*, Aequationes Math., **50** (1995), 143–190.
- [11] P. GĂVRUȚĂ, *A generalization of the Hyers–Ulam–Rassias stability of approximately additive mappings*, J. Math. Anal. Appl., **184** (1994), 431–436.
- [12] A. GIANYI, Z. KAISER, Z. PALES, *Estimates to the stability of functional equations*, Aequationes Math., **73** (2007), 125–143.
- [13] O. HADŽIĆ, E. PAP, *Fixed Point Theory in PM Spaces*, Kluwer Academic Publ., 2001.
- [14] D. H. HYERS, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. USA, **27** (1941), 222–224.
- [15] D. H. HYERS, G. ISAC, TH. M. RASSIAS, *Stability of Functional Equations in Several Variables*, Basel, 1998.
- [16] D. H. HYERS, G. ISAC, TH. M. RASSIAS, *Topics in Nonlinear Analysis and Applications*, World Scientific Publishing Company, Singapore, New Jersey, London, 1997.
- [17] I. ISTRĂȚESCU, *On generalized Menger spaces*, Boll. UMI, **5**, 13-A (1976), 95–100.
- [18] C. F. K. JUNG, *On generalized complete metric spaces*, Bull. A.M.S., **75** (1969), 113–116.
- [19] Y.-S. JUNG, *On the stability of the functional equation $f(x+y-xy) + xf(y) + yf(x) = f(x) + f(y)$* , Mathematical Inequalities and Appl. MIA, **7**, 1 (2004), 79–85.
- [20] W. A. J. LUXEMBURG, *On the convergence of successive approximations in the theory of ordinary differential equations, II*, Nederl. Akad. Wetensch. Proc. Ser. A 61 = Indag. Math., **20** (1958), 540–546.
- [21] B. MARGOLIS, J. B. DIAZ, *A fixed point theorem of the alternative for contractions on a generalized complete metric space*, Bull. Amer. Math. Soc., **74** (1968), 305–309.
- [22] D. MIHEȚ, *B-contractions on σ -Menger spaces*, Seminar on Probability Theory and Applications (STPA) West Univ. of Timișoara **109** (1994).
- [23] D. MIHEȚ, *The fixed point method for fuzzy stability of the Jensen functional equation*, Fuzzy Sets and Systems, **160** (2009), 1663–1667.
- [24] D. MIHEȚ, *The probabilistic stability for a functional equation in a single variable*, Acta Mathematica Hungarica, **123** (2009), 249–256.
- [25] D. MIHEȚ, V. RADU, *A fixed point theorem for mappings with contractive iterate in PM spaces*, An. Univ. A. I. Cuza Iași, **XLII** (1996), 311–314.
- [26] D. MIHEȚ, V. RADU, *On the stability of the additive Cauchy functional equation in random normed spaces*, Journal of Mathematical Analysis and Applications, **343** (2008), 567–572.
- [27] A. K. MIRMOSTAFEE, M. MIRZAVAZIRI, M. S. MOSLEHIAN, *Fuzzy stability of the Jensen functional equation*, Fuzzy Sets and Systems, **159** (2008), 730–738.
- [28] A. K. MIRMOSTAFEE, M. S. MOSLEHIAN, *Fuzzy versions of Hyers-Ulam-Rassias theorem*, Fuzzy Sets and Systems, **159**, 6 (2008), 720–729.

- [29] A. F. MONNA, *Sur un theoreme de M. Luxembourg concernant les points fixes d'un classe d'applications d'un espace metrique dans lui meme*, Nederl. Akad. Wet., Proc., Ser. A, **64** (1961), 89–96.
- [30] D. POPA, *Functional inclusions on square-symmetric grupoids and Hyers-Ulam stability*, Mathematical Inequalities and Appl. MIA, **2** (2004), 125–134.
- [31] V. RADU, *The fixed point alternative and the stability of functional equations*, Fixed Point Theory, Cluj-Napoca, **IV**, 1 (2003), 91–96.
- [32] TH. M. RASSIAS, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc., **72** (1978), 297–300.
- [33] B. SCHWEIZER, A. SKLAR, *Probabilistic Metric Spaces*, North Holland, 1983.
- [34] V. M. SEHGAL, A. T. BHARUCHA-REID, *Fixed points of contraction mappings on PM-Spaces*, Math. Syst. Theory, **6** (1972), 97–100.
- [35] N. X. TAN, *Generalized Probabilistic Metric Spaces and Fixed Point Theorems*, Mathematische Nachrichten, **129**, 1 (1986), 205–218.
- [36] T. TRIF, *On the stability of a functional equation deriving from an inequality of Popoviciu for convex functions*, Journal of Mathematical Analysis and Applications, **272** (2002), 604–616.

(Received September 22, 2008)

Dorel Mihet
West University of Timișoara
Faculty of Mathematics and Computer Science
Department of Mathematics
Vasile Pârvan 4
300223, Timișoara
Romania
e-mail: mihet@math.uvt.ro