

## RESTRICTED CURVATURE IN THE MINKOWSKI PLANE

MOSTAFA GHANDEHARI

(Communicated by S. Gomis)

*Abstract.* A geometric proof for the following problem is given: Let  $E$  be the unit circle in a Minkowski plane. Let  $C$  be any continuously differentiable closed curve with length  $l(C)$  (measured in Minkowski metric). Assume  $|\kappa_e(C, \cdot)| \leq k\kappa_e(E, \cdot)$  and  $\kappa_e(E, \cdot)$  denote Euclidean curvatures. Then  $C$  can be contained in a similar copy of the unit disk translated and magnified by a factor of

$$\frac{l(C)}{4} - \frac{1}{4k}(l(E) - 4).$$

### 1. Introduction

Our main purpose in this article is to generalize the following theorem from the Euclidean plane to Minkowski planes. Minkowski spaces are simply finite dimensional normal linear spaces.

**THEOREM 1.** *A closed curve in  $R^2$  of length  $L$  and curvature bounded by  $K$  can be contained in a circle of radius  $\frac{L}{4} - \frac{(\pi-2)}{2K}$ .*

Johnson [17] used optimal control theory to prove Theorem 1. Chakerian, Johnson, and Vogt [6] gave a geometric proof of Theorem 1. Melzak [21] gave an interesting treatment of plane motion with curvature limitations. Isaacs [16] discusses pursuer evader games where the radius of curvature of the pursuer is bounded.

Preliminary definitions and concepts are discussed in Section 2. A generalization of Theorem 1 is proved in Section 3.

### 2. Preliminaries

By a plane convex body we shall mean a compact, convex subset of the Euclidean plane having a non-empty interior. We shall take a “unit circle”  $E$  for the Minkowski plane to be a centrally symmetric convex body with its center at the origin in the Euclidean plane. The *Minkowski distance* from  $x$  to  $y$  is defined by

$$\|x - y\| = \frac{\|x - y\|_e}{r}$$

---

*Mathematics subject classification* (2010): 52A38.

*Keywords and phrases:* geometric inequalities; Minkowskian geometry; restricted curvature.

where  $\|x - y\|_e$  is the Euclidean length from  $x$  to  $y$ , and  $r$  is the Euclidean radius of  $E$  in the direction of the vector  $y - x$ .

Minkowski distance defined by means of a convex body was developed by Minkowski [22]. The articles by Busemann [5] and Petty [24] contain basic concepts for the study of Minkowski geometry, as does Chapter 6 of Benson’s book [1] and Chapter 4 of Valentine’s book [30]. These last two references also contain useful background material from the theory of convex sets.

We use techniques from integral geometry. Santaló [27] is a good reference for integral geometry in the Euclidean spaces. Given a curve  $C$  in the Euclidean plane, let  $L$  denote the length of  $C$ . Crofton’s simplest formula (see Santaló [27]) is

$$\iint n \, dp \, d\theta = 2L,$$

where the integral is taken over all lines intersecting the curve. Assume  $(p, \theta)$  is the polar coordinate representation of the foot of the perpendicular from the origin to the line, and  $n$  is the number of intersections of a line with coordinates  $(p, \theta)$  with  $C$ . The differential element  $dG = dp \, d\theta$  is the integral geometric density for lines.

Chakerian [9] treats integral geometry in the Minkowski plane. We sketch the definitions he uses to develop Crofton’s simplest formula in the Minkowski plane. Assume  $E$  is “sufficiently” differentiable and has positive finite curvature everywhere. Parameterize  $E$  by twice its sectional area  $\phi$  and write the equation of  $E$  as

$$t = t(\phi), \quad 0 \leq \phi \leq 2\pi \quad \|t\| = \|t - 0\| = 1.$$

$E$  is called the *indicatrix*. Define the *isoperimetrix*  $T$  by the parametric representation

$$n(\phi) = \frac{dt(\phi)}{d(\phi)}, \quad 0 \leq \phi \leq 2\pi.$$

Define  $\lambda(\phi)$  by  $\frac{dn(\phi)}{d(\phi)} = -\lambda^{-1}(\phi)t(\phi)$ . Then the density for lines in two-dimensional Minkowski spaces is defined as follows. Let  $G = G(p, \phi)$  be parallel to the direction  $t(\phi)$ .

The equation of  $G$  is

$$[t(\phi), x] = p.$$

where  $[x - y] = x_1y_2 - x_2y_1$ . Then the density  $dG$  for lines is

$$dG = \lambda^{-1}(\phi) \, dp \, d\phi.$$

It is then shown in Chakerian [9] that the simplest formula of Crofton holds:

$$\int n \, dG = 2l$$

where  $n$  is the number of intersections of a line  $G$  with a curve  $C$ , integration is taken over all lines intersecting  $C$  and  $l$  is the Minkowskian length of  $C$ .

### 3. The Main Result

The following Theorem 2 is the generalization of Theorem 1 to the Minkowski plane. The author has used the theory of optimal control to prove Theorem 2 in [13].

**THEOREM 2.** *Let  $E$  be the unit circle in a Minkowski plane. Let  $C$  be any continuously differentiable closed curve with length  $l(C)$  (measured in the Minkowski metric). Assume  $|\kappa_e(C, \cdot)| \leq k\kappa_e(E, \cdot)$  where  $\kappa_e(C, \cdot)$  and  $\kappa_e(E, \cdot)$  denote Euclidean curvatures of the respective curves. Then  $C$  can be contained in a similar copy of the unit disk translated and magnified by a factor  $\mu \geq \frac{l(C)}{4} - \frac{1}{4k}(l(E) - 4)$*

In order to prove Theorem 2, we need the case with no restriction on curvature and two lemmas. Theorem 3 gives a bound on the size of the unit ball containing a closed curve of Minkowski length less than or equal to  $\frac{l}{4}$ .

**THEOREM 3.** *Any continuous closed curve  $C$  of Minkowski length  $l$  in an  $n$ -dimensional Minkowski space can be enclosed by a similar copy of the unit ball magnified by a factor less than or equal to  $\frac{l}{4}$ .*

The case of equality is discussed after the proof of Theorem 3. The Euclidean version of Theorem 3 was first proved in a more general form by Segre [28] and independently by Rustishauser and Samelson [26]. Nitsche [23] gives an elementary proof in the Euclidean 3-space. The proof given here is the same as the proof given in Chakerian and Klamkin [8] where they prove Theorem 3 in Euclidean space, and they give complete references related to Theorem 3. They also consider minimal covers other than the ball.

*Proof of Theorem 3.* Let  $u \neq v$  be two points on  $C$  dividing it into two arcs each of Minkowskian length  $\frac{l}{2}$ . Let  $p$  be the midpoint of the segment joining  $u$  and  $v$ . For  $\omega \in C$ , we have

$$\|\omega - p\| \leq \frac{1}{2} [\|u - \omega\| + \|v - \omega\|] \leq \frac{l}{4}. \tag{1}$$

To see the first inequality consider the central reflection of  $\omega$  through  $p$  to a point  $\omega^*$  with  $\|\omega^* - p\| = \|\omega - p\|$ . The first inequality is the a consequence of the triangle inequality applied to the triangle  $\omega^*u\omega$ . The second inequality follows since the straight line segments joining  $u$  and  $v$  to  $\omega$  have lengths less than or equal to the length from  $u$  to  $v$  along the curve.  $\square$

In Euclidean spaces, equality in (1) holds if and only if  $\omega$  is collinear with  $u$  and  $v$  in which case  $C$  is a “needle,” i.e., a line segment of length  $\frac{l}{2}$  traversed twice. If the unit ball of the Minkowski space does not contain a line segment, then the same argument applies. However, if the unit ball contains the line segments then we do not necessarily have the case of a needle.  $E$ .

Theorem 3 implies that a triangle inscribed in the unite circle of a Minkowski plane and having the center as an interior point has perimeter greater than 4. This result is due to Laugwitz [19]. We now proceed to prove two lemmas needed to prove Theorem 2. We also require a generalization of Blasche’s Rolling Theorem [3, pp. 114–119] due to Koutroufiotis [18]. Theorem 4 is Koutroufiotis’ generation of Lemma 2 which will follow.

Assuming the boundary of the unit circle  $E$  is smooth, we use integral geometry to prove the following Lemma 1. The proof given here is the same as the proof given in Chakerian [7] extended to Minkowski planes.

LEMMA 1. Consider a Minkowski plane with smooth unit circle  $E$ . Let  $\bar{C}$  be the convex hull of a closed curve  $C$ . Then

$$l(\bar{C}) \leq l(C).$$

*Proof.* Using Crofton’s formula for Minkowski planes (see Chakerian [9]), we can compute  $l(\bar{C})$  by the measure of lines intersecting  $\bar{C}$ . Any line  $G$  meeting  $\bar{C}$  in two points must  $C$  in at least two points. If not,  $C$  would be contained in the closed half plane  $H$  and determined by  $G$  and hence in the proper convex subset of  $\bar{C}$  determined by  $H$  and the boundary of  $\bar{C}$ . This yields a contradiction.

Schaer [29] gives a proof of Lemma 1 for the Euclidian plane. A continuously differentiable curve  $X$  in  $R^n$ , parameterized by arc length  $s$ , is called a  $K$ -curve if and only if  $\|X'(s_2) - X'(s_1)\| \leq K |s_2 - s_1|$  for all  $s_1$  and  $s_2$ . Hence  $K$ -curves are generalization of  $C^2$  curves with curvature bounded by  $K$ . Dubins [10] showed that among  $K$ -curves with prescribed initial and terminal points and prescribed initial and terminal vectors there exists a  $K$ -curve of minimal length. Chakerian, Johnson, and Vogt [6] prove that the convex hull of a closed  $K$ -curve in  $R^n$  is a closed  $K$ -curve. The same proof is extended to the Minkowski plane with unit circle  $E$  in lemma 2.  $\square$

Parametrize a given curve  $C$  with a yielding  $C(s)$ ,  $s$  is the Euclidean arc length along  $C$ .

Let  $E(s)$  be the point on  $E$  such that the unit tangent to  $E$  makes the same angle  $\theta(s)$  with the horizontal as the tangent to  $C$  at  $C(s)$ . Hence  $E(s)$  is the relative normal to  $C$  at  $C(s)$ .

$$\left\| \frac{dE}{ds} \right\|_e = \frac{ds_E}{ds} \tag{2}$$

where  $ds_E$  is the Euclidean arc length along  $E$ . But,

$$\lim_{s_2 \rightarrow s} \frac{\|E(s_2) - E(s)\|_e}{s_2 - s} = \left\| \frac{dE}{ds} \right\|_e = \left\| \frac{dE}{ds} \right\|_e \frac{ds_E}{ds} = \frac{ds_E}{ds} = \frac{ds_E}{d\theta} = \frac{\kappa_e(C, s)}{\kappa_e(E, s)} = \kappa_m(C, s) \tag{3}$$

Thus we define a continuously differentiable curve to be a Minkowskian  $K$ -curve if and only if

$$\|E(s_1) - E(s_2)\| \leq K |s_1 - s_2| \tag{4}$$

For all  $s_1$  and  $s_2$ . As a consequence of (3) and (4) we see that a  $C^2$  curve  $C$  with Minkowskian curvature bounded by  $K$  is Minkowskian  $K$ -curve.

Lemma 2 below shows that the convex hull of closed Minkowski  $K$ -curve is a Minkowskian  $K$ -curve.

LEMMA 2. *Let  $C$  be a closed Minkowskian  $K$ -curve in a Minkowski plane unit circle  $E$ . Let  $\bar{C}$  be the convex hull of  $C$ , then  $\bar{C}$  is a Minkowskian  $K$ -curve.*

*Proof.* Every point on  $\bar{C}$  is either a point  $C$  or else an interior point of a line segment in  $\bar{C}$  whose endpoints lie on  $C$ . At the endpoints of the line segment,  $\bar{C}$  has its tangent line parallel to the line segment. At points of  $\bar{C} \cap C$  there is a unique supporting line of  $C$ . If the supporting lines formed a cone at such a point,  $C$  could not have a derivative there.

Let  $\tau$  be an arc length parameter for  $C$  and let  $s$  be an arc length parameter for  $\bar{C}$ . Let  $E(\tau)$  and  $E(s)$  be the points on the unit circle corresponding to  $C(\tau)$  and  $\bar{C}(s)$  respectively. Let  $\varepsilon$  be a positive number and  $s_0$  a particular value of  $s$ . We show that for  $s$  sufficiently close to  $s_0$

$$|E(s_1) - E(s_0)|_e \leq (K + \varepsilon) |s_1 - s_0| .$$

Suppose not. There exists a sequence  $\{s_n\}$  convergent to  $s_0$  such that  $\|E(s_n) - E(s_0)\|_e > (K + \varepsilon) |s_n - s_0|$  for all  $n$ . If  $\bar{C}(s_0)$  is not on  $C$ , then for  $n$  large  $\bar{C}(s_n)$  is on the line segment of  $\bar{C}$  through  $\bar{C}(s_0)$ . Then  $E(s_n) = E(s_0)$  for a contradiction. Hence  $\bar{C}(s_0)$  must belong to  $\bar{C} \cap C$ . We can also suppose that for each  $n$ ,  $\bar{C}(s_n)$  belongs to  $\bar{C} \cap C$ . Otherwise, the point  $\bar{C}(s_n)$  would be interior points of the line segments of  $\bar{C}$ . Without affecting  $E(s_n)$  or increasing  $|s_n - s_0|$ , we can shift points along the segments until they meet  $C$ .

The curve  $C$  has only a finite number of branches which go through  $\bar{C}(s_0)$ . By passing to a subsequence, we can suppose that the points  $\bar{C}(s_n)$  all lie on the same branch of  $C$ . Hence there is a sequence  $\{\tau_n\}$  convergent to a parameter value of  $\tau_0$  with  $C(\tau_0) = \bar{C}(s_0)$  and  $C(\tau_n) = \bar{C}(s_n)$  for all  $n$ . By passing to a subsequence once more, we can assume that for all  $n$ ,  $E(s_n) = uE(\tau_n)$  where  $u = \pm 1$  is fixed. But  $\{E(\tau_n)\}$  converges to  $E(\tau_0)$ . Hence  $\{E(s_n)\}$  converges to a limit which, up to sign, equals  $E(s_0)$ . Since  $\bar{C}$  is a closed convex curve, it cannot reverse direction abruptly. So  $\{E(s_n)\}$  converges to  $E(s_0)$  and  $E(s_0) = uE(\tau_0)$ . Then

$$\begin{aligned} (K + \varepsilon) |s_n - s_0| < \|E(s_n) - E(s_0)\|_e &= \|uE(\tau_n) - E(\tau_0)\|_e \\ &= \|E(\tau_n) - E(\tau_0)\|_e \leq K |\tau_n - \tau_0| \end{aligned}$$

But for any positive number  $\alpha < 1$  and for  $n$  sufficiently large,

$$\|C(\tau_n) - C(\tau_0) / (\tau_n - \tau_0)\|_e > 1 - \alpha$$

Hence

$$\begin{aligned} (K + \varepsilon) |s_n - s_0| < K |\tau_n - \tau_0| < K \|C(\tau_n) - C(\tau_0)\|_e / 1 - \alpha \\ &= K \left\| \bar{C}(s_n) - \bar{C}(s_0) \right\|_e / 1 - \alpha \leq K |s_n - s_0| / 1 - \alpha . \end{aligned}$$

Hence  $K + \varepsilon < \frac{K}{1 - \alpha}$ . Letting  $\alpha$  vary, we conclude  $K + \varepsilon \leq K$  which gives a contradiction. Thus for each  $s_0$  and for each  $s$  sufficiently close to  $s_0$ ,  $\|E(s) - E(s_0)\|_e \leq (K + \varepsilon) |s - s_0|$ .

If  $s_1$  and  $s_2$  are any two values, by a compactness argument and partitioning the interval  $[s_1, s_2]$  into small enough subintervals and repeated application of the triangle inequality, it follows that  $\bar{C}$  is Minkowskian  $K$ -curve.  $\square$

Blaschke's Rolling Theorem states that if  $C$  is a  $C^2$  simple convex curve with curvature  $\kappa$  satisfying  $\kappa \leq K$  then a circle of radius  $\frac{1}{\kappa}$  rolls freely inside  $C$  in the sense that, if it touches  $C$  from inside at any point, it lies entirely within the closed convex set bounded by  $C$ .

The following Theorem 4 is a generalization of Blaschke's Rolling Theorem which will be used for the proof of Theorem 2.

**THEOREM 4.** (Koutroufiotis [18]) *Let the plane convex set  $D_1$  have as boundary a  $C^2$  curve  $C_1$  with curvature  $\kappa_1$ . Let  $C_2$  be a regular convex curve with curvature  $\kappa_2$ . Assume that  $C_1$  and  $C_2$  are tangent to each other at  $p_0$  with the same unit normal and that  $\kappa_1(p_1) < \kappa_2(p_2)$  if the unit tangents at  $p_1$  and  $p_2$  are equal. Then  $C_2$ , except for  $p_0$ , lies in  $D_1$ .*

*Proof of Theorem 2.* Let  $\bar{C}$  be the convex hull of  $C$ . Lemma 2 implies  $|\kappa_e(\bar{C}, \cdot)| \leq \kappa_e(E, \cdot) = \kappa_e(\frac{1}{\kappa}E, \cdot)$ . Let  $\bar{D}$  be the region bounded by  $\bar{C}$ . Using Theorem 4, we conclude  $\bar{D} = \frac{1}{\kappa}E + D$  for some  $D$ . Hence  $l(D) = l(\bar{C}) - \frac{1}{\kappa}l(E)$ . By Theorem 3,  $D$  can be covered by a copy of the unit circle magnified by a factor of  $\frac{1}{4}l(\bar{C}) - \frac{1}{4\kappa}l(E) + \frac{1}{\kappa}$ . Using Lemma 1,  $l(\bar{C}) \leq l(C)$ , and the fact that any region containing  $\bar{C}$  also contains  $C$ , we conclude that  $C$  can be covered by a copy of the unit circle  $E$  magnified by a factor of  $\frac{1}{4}l(\bar{C}) - \frac{1}{4\kappa}l(E) + \frac{1}{\kappa} = \frac{l(C)}{4} - \frac{1}{4\kappa}(l(E) - 4)$ .  $\square$

#### REFERENCES

- [1] R. V. BENSON, *Euclidean Geometry and Convexity*, McGraw-Hill, 1966.
- [2] O. BIBERSTEIN, *Éléments de géométrie différentielle Minkowskienne, Thèse*, Université de Montréal, 1957.
- [3] W. BLASCHKE, *Kreis und Kugel*, Viet, Leipzig, 1916; reprint, Chelsea, New York, 1949.
- [4] V. G. BOLTYANSKII, *Mathematical Methods of Optimal Control*, Nauka, Moscow, 1966; English transl., Holt, Rinehart, and Winston, New York, 1948.
- [5] H. BUSEMANN, *The foundations of Minkowskian geometry*, *Comm. Math. Helv.*, 24 (1950), pp. 156–187.
- [6] G. D. CHAKERIAN, H. H. JOHNSON, AND A. VOGT, *A geometric inequality for plane curves with restricted curvature*, *Proc. Amer. Math. Soc.*, 57 (1976), pp. 122–126.
- [7] G. D. CHAKERIAN, *A characterization of curves of constant width*, *Amer. Math. Monthly*, 81 (1974), pp. 146–149.
- [8] G. D. CHAKERIAN, AND M. S. KLAMKIN, *Minimal covers for closed curves*, *Math. Mag.*, 46 (1973), pp. 55–61.
- [9] G. D. CHAKERIAN, *Integral geometry in the Minkowski plane*, *Duke Math. J.*, 29 (1962), pp. 375–382.
- [10] L. E. DUBINS, *On curves of minimal length with a constraint on average curvature, and with prescribed initial and terminal positions and tangents*, *Amer. J. Math.*, 79(1957), pp. 497–516.
- [11] W. H. FLEMING, AND R. W. RISHEL, *Deterministic and stochastic Optimal Control*, Springer-Verlag, 1975.
- [12] I. M. GELFAND, AND S. V. FOMIN, *Calculus of Variations*, Prentice Hall, 1963.

- [13] M. GHANDEHARI, *Controlling Curvature in the Minkowski Plane*, (English summary), J. Math. Anal. Appl. 252 (2000), no. 2, 951–958.
- [14] P. A. GRIFFITHS, *Exterior Differential Systems and the Calculus of Variations*, Progress in Mathematics, Birkhäuser, 1983.
- [15] R. HERMANN, *Differential Geometry and the Calculus of Variations*, Academic Press, 1968.
- [16] R. ISAACS, *Differential Games*, Wiley, 1965, pp. 8–13.
- [17] H. H. JOHNSON, *An application of the maximum principle to the geometry of plane curves*, Proc. Amer. Math. Soc., 44(1974), pp. 432–435.
- [18] D. KOUTROFIOTIS, *On Blaschke's rolling theorems*, Arch. Math., 23 (1972), pp. 655–660.
- [19] D. LAUGWITZ, *Konvexe mittelpunktsbereiche und normierte Räume*, Math. Z., 61 (1954), pp. 235–244.
- [20] G. LEITMANN, *The Calculus of Variations and optimal Control*, Plenum press, 1981.
- [21] Z. A. MELZAK, *Plane motion with curvature limitation*, J. Soc. Indust. Appl. Math., 9 (1961), pp. 422–432.
- [22] H. MINKOWSKI, *Theorie der Konvexen Körper, Insbesondere Begründung Ihres Oberflächen Bgiffs*, Ges. Abhandl., Leipzig-Berlin, Vol. 2 (1911), pp. 131–229.
- [23] J. C. C. NITSCHKE, *The smallest sphere containing a rectifiable curve*, Amer. Math. Monthly, 78 (1971), pp. 881–888.
- [24] C. M. PETTY, *On the geometry of the Minkowski plane*, Riv. Mat. Univ. Parma, Vol. 6 (1955), pp. 269–292.
- [25] L. S. PONTRYAGIN, V. G. BOLTYANSKII, R. V. GAMKERLIDZE, AND E. F. MISHCHENKO, *The Mathematical Theory of Optimal Processes*, translated from the Russian by K. N. Tirogoff, edited by L. W. Neustadt, Interscience Publishers, New York, 1962.
- [26] H. RUTISHAUSER, AND H. SAMELSON, *Sur le rayon d'une sphere dont la surface contient une courbe fermée*, C. R. Acad. Sci. Paris, 227 (1948), pp. 755–757.
- [27] S. L. A. SANTALÓ, *Introduction to Integral Geometry*, Paris, Hermann, 1953.
- [28] B. SEGRE, *Sui circoli heodetici di una superfilcie a curvatura, che contengono nell'interno una linea assegnata*, Boll. Un. Mat. Ital., 13 (1934), pp. 279–283.
- [29] J. SCHAEER, *An "obvious" but useful theorem about closed curves*, Math. Mag., (1972), pp. 154–155.
- [30] F. A. VALENTINE, *Convex Sets*, Krieger, 1976.

(Received February 10, 2009)

Mostafa Ghandehari  
Department of Mathematics  
University of Texas at Arlington  
Arlington, Texas 76019  
U.S.A.