

## SOME PROPERTIES FOR A CLASS OF SYMMETRIC FUNCTIONS AND APPLICATIONS

YU-MING CHU, WEI-FENG XIA AND TIE-HONG ZHAO

(Communicated by G. Toader)

*Abstract.* For  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ , the symmetric function  $F_n(x, r)$  is defined by

$$F_n(x, r) = F_n(x_1, x_2, \dots, x_n; r) = \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r x_{i_j}}{\sum_{j=1}^r (1 + x_{i_j})},$$

where  $r = 1, 2, \dots, n$  and  $i_1, i_2, \dots, i_r$  are positive integers. In this article, the Schur convexity, Schur harmonic convexity and Schur multiplicative convexity of  $F_n(x, r)$  are discussed. As applications, some inequalities are established by use of the theory of majorization.

### 1. Introduction

Throughout the paper we use  $\mathbb{R}^n$  denote the  $n$ -dimensional Euclidean space, and  $\mathbb{R}_+^n = \{(x_1, x_2, \dots, x_n) : x_i > 0, i = 1, 2, \dots, n\}$ . In particular, we use  $\mathbb{R}$  to denote  $\mathbb{R}^1$ .

For  $x = (x_1, x_2, \dots, x_n)$ ,  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}_+^n$  and  $\alpha > 0$ , let

$$\begin{aligned} x + y &= (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n), \\ xy &= (x_1 y_1, x_2 y_2, \dots, x_n y_n), \\ \alpha x &= (\alpha x_1, \alpha x_2, \dots, \alpha x_n), \\ x^\alpha &= (x_1^\alpha, x_2^\alpha, \dots, x_n^\alpha), \\ \frac{1}{x} &= \left( \frac{1}{x_1}, \frac{1}{x_2}, \dots, \frac{1}{x_n} \right) \\ \log x &= (\log x_1, \log x_2, \dots, \log x_n) \end{aligned}$$

and

$$e^x = (e^{x_1}, e^{x_2}, \dots, e^{x_n}).$$

---

*Mathematics subject classification* (2010): 05E05, 26B25.

*Keywords and phrases:* Symmetric function, Schur convex, Schur harmonic convex, Schur multiplicatively convex, theory of majorization.

This work was supported by NNSF Grant of China (No. 60850005) and the NSF Grant of Zhejiang Province (No. D7080080, Y7080185, Y607128).

For  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ ,  $r \in N$  and  $r \leq n$ , the Hamy symmetric function  $H_n(x, r)$  is defined by T. Hara, M. Uchiyama and S. Takahasi [1] as follows:

$$H_n(x, r) = H_n(x_1, x_2, \dots, x_n; r) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \left( \prod_{j=1}^r x_{i_j} \right)^{\frac{1}{r}},$$

where  $i_1, i_2, \dots, i_n \in N$ .

Corresponding to this is the  $r$ -th order Hamy mean

$$\sigma_n(x, r) = \sigma_n(x_1, x_2, \dots, x_n; r) = \frac{(n-r)!r!}{n!} H_n(x, r).$$

T. Hara, M. Uchiyama and S. Takahasi [1] established the following refinement of the classical arithmetic and geometric means inequalities:

$$G_n(x) = \sigma_n(x, n) \leq \sigma_n(x, n-1) \leq \dots \leq \sigma_n(x, 2) \leq \sigma_n(x, 1) = A_n(x),$$

where  $A_n(x) = \frac{1}{n} \sum_{i=1}^n x_i$  and  $G_n(x) = \left( \prod_{i=1}^n x_i \right)^{\frac{1}{n}}$  denote the classical arithmetic and geometric means of  $x$ , respectively.

The paper [2] by H. T. Ku, M. C. Ku and X. M. Zhang contains some interesting inequalities including the fact that  $(\sigma_n(x, r))^{\frac{1}{r}}$  is log-concave. More results can be found in the book [3] by P. S. Bullen.

Recently, the Schur convexity of the Hamy symmetric function  $H_n(x, r)$  was discussed and some analytic inequalities were established by K. Z. Guan [4].

The main purpose of this paper is to discuss the Schur convexity, Schur harmonic convexity and Schur multiplicative convexity for the symmetric function

$$F_n(x, r) = F_n(x_1, x_2, \dots, x_n; r) = \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r x_{i_j}}{\sum_{j=1}^r (1 + x_{i_j})}. \quad (1.1)$$

As applications, some inequalities are established by use of the theory of majorization.

For the reader convenience, we recall several definitions.

DEFINITION 1.1. Let  $E \subseteq \mathbb{R}^n$  be a set, a real-valued function  $F$  on  $E$  is said to be Schur convex if

$$F(x_1, x_1, \dots, x_n) \leq F(y_1, y_2, \dots, y_n)$$

for each pair of  $n$ -tuples  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  in  $E$ , such that  $x \prec y$ , that is

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}, \quad k = 1, 2, \dots, n-1$$

and

$$\sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]},$$

where  $x_{[i]}$  denotes the  $i$ th largest component in  $x$ .  $F$  is called Schur concave if  $-F$  is Schur convex.

DEFINITION 1.2. Let  $E \subseteq \mathbb{R}_+^n$  be a set, a real-valued function  $F$  on  $E$  is said to be Schur multiplicatively convex if

$$F(x_1, x_2, \dots, x_n) \leq F(y_1, y_2, \dots, y_n)$$

for each pair of  $n$ -tuples  $x = (x_1, x_2, \dots, x_n)$  and  $y = (y_1, y_2, \dots, y_n)$  in  $E$ , such that  $\log x \prec \log y$ .  $F$  is called Schur multiplicatively concave if  $\frac{1}{F}$  is Schur multiplicatively convex.

DEFINITION 1.3. Let  $E \subseteq \mathbb{R}_+^n$  be a set, a real-valued function  $F$  on  $E$  is said to be Schur harmonic convex if

$$F(x_1, x_2, \dots, x_n) \leq F(y_1, y_2, \dots, y_n) \quad (1.2)$$

for each pair of  $n$ -tuples  $x = (x_1, x_2, \dots, x_n)$  and  $y = (y_1, y_2, \dots, y_n)$  in  $E$ , such that  $\frac{1}{x} \prec \frac{1}{y}$ .  $F$  is called a Schur harmonic concave on  $E$  if inequality (1.2) is reversed.

The Schur convexity was introduced by I. Schur [5] in 1923, G. H. Hardy, J. E. Littlewood and G. Pólya were also interested in some inequalities that are related to the Schur convexity [6]. Recently, the Schur multiplicative convexity was introduced and investigated in paper [7, 8].

Very recently, the Schur harmonic convexity was introduced by Y. M. Chu and Y. P. Lv [9], and the Schur harmonic convexity for the Hamy Symmetric function  $H_n(x, r)$  was discussed. To investigate the Schur harmonic convexity for the symmetric function  $F_n(x, r)$  is one of the main purpose in this article.

## 2. Lemmas

In order to establish our main results we need several lemmas, which we present in this section.

LEMMA 2.1. ([5]) *Suppose that  $f : \mathbb{R}_+^n \rightarrow \mathbb{R}_+$  is a continuous symmetric function. If  $f$  is differentiable in  $\mathbb{R}_+^n$ , then  $f$  is Schur convex in  $\mathbb{R}_+^n$  if and only if*

$$(x_i - x_j) \left( \frac{\partial f}{\partial x_i} - \frac{\partial f}{\partial x_j} \right) \geq 0 \quad (2.1)$$

for all  $i, j = 1, 2, \dots, n$  and  $x = (x_1, \dots, x_n) \in \mathbb{R}_+^n$ . And  $f$  is Schur concave in  $\mathbb{R}_+^n$  if and only if inequality (2.1) is reversed. Here  $f$  is a symmetric function in  $\mathbb{R}_+^n$  which means that  $f(Px) = f(x)$  for any  $x \in \mathbb{R}_+^n$  and any  $n \times n$  permutation matrix  $P$ .

REMARK 2.1. Since  $f$  is symmetric, the Schur's condition in Lemma 2.1, that is (2.1) can be reduced to

$$(x_1 - x_2) \left( \frac{\partial f}{\partial x_1} - \frac{\partial f}{\partial x_2} \right) \geq 0.$$

LEMMA 2.2. ([7, 8]) Suppose that  $f : \mathbb{R}_+^n \rightarrow \mathbb{R}_+$  is a continuous symmetric function. If  $f$  is differentiable in  $\mathbb{R}_+^n$ , then  $f$  is Schur multiplicatively convex in  $\mathbb{R}_+^n$  if and only if

$$(\log x_1 - \log x_2) \left( x_1 \frac{\partial f}{\partial x_1} - x_2 \frac{\partial f}{\partial x_2} \right) \geq 0 \quad (2.2)$$

for all  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ . And  $f$  is Schur multiplicatively concave in  $\mathbb{R}_+^n$  if and only if inequality (2.2) is reversed.

LEMMA 2.3. ([9]) Suppose that  $f : \mathbb{R}_+^n \rightarrow \mathbb{R}_+$  is a continuous symmetric function. If  $f$  is differentiable in  $\mathbb{R}_+^n$ , then  $f$  is Schur harmonic convex in  $\mathbb{R}_+^n$  if and only if

$$(x_1 - x_2) \left( x_1^2 \frac{\partial f}{\partial x_1} - x_2^2 \frac{\partial f}{\partial x_2} \right) \geq 0 \quad (2.3)$$

for all  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ . And  $f$  is Schur harmonic concave in  $\mathbb{R}_+^n$  if and only if inequality (2.3) is reversed.

LEMMA 2.4. ([4, 8, 10]) Suppose that  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$  and  $\sum_{i=1}^n x_i = s$ . If  $c \geq s$ , then

$$\frac{c-x}{\frac{nc}{s}-1} = \left( \frac{c-x_1}{\frac{nc}{s}-1}, \frac{c-x_2}{\frac{nc}{s}-1}, \dots, \frac{c-x_n}{\frac{nc}{s}-1} \right) \prec (x_1, x_2, \dots, x_n) = x.$$

LEMMA 2.5. ([10]) Suppose that  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$  and  $\sum_{i=1}^n x_i = s$ . If  $c \geq 0$ , then

$$\frac{c+x}{\frac{nc}{s}+1} = \left( \frac{c+x_1}{\frac{nc}{s}+1}, \frac{c+x_2}{\frac{nc}{s}+1}, \dots, \frac{c+x_n}{\frac{nc}{s}+1} \right) \prec (x_1, x_2, \dots, x_n) = x.$$

LEMMA 2.6. ([11]) Suppose that  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$  and  $\sum_{i=1}^n x_i = s$ . If  $0 \leq \lambda \leq 1$ , then

$$\frac{s-\lambda x}{n-\lambda} = \left( \frac{s-\lambda x_1}{n-\lambda}, \frac{s-\lambda x_2}{n-\lambda}, \dots, \frac{s-\lambda x_n}{n-\lambda} \right) \prec (x_1, x_2, \dots, x_n) = x.$$

### 3. Main Results

**THEOREM 3.1.**  $F_n(x, r)$  is Schur concave in  $\mathbb{R}_+^n$ .

*Proof.* The proof is divided into four cases.

*Case 1.* If  $r = 1$ , then (1.1) leads to

$$F_n(x, 1) = F_n(x_1, x_2, \dots, x_n; 1) = \prod_{i=1}^n \frac{x_i}{1+x_i} \tag{3.1}$$

and

$$(x_1 - x_2) \left( \frac{\partial F_n(x, 1)}{\partial x_1} - \frac{\partial F_n(x, 1)}{\partial x_2} \right) = -\frac{(x_1 - x_2)^2(1+x_1+x_2)}{x_1x_2(1+x_1)(1+x_2)} F_n(x, 1) \leq 0.$$

*Case 2.* If  $r = n$ , then

$$F_n(x, n) = F_n(x_1, x_2, \dots, x_n; n) = \frac{\sum_{i=1}^n x_i}{\sum_{i=1}^n (1+x_i)} \tag{3.2}$$

and

$$(x_1 - x_2) \left( \frac{\partial F_n(x, n)}{\partial x_1} - \frac{\partial F_n(x, n)}{\partial x_2} \right) = 0.$$

*Case 3.* If  $n \geq 3$  and  $r = 2$ , then

$$\begin{aligned} F_n(x, 2) &= F_n(x_1, x_2, \dots, x_n; 2) \\ &= F_{n-1}(x_2, x_3, \dots, x_n; 2) \frac{(x_1 + x_2) \prod_{j=3}^n (x_1 + x_j)}{[(1+x_1) + (1+x_2)] \prod_{j=3}^n [(1+x_1) + (1+x_j)]} \\ &= F_{n-1}(x_1, x_3, \dots, x_n; 2) \frac{(x_2 + x_1) \prod_{j=3}^n (x_2 + x_j)}{[(1+x_1) + (1+x_2)] \prod_{j=3}^n [(1+x_2) + (1+x_j)]} \end{aligned} \tag{3.3}$$

and

$$\begin{aligned} &(x_1 - x_2) \left( \frac{\partial F_n(x, 2)}{\partial x_1} - \frac{\partial F_n(x, 2)}{\partial x_2} \right) \\ &= -2(x_1 - x_2)^2 F_n(x, 2) \sum_{j=3}^n \frac{x_1 + x_2 + 2x_j + 2}{(x_1 + x_j)(x_2 + x_j)(2 + x_1 + x_j)(2 + x_2 + x_j)} \leq 0. \end{aligned}$$

*Case 4.* If  $n \geq 4$  and  $3 \leq r \leq n - 1$ , then

$$\begin{aligned} F_n(x, r) &= F_n(x_1, x_2, \dots, x_n; r) \\ &= F_{n-1}(x_2, x_3, \dots, x_n; r) \frac{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} (x_1 + \sum_{j=1}^{r-1} x_{i_j})}{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} [(1+x_1) + \sum_{j=1}^{r-1} (1+x_{i_j})]} \\ &\quad \times \frac{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} (x_1 + x_2 + \sum_{j=1}^{r-2} x_{i_j})}{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} [(1+x_1) + (1+x_2) + \sum_{j=1}^{r-2} (1+x_{i_j})]} \end{aligned} \tag{3.4}$$

$$\begin{aligned}
&= F_{n-1}(x_2, x_3, \dots, x_n; r) \frac{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} (x_2 + \sum_{j=1}^{r-1} x_{i_j})}{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} [(1+x_2) + \sum_{j=1}^{r-1} (1+x_{i_j})]} \\
&\quad \times \frac{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} (x_2 + x_1 + \sum_{j=1}^{r-2} x_{i_j})}{\prod_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} [(1+x_1) + (1+x_2) + \sum_{j=1}^{r-2} (1+x_{i_j})]}.
\end{aligned}$$

and

$$\begin{aligned}
&(x_1 - x_2) \left( \frac{\partial F_n(x, r)}{\partial x_1} - \frac{\partial F_n(x, r)}{\partial x_2} \right) \\
&= -r(x_1 - x_2)^2 F_n(x, r) \sum_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} \frac{r + x_1 + x_2 + 2 \sum_{j=1}^{r-1} x_{i_j}}{\Delta} \leq 0,
\end{aligned}$$

where

$$\Delta = \left( x_1 + \sum_{j=1}^{r-1} x_{i_j} \right) \left( x_2 + \sum_{j=1}^{r-1} x_{i_j} \right) \left( r + x_1 + \sum_{j=1}^{r-1} x_{i_j} \right) \left( r + x_2 + \sum_{j=1}^{r-1} x_{i_j} \right). \quad (3.5)$$

□

Therefore, Theorem 3.1 follows from Lemma 2.1 and Remark 2.1 together with Cases 1-4.

**THEOREM 3.2.**  $F_n(x, r)$  is Schur harmonic convex in  $\mathbb{R}_+^n$ .

*Proof.* We divided the proof into four cases.

*Case I.* If  $r = 1$ , then (3.1) leads to

$$(x_1 - x_2) \left( x_1^2 \frac{\partial F_n(x, 1)}{\partial x_1} - x_2^2 \frac{\partial F_n(x, 1)}{\partial x_2} \right) = \frac{(x_1 - x_2)^2}{(1+x_1)(1+x_2)} F_n(x, 1) \geq 0.$$

*Case II.* If  $r = n$ , then (3.2) yields

$$(x_1 - x_2) \left( x_1^2 \frac{\partial F_n(x, n)}{\partial x_1} - x_2^2 \frac{\partial F_n(x, n)}{\partial x_2} \right) = \frac{n(x_1 - x_2)^2 (x_1 + x_2)}{[\sum_{i=1}^n (1+x_i)]^2} \geq 0.$$

*Case III.* If  $n \geq 3$  and  $r = 2$ , then (3.3) implies

$$\begin{aligned}
&(x_1 - x_2) \left( x_1^2 \frac{\partial F_n(x, 2)}{\partial x_1} - x_2^2 \frac{\partial F_n(x, 2)}{\partial x_2} \right) \\
&= 2(x_1 - x_2)^2 F_n(x, 2) \left[ \frac{1}{x_1 + x_2 + 2} \right. \\
&\quad \left. + \sum_{j=3}^n \frac{2x_1x_2 + 2x_1x_j + 2x_2x_j + 2x_1x_2x_j + x_1x_j^2 + x_2x_j^2}{(x_1+x_j)(x_2+x_j)(2+x_1+x_j)(2+x_2+x_j)} \right] \geq 0.
\end{aligned}$$

Case IV. If  $n \geq 4$  and  $3 \leq r \leq n - 1$ , then from (3.4) we get

$$\begin{aligned} & (x_1 - x_2) \left( x_1^2 \frac{\partial F_n(x, r)}{\partial x_1} - x_2^2 \frac{\partial F_n(x, r)}{\partial x_2} \right) \\ &= r(x_1 - x_2)^2 F_n(x, r) \\ & \quad \times \left[ \sum_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} \frac{x_1 + x_2}{(x_1 + x_2 + \sum_{j=1}^{r-2} x_{i_j})(r + x_1 + x_2 + \sum_{j=1}^{r-2} x_{i_j})} \right. \\ & \quad \left. + \sum_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} \frac{\Theta}{\Delta} \right] \geq 0, \end{aligned}$$

where  $\Theta = (x_1 + x_2) \left( \sum_{j=1}^{r-1} x_{i_j} \right)^2 + 2x_1 x_2 \sum_{j=1}^{r-1} x_{i_j} + r(x_1 + x_2) \sum_{j=1}^{r-1} x_{i_j} + r x_1 x_2$  and  $\Delta$  is defined as in (3.5).

Therefore, Theorem 3.2 follows from Lemma 2.3 and Cases I-IV.  $\square$

Next, we denote by

$$\Omega_n(t, r) = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : t \leq x_i \leq \sqrt{(r-1)^2 t^2 + r(r-1)t}\}$$

for  $t > 0$  and  $2 \leq r \leq n - 1$ .

For the Schur multiplicative convexity or concavity of the symmetric function  $F_n(x, r)$ , we have the following Theorem 3.3.

- THEOREM 3.3.** (i)  $F_n(x, 1)$  is Schur multiplicatively concave in  $\mathbb{R}_+^n$ ;  
(ii)  $F_n(x, n)$  is Schur multiplicatively convex in  $\mathbb{R}_+^n$ ;  
(iii) If  $n \geq 3$  and  $2 \leq r \leq n - 1$ , then  $F_n(x, r)$  is Schur multiplicatively convex in  $\Omega_n(t, r)$  for any  $t > 0$ .

*Proof.* (i) From (3.1) we clearly see that

$$\begin{aligned} & (\log x_1 - \log x_2) \left( x_1 \frac{\partial F_n(x, 1)}{\partial x_1} - x_2 \frac{\partial F_n(x, 1)}{\partial x_2} \right) \\ &= - \frac{(x_1 - x_2)(\log x_1 - \log x_2)}{(1 + x_1)(1 + x_2)} F_n(x, 1) \leq 0. \end{aligned} \tag{3.6}$$

Therefore, Theorem 3.3 (i) follows from (3.6) and Lemma 2.2.

(ii) Equation (3.2) leads to

$$\begin{aligned} & (\log x_1 - \log x_2) \left( x_1 \frac{\partial F_n(x, n)}{\partial x_1} - x_2 \frac{\partial F_n(x, n)}{\partial x_2} \right) \\ &= \frac{n(x_1 - x_2)(\log x_1 - \log x_2)}{[\sum_{i=1}^n (1 + x_i)]^2} \geq 0. \end{aligned} \tag{3.7}$$

Therefore, Theorem 3.3 (ii) follows from (3.7) and Lemma 2.2.

(iii) We divided the proof into two cases.

Case A. If  $n \geq 3$  and  $r = 2$ , then (3.3) implies

$$\begin{aligned}
 & (\log x_1 - \log x_2) \left( x_1 \frac{\partial F_n(x, 2)}{\partial x_1} - x_2 \frac{\partial F_n(x, 2)}{\partial x_2} \right) \\
 &= 2(x_1 - x_2)(\log x_1 - \log x_2) F_n(x, 2) \left[ \frac{1}{(x_1 + x_2)(x_1 + x_2 + 2)} \right. \\
 & \quad \left. + \sum_{j=3}^n \frac{x_j^2 + 2x_j - x_1 x_2}{(x_1 + x_j)(x_2 + x_j)(x_1 + x_j + 2)(x_2 + x_j + 2)} \right] \geq 0
 \end{aligned} \tag{3.8}$$

for any  $t > 0$  and  $x = (x_1, x_2, \dots, x_n) \in \Omega_n(t, 2)$ .

Case B. If  $n \geq 4$  and  $3 \leq r \leq n - 1$ , then (3.4) yields

$$\begin{aligned}
 & (\log x_1 - \log x_2) \left( x_1 \frac{\partial F_n(x, r)}{\partial x_1} - x_2 \frac{\partial F_n(x, r)}{\partial x_2} \right) \\
 &= r(x_1 - x_2)(\log x_1 - \log x_2) F_n(x, r) \\
 & \quad \times \left[ \sum_{3 \leq i_1 < i_2 < \dots < i_{r-2} \leq n} \frac{1}{(x_1 + x_2 + \sum_{j=1}^{r-2} x_{i_j})(r + x_1 + x_2 + \sum_{j=1}^{r-2} x_{i_j})} \right. \\
 & \quad \left. + \sum_{3 \leq i_1 < i_2 < \dots < i_{r-1} \leq n} \frac{r \sum_{j=1}^{r-1} x_{i_j} + (\sum_{j=1}^{r-1} x_{i_j})^2 - x_1 x_2}{\Delta} \right] \geq 0
 \end{aligned}$$

for any  $t > 0$  and  $x = (x_1, x_2, \dots, x_n) \in \Omega_n(t, r)$ , where  $\Delta$  is defined as in (3.5).

Therefore, Theorem 3.3 (iii) follows from Cases A and B together with Lemma 2.2.  $\square$

#### 4. Applications

In this section, we establish some inequalities by use of Theorems 3.1–3.3 and the theory of majorization.

The following Theorem 4.1 easily follows from Theorems 3.1 and 3.2 together with Lemmas 2.4–2.6.

**THEOREM 4.1.** *Suppose that  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$  and  $\sum_{i=1}^n x_i = s$ . If  $c_1 \geq s, c_2 \geq 0, 0 \leq \lambda \leq 1$  and  $r \in \{1, 2, \dots, n\}$ , then*

- (1)  $F_n(x, r) \leq F_n\left(\frac{c_1 - x}{\frac{nc_1}{s} - 1}, r\right)$ ;
- (2)  $F_n\left(\frac{1}{x}, r\right) \geq F_n\left(\frac{\frac{nc_1}{s} - 1}{c_1 - x}, r\right)$ ;
- (3)  $F_n(x, r) \leq F_n\left(\frac{c_2 + x}{\frac{nc_2}{s} + 1}, r\right)$ ;



$$(4) F_n\left(\frac{1}{x}, r\right) \geq F_n\left(\frac{\frac{nc_2}{s} + 1}{c_2 + x}, r\right);$$

$$(5) F_n(x, r) \leq F_n\left(\frac{s - \lambda x}{n - \lambda}, r\right);$$

$$(6) F_n\left(\frac{1}{x}, r\right) \geq F_n\left(\frac{n - \lambda}{s - \lambda x}, r\right).$$

For  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ , if we denote  $H_n(x) = \frac{n}{\sum_{i=1}^n \frac{1}{x_i}}$ , then we clearly see that

$$(A_n(x), A_n(x), \dots, A_n(x)) \prec (x_1, x_2, \dots, x_n) \tag{4.1}$$

and

$$\left(\frac{1}{H_n(x)}, \frac{1}{H_n(x)}, \dots, \frac{1}{H_n(x)}\right) \prec \left(\frac{1}{x_1}, \frac{1}{x_2}, \dots, \frac{1}{x_n}\right). \tag{4.2}$$

It follows from (1.1) and (4.1)–(4.2) together with Theorems 3.1–3.2 that the following Theorem 4.2 is obvious.

**THEOREM 4.2.** *Suppose that  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ . If  $r \in \{1, 2, \dots, n\}$ , then*

$$(1) \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r x_{i_j}}{r + \sum_{j=1}^r x_{i_j}} \leq \left[ \frac{A_n(x)}{A_n(1+x)} \right]^{\frac{n!}{r!(n-r)!}};$$

$$(2) \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r \frac{1}{x_{i_j}}}{r + \sum_{j=1}^r \frac{1}{x_{i_j}}} \geq \left[ \frac{1}{A_n(1+x)} \right]^{\frac{n!}{r!(n-r)!}};$$

$$(3) \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r \frac{1}{x_{i_j}}}{r + \sum_{j=1}^r \frac{1}{x_{i_j}}} \leq \left[ \frac{1}{1 + H_n(x)} \right]^{\frac{n!}{r!(n-r)!}};$$

$$(4) \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \frac{\sum_{j=1}^r x_{i_j}}{r + \sum_{j=1}^r x_{i_j}} \geq \left[ \frac{H_n(x)}{1 + H_n(x)} \right]^{\frac{n!}{r!(n-r)!}}.$$

If we take  $r = 1$  in Theorem 4.2(1), (3) and (4), respectively, then we get

**COROLLARY 4.1.** *If  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ , then*

$$(1) \frac{G_n(x)}{G_n(1+x)} \leq \frac{A_n(x)}{A_n(1+x)};$$

$$(2) G_n(1+x) \geq 1 + H_n(x);$$

$$(3) \frac{G_n(x)}{G_n(1+x)} \geq \frac{H_n(x)}{1 + H_n(x)}.$$

**REMARK 4.1.** Inequality in Corollary 4.1(1) was proved by V. Govedarica and M. V. Jovanović in [12].

REMARK 4.2. If we take  $\sum_{i=1}^n x_i = 1$  in Corollary 4.1(1), then we get the Weierstrass inequality [13, p. 260]

$$\prod_{i=1}^n (x_i^{-1} + 1) \geq (n+1)^n.$$

THEOREM 4.3. Let  $\mathcal{A} = A_1 A_2 \cdots A_{n+1}$  be a  $n$ -dimensional simplex in  $\mathbb{R}^n$  and  $P$  be an arbitrary point in the interior of  $\mathcal{A}$ . If  $B_i$  is the intersection point of straight line  $A_i P$  and hyperplane  $\Sigma_i = A_1 A_2 \cdots A_{i-1} A_{i+1} \cdots A_{n+1}$ ,  $i = 1, 2, \dots, n+1$ . Then for  $r \in \{1, 2, \dots, n+1\}$  we have

$$(1) \quad \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n+1} \frac{\sum_{j=1}^r \frac{PB_{i_j}}{A_{i_j} B_{i_j}}}{r + \sum_{j=1}^r \frac{PB_{i_j}}{A_{i_j} B_{i_j}}} \leq \left( \frac{1}{n+2} \right)^{\frac{(n+1)!}{r!(n-r+1)!}};$$

$$(2) \quad \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n+1} \frac{\sum_{j=1}^r \frac{PA_{i_j}}{A_{i_j} B_{i_j}}}{r + \sum_{j=1}^r \frac{PA_{i_j}}{A_{i_j} B_{i_j}}} \leq \left( \frac{n}{2n+1} \right)^{\frac{(n+1)!}{r!(n-r+1)!}};$$

$$(3) \quad \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n+1} \frac{\sum_{j=1}^r \frac{A_{i_j} B_{i_j}}{PB_{i_j}}}{r + \sum_{j=1}^r \frac{A_{i_j} B_{i_j}}{PB_{i_j}}} \geq \left( \frac{n+1}{n+2} \right)^{\frac{(n+1)!}{r!(n-r+1)!}};$$

$$(4) \quad \prod_{1 \leq i_1 < i_2 < \dots < i_r \leq n+1} \frac{\sum_{j=1}^r \frac{A_{i_j} B_{i_j}}{PA_{i_j}}}{r + \sum_{j=1}^r \frac{A_{i_j} B_{i_j}}{PA_{i_j}}} \geq \left( \frac{n+1}{2n+1} \right)^{\frac{(n+1)!}{r!(n-r+1)!}}.$$

*Proof.* It is easy to see that  $\sum_{i=1}^{n+1} \frac{PB_i}{A_i B_i} = 1$  and  $\sum_{i=1}^{n+1} \frac{PA_i}{A_i B_i} = n$ , these identities imply that

$$\left( \frac{1}{n+1}, \frac{1}{n+1}, \dots, \frac{1}{n+1} \right) \prec \left( \frac{PB_1}{A_1 B_1}, \frac{PB_2}{A_2 B_2}, \dots, \frac{PB_{n+1}}{A_{n+1} B_{n+1}} \right) \quad (4.3)$$

and

$$\left( \frac{n}{n+1}, \frac{n}{n+1}, \dots, \frac{n}{n+1} \right) \prec \left( \frac{PA_1}{A_1 B_1}, \frac{PA_2}{A_2 B_2}, \dots, \frac{PA_{n+1}}{A_{n+1} B_{n+1}} \right). \quad (4.4)$$

Therefore, Theorem 4.3 follows from (4.3), (4.4), Theorem 3.1, Theorem 3.2 and (1.1).  $\square$

REMARK 4.3. D. S. Mitrinović, J. E. Pečarić and V. Volenec [14, p. 473–479] established a series of inequalities for  $\frac{PA_i}{A_i B_i}$  and  $\frac{PB_i}{A_i B_i}$ ,  $i = 1, 2, \dots, n+1$ . Obviously, our inequalities in Theorem 4.3 are different from theirs.

## REFERENCES

- [1] T. HARA, M. UCHIYAMA, S. TAKAHASI, *A refinement of various mean inequalities*, J. Inequal. Appl., **2** (1998), 387–395.
- [2] H. T. KU, M. C. KU, X. M. ZHANG, *Inequalities for symmetric means, symmetric harmonic means, and their applications*, Bull. Austral. Math. Soc., **56** (1997), 409–420.
- [3] P. S. BULLEN, *Handbook of Means and Their Inequalities*, Kluwer Academic Publishers Group, Dordrecht, 2003.
- [4] K. Z. GUAN, *The Hamy symmetric function and its generalization*, Math. Inequal. Appl., **9** (2006), 797–805.
- [5] A. W. MARSHALL, I. OLKIN, *Inequalities: Theory of Majorization and Its Applications*, Academic Press, New York, 1979.
- [6] G. H. HARDY, J. E. LITTLEWOOD, G. PÓLYA, *Some simple inequalities satisfied by convex functions*, Messenger Math., **58** (1929), 145–152.
- [7] Y. M. CHU, X. M. ZHANG, G. D. WANG, *The Schur geometrical convexity of the extended mean values*, J. Convex Anal., **15** (2008), 707–718.
- [8] K. Z. GUAN, *A class of symmetric functions for multiplicatively convex function*, Math. Inequal. Appl., **10** (2007), 745–753.
- [9] Y. M. CHU, Y. P. LV, *The Schur harmonic convexity of the Hamy symmetric function and its applications*, J. Inequal. Appl., **2009** (2009), Art. ID 838529, 10pp.
- [10] K. Z. GUAN, *Schur-convexity of the complete symmetric function*, Math. Inequal. Appl., **9** (2006), 567–576.
- [11] S. H. WU, *Generalization and sharpness of the power means inequality and their applications*, J. Math. Anal. Appl., **312** (2005), 637–652.
- [12] V. GOVEDARICA, M. JOVANOVIĆ, *On the inequalities of Ky Fan, Wang-Wang and Alzer*, J. Math. Anal. Appl., **270** (2002), 709–712.
- [13] P. S. BULLEN, *A Dictionary of Inequalities*, Longman, Harlow, 1998.
- [14] D. S. MITRINOVIĆ, J. E. PEČARIĆ, V. VOLENEC, *Recent Advances in Geometric Inequalities*, Kluwer Academic Publishers Group, Dordrecht, 1989.

(Received July 30, 2009)

Yu-Ming Chu  
 Department of Mathematics  
 Huzhou Teachers College  
 Huzhou 313000, P.R.China  
 e-mail: chuyuming@hutc.zj.cn

Wei-Feng Xia  
 School of Teachers Education  
 Huzhou Teachers College  
 Huzhou 313000, P.R.China  
 e-mail: xwf212@hutc.zj.cn

Tie-Hong Zhao  
 Institut de Mathématiques  
 Université Pierre et Marie Curie  
 4 place Jussieu  
 Paris F-75252, France  
 e-mail: zhao@math.jussieu.fr