

A SELF-ADAPTIVE PROJECTION METHOD FOR A CLASS OF VARIANT VARIATIONAL INEQUALITIES

ABDELLAH BNOUHACHEM, MUHAMMAD ASLAM NOOR, MOHAMED KHALFAOUI
AND SHENG ZHAOHAN

(Communicated by A. Čižmešija)

Abstract. In this paper, we consider the general variant variational inequality of the type: Find a vector $u^* \in \mathbb{R}^n$, such that

$$Q(u^*) \in \Omega, \quad \langle v - Q(u^*), Tu^* \rangle \geq 0, \quad \forall v \in \Omega,$$

where T, Q are operators. We suggest and analyze a very simple self-adaptive iterative method for solving this class of general variational inequalities. Under certain conditions, the global convergence of the proposed method is proved. An example is given to illustrate the efficiency and implementation of the proposed method. Preliminary numerical results show that the proposed method is applicable.

1. Introduction

The classical variational inequality, denoted by $VI(\Omega, F)$, is to determine a vector u^* in a nonempty closed convex subset Ω of the n -dimensional Euclidean space \mathbb{R}^n , such that

$$\langle v - u^*, F(u^*) \rangle \geq 0, \quad \forall v \in \Omega,$$

where F is an operator from \mathbb{R}^n into itself. Variational inequality problems are of fundamental importance for a wide range of problems in science and technology, such as mathematical programming, traffic engineering, economics and equilibrium problems, see [1–35].

It is well-known that $VI(\Omega, F)$ is equivalent to the projection equation

$$u = P_\Omega[u - \beta_k F(u)],$$

where $P_\Omega(\cdot)$ denotes the orthogonal projection map on Ω and β_k is a judiciously chosen positive step size. Thus the solution of the $VI(\Omega, F)$ is equivalent to finding a zero of the residue function

$$e(u, \beta) := u - P_\Omega[u - \beta_k F(u)].$$

We now have a variety of techniques to suggest and analyze various iterative algorithms for solving variational inequalities [1–30]. The simplest is the Goldstein's projection

Mathematics subject classification (2010): 90C33, 49J40.

Keywords and phrases: Variational inequalities, self-adaptive rules, projection method.

method [6] which, starting with any $u^0 \in \mathbb{R}^n$, iteratively updates u^{k+1} according to the formula

$$u^{k+1} = P_{\Omega}[u^k - \beta_k F(u^k)], \quad (1.1)$$

where β_k is a chosen positive step-size. In contrast with Douglas-Rachford operator splitting method [9, 17] for VI(Ω, F), this projection method can be viewed as a simple explicit method.

In this paper, we consider a class of general variational inequalities of finding $u \in \mathbb{R}^n$ such that

$$Q(u) \in \Omega, \quad \langle v - Q(u), Tu \rangle \geq 0, \quad \forall v \in \Omega, \quad (1.2)$$

where $Q, T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are given nonlinear operators and $\Omega \subset \mathbb{R}^n$ is a nonempty closed convex set. Problem of the type (1.2) was introduced and considered by Noor [19]. For the formulation, applications, numerical methods and other aspects of the general variational inequalities (1.2), see [9–29] and the references therein. In this paper, our main aim is to develop a simple method for solving problem (1.2). In order to obtain the new iterate of the proposed method, we do not need to compute additionally the value of function Q . This is very important especially in practical problems in which the cost of computing or approximating the value of function Q is very expensive. Furthermore, numerical experiments show that the proposed method may be efficient for some large scale problems, which clearly illustrates its simplicity and its efficiency.

Throughout this paper, we assume that the operator Q is strongly monotone with respect to the operator T with a positive modulus $\alpha > 0$ such that

$$\langle Tx - Ty, Q(x) - Q(y) \rangle \geq \alpha \|x - y\|^2, \quad \forall x, y \in \mathbb{R}^n$$

and Lipschitz continuous with a positive constant $L > 0$ such that

$$\|Q(x) - Q(y)\| \leq L \|x - y\|, \quad \forall x, y \in \mathbb{R}^n.$$

Note that, for $L = 1$, the operator Q is called nonexpansive. Without loss of generality, we assume that the operator T is nonexpansive, unless otherwise specified. We would like to mention that we need neither the Lipschitz constant L nor the strong monotone α to design the algorithm. We assume that the solution set of (1.2), denoted by S^* , is nonempty.

2. Preliminaries

In this section, we summarize some preliminary results which are useful in the following analysis. At first, we give some basic properties of the projection mapping. In the next step, we present an useful equivalent expression of the variational inequality problem.

LEMMA 2.1. *For a given $u \in \Omega$ and $z \in \mathbb{R}^n$, the inequality*

$$\langle u - z, v - u \rangle \geq 0, \quad \forall v \in \Omega$$

holds if and only if $u = P_{\Omega}(z)$, where P_{Ω} is the projection operator.

It follows from Lemma 2.1 that

$$\langle z - P_{\Omega}(z), P_{\Omega}(z) - v \rangle \geq 0, \quad \forall z \in \mathbb{R}^n, v \in \Omega. \tag{2.1}$$

LEMMA 2.2. *Let Ω be a closed convex set in \mathbb{R}^n , then we have*

$$\|(v - P_{\Omega}(v)) - (w - P_{\Omega}(w))\| \leq \|v - w\| \quad \forall v, w \in \mathbb{R}^n. \tag{2.2}$$

Proof. Using (2.1), we can prove that

$$\langle v - w, P_{\Omega}(v) - P_{\Omega}(w) \rangle \geq \|P_{\Omega}(v) - P_{\Omega}(w)\|^2 \quad \forall v, w \in \mathbb{R}^n,$$

by using the above inequality, we obtain

$$\begin{aligned} & \|(v - P_{\Omega}(v)) - (w - P_{\Omega}(w))\|^2 \\ &= \|v - w\|^2 - 2\langle v - w, P_{\Omega}(v) - P_{\Omega}(w) \rangle + \|P_{\Omega}(v) - P_{\Omega}(w)\|^2 \\ &\leq \|v - w\|^2 - \|P_{\Omega}(v) - P_{\Omega}(w)\|^2. \end{aligned}$$

This implies that

$$\|(v - P_{\Omega}(v)) - (w - P_{\Omega}(w))\| \leq \|v - w\|.$$

We obtain the desired result. \square

LEMMA 2.3. [32] *Denote $R_{\gamma}(u) = Q(u) - P_{\Omega}[Q(u) - \frac{1}{\gamma}Tu]$, then for all $u \in \mathbb{R}^n$ and $\gamma' \geq \gamma > 0$, it holds that*

$$\|R_{\gamma'}(u)\| \leq \|R_{\gamma}(u)\|. \tag{2.3}$$

LEMMA 2.4. [19] *$u^* \in \mathbb{R}^n$ is solution of problem (1.2) if and only if $u^* \in \mathbb{R}^n$ satisfies the relation:*

$$Q(u^*) = P_K[Q(u^*) - \beta Tu^*], \tag{2.4}$$

From Lemma 2.4, it is clear that u is solution of (1.2) if and only if u is a zero point of the function

$$r(u, \beta) := \frac{1}{\beta} \{Q(u) - P_{\Omega}[Q(u) - \beta Tu]\}. \tag{2.5}$$

It is clear that

$$u \in S^* \Leftrightarrow r(u, \beta) = 0.$$

For a given $\gamma > 0$, the equation (2.5) can be written as

$$u = u - \gamma r(u, \beta) \tag{2.6}$$

which plays an important part in the whole analysis of the proposed method.

3. Self-adaptive Method

In this section, we describe the proposed method and we prove some lemmas, which are used in the next section. Note that the Goldstein's projection scheme (1.1) can be viewed as

$$u^{k+1} = u^k - e(u^k, \beta_k). \quad (3.1)$$

By analogy, based on $r(u, \beta)$ we can construct a similar method for solving problem (1.2). Then, we consider the following iterative scheme:

ALGORITHM 3.1.

Step 0. Given a non negative sequence $\eta_k > 0$ with $\sum_{k=1}^{+\infty} \eta_k < +\infty, \beta_0 > 0, \varepsilon > 0,$

$\mu \in [0.5, 1), \delta \in (0, 1), \delta_0 \in (0, 1), \varepsilon > 0,$ and $u^0 \in \mathbb{R}^n,$ set $\gamma_0 := \beta_0$ and $k = 0.$

Step 1. If $\|r(u^k, 1)\| \leq \varepsilon$ then stop. Otherwise go to Step 2.

Step 2. If $u^k \notin S^*,$ then find the smallest non negative integer $l_k,$ such that

$$\beta_{k+1} := \frac{\gamma_k}{\mu^{l_k}} \quad \text{and} \quad u^{k+1} := u^k - r(u^k, \beta_{k+1}),$$

satisfies

$$\frac{1}{\beta_{k+1}} \|Q(u^k) - Q(u^{k+1})\|^2 \leq (2 - \delta) \langle Tu^k - Tu^{k+1}, Q(u^k) - Q(u^{k+1}) \rangle.$$

Step 3. Selection of $\gamma_{k+1},$ if

$$\frac{1}{\beta_{k+1}} \|Q(u^k) - Q(u^{k+1})\| \leq \delta_0 (\langle Tu^k - Tu^{k+1}, Q(u^k) - Q(u^{k+1}) \rangle),$$

then

$$\gamma_{k+1} := \frac{\beta_{k+1}}{1 + \eta_{k+1}},$$

otherwise

$$\gamma_{k+1} := \beta_{k+1}.$$

Set $k := k + 1,$ and go to Step 1.

REMARK 3.1. For the given $u^k,$ the new iterate of Li and Yuan's algorithm in [15] is generated by

$$u^{k+1} = u^k - \alpha_k d(u^k, \tilde{u}^k, \beta_k)$$

where

$$\tilde{u}^k = u^k - r(u^k, \beta_k)$$

$$d(u^k, \tilde{u}^k, \beta_k) = \frac{1}{\beta_k} \{Q(\tilde{u}^k) - P_\Omega[Q(u^k) - \beta_k Tu^k]\}$$

and

$$\alpha_k = \frac{\gamma_k \langle r(u^k, \beta_k), d(u^k, \tilde{u}^k, \beta_k) \rangle}{\|d(u^k, \tilde{u}^k, \beta_k)\|^2}, \quad \gamma_k \in (0, 2).$$

From Step 2, we have

$$u^{k+1} = -\frac{1}{\beta_{k+1}}\{(Q(u^k) - \beta_{k+1}Tu^k) - P_\Omega[Q(u^k) - \beta_{k+1}Tu^k]\}. \quad (3.2)$$

The main amount of computational effort when computing the new iterate is to obtain or approximate the value of the function Q at a given vector. The algorithm of Li and Yuan from [15] needs the value of Q both at \tilde{u}^k as well as at u^k . In contrast, our algorithm computes (or approximates) $Q(u^k)$ and computes the next iterate by an additional post processing step which solely involves the application of P_Ω . The cost of this post processing step is relatively minor in comparison with the cost of evaluating $Q(\tilde{u}^k)$. This is the main advantage of our method.

REMARK 3.2. The numerical performance of the method depends significantly on the initial step-size parameter β_0 . The method converges quite quickly when a proper fixed step-size parameter is chosen. However, this proper step-size parameter is unknown beforehand. If the step-size parameter is too large or too small, the number of iterations can increase significantly. To overcome this difficulty, in the case of our present work, we have used a self-adaptive technique to adjust the step-size parameter at each iteration. The main advantage of this technique is that it can adjust the step-size parameter automatically.

In the following lemma, we show that the sequence β_k is bounded both from above as well as from below.

LEMMA 3.1. *In each iteration of Algorithm 3.1, the procedure of searching β_{k+1} will terminate in finite steps. Furthermore, there is a constant $M > 0$ and there are two positive real numbers denoted by $\beta_{\min} := \beta_0 M$ and $\beta_{\max} := \frac{L^2}{(2-\delta)\alpha\mu}$ such that:*

$$\beta_{\max} \geq \beta_{k+1} \geq \beta_{\min} > 0, \quad \forall k > 0.$$

Proof. By Step 2 in the proposed method, we derive that

$$\beta_{k+1} \geq \frac{\beta_k}{1 + \eta_k}, \quad \forall k > 0.$$

Since

$$\sum_{k=1}^{+\infty} \eta_k < +\infty,$$

it follows that there is a constant $M > 0$ such that

$$\frac{1}{\prod_{k=1}^{+\infty} (1 + \eta_k)} \geq M.$$

Then

$$\beta_{k+1} \geq \beta_{\min} := \beta_0 M, \quad \forall k > 0.$$

Since the mapping Q is Lipschitz continuous with a constant L on the feasible set Ω , we have

$$\frac{1}{\beta_{k+1}} \|Q(u^k) - Q(u^{k+1})\|^2 \leq \frac{1}{\beta_{k+1}} L^2 \|u^k - u^{k+1}\|^2. \tag{3.3}$$

Because Q is strongly monotone with respect to the operator T with a constant modulus $\alpha > 0$, it yields

$$(2 - \delta)\alpha \|u^k - u^{k+1}\|^2 \leq (2 - \delta) \langle Tu^k - Tu^{k+1}, Q(u^k) - Q(u^{k+1}) \rangle. \tag{3.4}$$

From (3.3) and (3.4), it follows that the inequality of Step 2 is satisfied if

$$\beta_{k+1} \geq \frac{L^2}{(2 - \delta)\alpha}.$$

The parameter l_k in the algorithm is the minimum non negative integer fulfilling the condition of Step 2, this means that

$$\beta_{k+1} \leq \beta_{\max} := \frac{L^2}{(2 - \delta)\alpha\mu}, \quad \forall k > 0.$$

The proof is completed. \square

In the next lemma we prove that $\|r(u, \beta)\|$ is a non increasing function for $\beta > 0$. Note that the proof of this Lemma is different from that in [15, Lemma 2].

LEMMA 3.2. *For a given $u \in \mathbb{R}^n$, let $0 < \beta < \beta'$. Then it holds*

$$\|r(u, \beta')\| \leq \|r(u, \beta)\| \leq \frac{\beta'}{\beta} \|r(u, \beta')\|.$$

Proof. First, for all $v, w \in \mathbb{R}^n$ and for any nonempty closed convex set Ω , we have

$$\langle P_{\Omega}(v) - P_{\Omega}(w), v - P_{\Omega}(v) \rangle \geq 0.$$

Setting $v = Q(u) - \beta Tu$, $w = Q(u) - \beta' Tu$ and respectively, we obtain

$$\langle P_{\Omega}[Q(u) - \beta Tu] - P_{\Omega}[Q(u) - \beta' Tu], \beta' \{Q(u) - \beta Tu - P_{\Omega}[Q(u) - \beta Tu]\} \rangle \geq 0, \tag{3.5}$$

and

$$\langle P_{\Omega}[Q(u) - \beta' Tu] - P_{\Omega}[Q(u) - \beta Tu], \beta \{Q(u) - \beta' Tu - P_{\Omega}[Q(u) - \beta' Tu]\} \rangle \geq 0. \tag{3.6}$$

Using (2.5), adding (3.5) and (3.6), we get

$$\langle \beta' r(u, \beta') - \beta r(u, \beta), \beta' \beta r(u, \beta) - \beta' \beta r(u, \beta') \rangle \geq 0.$$

Thus, we have

$$\langle \beta' + \beta \rangle r(u, \beta'), r(u, \beta) \rangle \geq \beta' \|r(u, \beta')\|^2 + \beta \|r(u, \beta)\|^2.$$

It's easily to show that

$$(\beta' + \beta)\|r(u, \beta')\|^2 + (\beta' + \beta)\|r(u, \beta)\|^2 \geq 2\beta'\|r(u, \beta')\|^2 + 2\beta\|r(u, \beta)\|^2.$$

And

$$\|r(u, \beta)\|^2 \geq \|r(u, \beta')\|^2,$$

i.e.

$$\|r(u, \beta)\| \geq \|r(u, \beta')\|.$$

For the proof of the right-hand inequalities. Note that

$$\alpha\|r(u, \alpha)\| = \|R_{\frac{1}{\alpha}}(u)\|,$$

and from Lemma 2.3, we have

$$\beta\|r(u, \beta)\| = \|R_{\frac{1}{\beta}}(u)\| \leq \|R_{\frac{1}{\beta'}}(u)\| = \beta'\|r(u, \beta')\|, \tag{3.7}$$

from which, we have

$$\|r(u, \beta)\| \leq \frac{\beta'}{\beta}\|r(u, \beta')\|.$$

Therefore, the assertion of this lemma is proved. \square

4. Convergence Analysis

In this section, we consider the global convergence of the proposed method. For this purpose, we need the following result.

THEOREM 4.1. *The sequence $\{u^k\}$ generated by the proposed method satisfies*

$$\|r(u^{k+1}, \beta_{k+1})\| \leq \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right)^{\frac{1}{2}} (1 + \eta_k)\|r(u^k, \beta_k)\|. \tag{4.1}$$

Proof. It follows from (2.5) and (3.2) that

$$\begin{aligned} r(u^{k+1}, \beta_{k+1}) &= \frac{1}{\beta_{k+1}}\{(Q(u^{k+1}) - \beta_{k+1}Tu^{k+1}) - P_{\Omega}[Q(u^{k+1}) - \beta_{k+1}Tu^{k+1}]\} \\ &\quad - \frac{1}{\beta_{k+1}}\{(Q(u^k) - \beta_{k+1}Tu^k) - P_{\Omega}[Q(u^k) - \beta_{k+1}Tu^k]\}, \end{aligned} \tag{4.2}$$

substituting $v = Q(u^{k+1}) - \beta_{k+1}Tu^{k+1}$ and $w = Q(u^k) - \beta_{k+1}Tu^k$ in (2.2), we obtain

$$\begin{aligned} \|r(u^{k+1}, \beta_{k+1})\|^2 &\leq \|(Tu^{k+1} - Tu^k) - \frac{1}{\beta_{k+1}}(Q(u^{k+1}) - Q(u^k))\|^2 \\ &\leq \|Tu^{k+1} - Tu^k\|^2 - \frac{2}{\beta_{k+1}}\langle Tu^{k+1} - Tu^k, Q(u^{k+1}) - Q(u^k) \rangle \\ &\quad + \frac{1}{\beta_{k+1}^2}\|Q(u^{k+1}) - Q(u^k)\|^2. \end{aligned} \tag{4.3}$$

From Step 2, we have

$$\frac{1}{\beta_{k+1}^2} \|Q(u^k) - Q(u^{k+1})\|^2 \leq \frac{2-\delta}{\beta_{k+1}} \langle Tu^k - Tu^{k+1}, Q(u^k) - Q(u^{k+1}) \rangle. \quad (4.4)$$

Substituting (4.4) in (4.3) and using the strongly monotone of Q with respect to the operator T , we obtain

$$\begin{aligned} \|r(u^{k+1}, \beta_{k+1})\|^2 &\leq \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right) \|u^{k+1} - u^k\|^2 \\ &= \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right) \|r(u^k, \beta_{k+1})\|^2. \end{aligned} \quad (4.5)$$

Because $\beta_{k+1} \geq \frac{\beta_k}{1+\eta_k}$ and $\beta_k \geq \frac{\beta_k}{1+\eta_k}$, based on Lemma 3.2, we get

$$\|r(u^k, \beta_{k+1})\| \leq \left\| r\left(u^k, \frac{\beta_k}{1+\eta_k}\right) \right\| \leq \frac{\beta_k}{1+\eta_k} \|r(u^k, \beta_k)\|,$$

then

$$\|r(u^k, \beta_{k+1})\| \leq (1 + \eta_k) \|r(u^k, \beta_k)\|. \quad (4.6)$$

Combining (4.5) and (4.6), we have

$$\|r(u^{k+1}, \beta_{k+1})\| \leq \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right)^{\frac{1}{2}} (1 + \eta_k) \|r(u^k, \beta_k)\|.$$

Then (4.1) holds and the theorem is proved. \square

From the above theorem, we get the convergence of the proposed method as follows.

THEOREM 4.2. *The proposed method for solving (1.2) is globally convergent.*

Proof. From (4.1), we obtain

$$\|r(u^{k+1}, \beta_{k+1})\| \leq \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right)^{\frac{k+1}{2}} \prod_{i=0}^k (1 + \eta_i) \|r(u^0, \beta_0)\|.$$

Since $0 \leq 1 - \frac{\delta\alpha}{\beta_{\max}} < 1$ and $\prod_{i=0}^k (1 + \eta_i) < +\infty$, it follows that

$$\|r(u^k, \beta_k)\| \rightarrow 0,$$

and by using Lemma 3.2, we have

$$\|r(u^k, \beta_{\max})\| \rightarrow 0.$$

From $\lim_{k \rightarrow \infty} \eta_k = 0$, we know that there exist $k_0 > 0$ and $c_0 < 1$ such that

$$\left(1 - \frac{\delta\alpha}{\beta_{\max}}\right)^{\frac{1}{2}}(1 + \eta_k) \leq c_0, \quad \forall k \geq k_0, \tag{4.7}$$

which combined with (4.1) gives

$$\begin{aligned} \|r(u^{k+1}, \beta_{k+2})\| &\leq (1 + \eta_{k+1}) \left(1 - \frac{\delta\alpha}{\beta_{\max}}\right)^{\frac{1}{2}} \|r(u^k, \beta_{k+1})\| \\ &\leq c_0 \|r(u^k, \beta_{k+1})\| \quad \forall k \geq k_0. \end{aligned} \tag{4.8}$$

Note that $r(u^k, \beta_{k+1}) = u^k - u^{k+1}$, and from (4.8) we get

$$\|u^{k+2} - u^{k+1}\| \leq c_0 \|u^{k+1} - u^k\|, \quad \forall k \geq k_0.$$

Therefore, $\{u^k\}$ is a Cauchy sequence and converges to its cluster point, say u' . Because $\lim_{k \rightarrow \infty} u^k = u'$ and $r(u, \beta_{\max})$ is continuous on Ω , it follows that

$$r(u', \beta_{\max}) = \lim_{k \rightarrow \infty} r(u^k, \beta_{\max}) = 0,$$

and u' is a solution of (1.2). Since the problem has unique solution, then we have $u' = u^*$. Therefore, the generated sequence converges to the unique solution u^* . \square

5. Numerical Results

To verify the theoretical assertions, we consider the following least distance problem:

$$\begin{aligned} \min \quad & \frac{1}{2} \|x - c\|^2 \\ \text{s.t.} \quad & Ax \in \Omega \end{aligned}$$

where $A \in \mathbb{R}^{n \times n}$, $c \in \mathbb{R}^n$ and $\Omega \subset \mathbb{R}^n$ is a closed convex set. This problem can be written as

$$\begin{aligned} \min \quad & \frac{1}{2} \|x - c\|^2 \\ \text{s.t.} \quad & Ax - \xi = 0, \quad \xi \in \Omega. \end{aligned} \tag{5.1}$$

The Lagrangian function of problem (5.1) is

$$L(x, \xi, y) = \frac{1}{2} \langle x, x \rangle - \langle c, x \rangle - \langle y, Ax - \xi \rangle.$$

Thus, we have

$$L(x^*, \xi^*, y) \leq L(x^*, \xi^*, y^*) \leq L(x, \xi, y^*),$$

where $(x^*, \xi^*, y^*) \in \mathbb{R}^n \times \Omega \times \mathbb{R}^n$ is saddle point of the Lagrangian function. From the above inequalities for all $\xi \in \Omega$ we can obtain

$$\begin{cases} x^* = \langle A, y^* \rangle + c \\ \langle \xi - \xi^*, y^* \rangle \geq 0 \\ Ax^* = \xi^* \end{cases} \tag{5.2}$$

Substituting the first and the third equation in the second of system (5.2), we get

$$(AA^T y^* + Ac) \in \Omega, \quad \langle \xi - (AA^T y^* + Ac), y^* \rangle \geq 0 \quad \forall \xi \in \Omega, \tag{5.3}$$

which is exactly the general variational inequality (1.2) with

$$Q(u) = AA^T u + Ac \quad \text{and} \quad Ty = y.$$

We form the test problem as follows: The matrix A is $n \times n$ matrix whose entries are randomly generated in the interval $(-5,+5)$, the vector c is generated from a uniform distribution in the interval $(-500,500)$ and the closed convex set Ω is defined as

$$\Omega := \{u \in \mathbb{R}^n \mid \|u\| \leq a\}.$$

Then the projection on Ω in the sense of Euclidean-norm is very easy to carry out, for example,

$$\forall u \in \mathbb{R}^n, P_\Omega[u] = \begin{cases} u, & \text{if } \|u\| \leq a; \\ \frac{a}{\|u\|}u, & \text{if } \|u\| > a. \end{cases}$$

Note that in the case $\|Ac\| > a, \|AA^T u^* + Ac\| = a$ (otherwise $u^* = 0$ is the trivial solution). Therefore, we test the problem with $a = \rho \|Ac\|$ and $\rho \in (0, 1)$.

In all tests we take $\delta_0 = 0.5; \delta = 0.2$ and $\mu = 0.5$. All iterations start with $u^0 = (1, \dots, 1)^T$, $\beta_0 = 1$ and stopped whenever $\frac{\|r(u,1)\|}{a} \leq 10^{-5}$. The adjustment factor η_k for these test problems is adopted based in the criterion below,

$$\eta_{k+1} = \begin{cases} 1 & \text{if } u^{k+1} \text{ satisfies condition of Step 2;} \\ 0 & \text{otherwise,} \end{cases}$$

All codes were written in Matlab. The test results for problems (5.3) are reported in tables 1-2, k is the number of iterations and l denotes the number of evaluations of mapping Q .

Table 1. Numerical results for problem (5.3) with $n = 100$

ρ	Method [15]		Algorithm 3.1	
	k	l	k	l
0.1	58	185	51	79
0.3	13	53	13	26
0.5	8	45	7	19
0.7	5	33	5	16
0.9	3	27	4	15

Table 2. Numerical results for problem (5.3) with $n = 500$

ρ	Method [15]		Algorithm 3.1	
	k	l	k	l
0.1	39	103	39	52
0.3	9	43	9	22
0.5	5	35	5	18
0.7	4	33	4	17
0.9	3	31	3	16

6. Remarks

It is interesting to observe the difference of the convergent between Goldstein's projection method [6] and our method. We remark that the Goldstein's projection method updates u^{k+1} according to the formula

$$u^{k+1} = P_{\Omega}[u^k - \beta F(u^k)].$$

We note that if F is Lipschitz continuous (with a Lipschitz $L > 0$) and uniformly strong monotone (with a constant modulus α)

$$(u - v)^T (F(u) - F(v)) \geq \alpha \|u - v\|^2,$$

and β satisfies

$$0 < \beta < \frac{2\alpha}{L^2},$$

then this method is convergent. However, the estimation of α and L may lead to a slow convergence. The performance of our method for $T = I$, the identity operator, needs neither the Lipschitz L nor the strong monotone modulus α to design the algorithm, but only depends on the choice of scaling parameter β . Our preliminary numerical results show that the method may be efficient for some large scale problems. Moreover, it demonstrates computationally that the new method needs less evaluations of mapping Q . It is evident that the smaller the ratio $a/\|AC\|$ is, the more iterations it takes to meet the stopping criterion. On the other hand, it seems that the number of evaluations of mapping Q as well as the overall number of iterations is not very sensitive to the size of the problem, that is the dimension of the space \mathbb{R}^n .

Acknowledgements. This research was supported partly by NSFC Grants Nos: 70731002 and 70571034, "The Second Term '985' Engineering-Innovation Center for Computational Laboratory of Evolutionary Economic Theory and Its Applications" and "Study on the Evolution of Complex Economic System" at "Innovation Center of Economic Transition and Development of Nanjing University" of Ministry of Education, China. The research of Prof. M. Aslam Noor was supported by the Visiting Professor Program of King Saud University, Riyadh, Saudi Arabia and Research Grant No: KSU.VPP.108. Prof. M. Aslam Noor is grateful to Dr. S. M. Junaid Zaidi, Rector, CIIT, Islamabad, for providing excellent research facilities. The authors would like to thank the referee for his very constructive suggestions and comments.

REFERENCES

- [1] A. BNOUHACHEM, *A self-adaptive method for solving general mixed variational inequalities*, J. Math. Anal. Appl., **309** (2005), 136–150.
- [2] A. BNOUHACHEM, *A new projection and contraction method for linear variational inequalities*, J. Math. Anal. Appl., **314** (2006), 513–525.
- [3] R. E. BRUCK, *An iterative solution of a variational inequality for certain monotone operators in Hilbert space*, Bull. Amer. Math. Soc., **81** (1975), 890–892.
- [4] B. C. EAVES, *On the basic theorem of complementarity*, Mathematical Programming, **1** (1971), 68–75.
- [5] M. FUKUSHIMA, *A relaxed projection method for variational inequalities*, Mathematical Programming, **35** (1986), 58–70.
- [6] A. A. GOLDSTEIN, *Convex programming in Hilbert space*, Bul. Amer. Math. Soc., **70** (1964), 709–710.
- [7] P. T. HARKER AND J. S. PANG, *Finite-dimensional variational inequality and nonlinear complementarity problems: A survey of theory, algorithms and applications*, Mathematical Programming, **48** (1990), 161–220.
- [8] B. S. HE, *A modified projection and contraction method for a class of linear complementarity problems*, J. Computational Math., **14** (1996), 54–63.
- [9] B. S. HE, *Inexact implicit methods for monotone general variational inequalities*, Mathematical Programming, **86** (1999), 199–216.
- [10] B. S. HE, H. YANG, Q. MENG AND D. R. HAN, *Modified Goldstein-Levitin-Polyak projection method for asymmetric strongly monotone variational inequalities*, J. Optim. Theory Appl., **112** (2002), 129–143.
- [11] B. S. HE, AND L. Z. LIAO, *Improvement of some projection methods for monotone variational inequalities*, J. Optim. Theory Appl., **112** (2002), 111–128.
- [12] A. N. IUSEM AND B. F. SVAITER, *A variant of Korplevich's method for variational inequalities with a new search strategy*, Optimization, **42** (1997), 309–321.
- [13] E. N. KHOBOTOV, *Modification of the extragradient method for solving variational inequalities and certain optimization problems*, USSR Computational Mathematics and Mathematical Physics, **27** (1987), 120–127.
- [14] G. M. KORPLEVICH, *The extragradient method for finding saddle points and other problems*, Ekonomika i Matematicheskie Metody, **12** (1976), 747–756.
- [15] M. LI AND X. M. YUAN, *An improved Goldstein's type method for a class of variant variational inequalities*, J. Computational and Applied Math., **214** (2008), 304–312.
- [16] J. L. LIONS AND G. STAMPACCHIA, *Variational inequalities*, Comm. Pure Appl. Math., **20** (1967), 493–512.
- [17] J. L. LIONS AND B. MERCIER, *Splitting algorithms for the sum of two nonlinear operators*, SIAM J. Numerical Analysis, **16** (1979), 964–979.
- [18] M. A. NOOR, *General variational inequalities*, Appl. Math. Letters, **1** (1988), 119–121.
- [19] M. A. NOOR, *Quasi variational inequalities*, Appl. Math. Letters, **1** (1988), 267–270.
- [20] M. A. NOOR, *New extragradient-type methods for general variational inequalities*, J. Math. Anal. Appl., **277** (2003), 379–395.
- [21] M. A. NOOR, *Some developments in general variational inequalities*, Appl. Math. Computation, **152** (2004), 199–277.
- [22] M. A. NOOR, *Principles of Variational Inequalities*, Lap-Lambert Academic Publishing AG & Co, Saarbrücken, Germany, 2009.
- [23] M. A. NOOR, *On a class of general variational inequalities*, Journal of Advanced Mathematical studies, **1** (2009), 75–86.
- [24] M. A. NOOR, *An extragradient algorithm for solving general nonconvex variational inequalities*, Applied mathematics Letters, **23** (2010), 917–921.
- [25] M. A. NOOR, *On an implicit method for nonconvex variational inequalities*, Journal of Optimization Theory and Applications, **147** (2010), 411–417.
- [26] M. A. NOOR, *Some classes of general nonconvex variational inequalities*, Albanian Journal of Mathematics, **3** (2009), 175–188.

- [27] M. A. NOOR AND K. I. NOOR, *Self-adaptive projection algorithms for general variational inequalities*, Appl. Math. Computation, **151** (2004), 659–670.
- [28] M. A. NOOR, K. I. NOOR AND T. M. RASSIAS, *Some aspects of variational inequalities*, J. Comput. Appl. Math., **47** (1993), 285–312.
- [29] M. A. NOOR AND T. M. RASSIAS, *A class of projection methods for general variational inequalities*, J. Math. Anal. Appl., **268** (2002), 334–343.
- [30] J. V. OUTFRATA AND J. ZOWE, *A Newton method for a class of quasi-variational inequalities*, Computational Optimization and Applications, **4** (1995), 5–21.
- [31] J. S. PANG AND L. QI, *Non smooth equations: motivation and algorithms*, SIAM J. Optimization, **3** (1993), 443–465.
- [32] J. M. PENG AND M. FUKUSHIMA, *A hybrid Newton method for solving the variational inequality problem via the D-gap function*, Mathematical Programming, **86**, 2 (1999), 367–386.
- [33] M. V. SOLODOV AND P. TSENG, *Modified projection-type methods for monotone variational inequalities*, SIAM J. Control and Optim., **34** (1996), 1814–1830.
- [34] M. V. SOLODOV AND B. F. SVAITER, *A new projection method for variational inequality problems*, SIAM J. Control and Optim., **37** (1999), 765–776.
- [35] N. XIU, J. ZHANG AND M. A. NOOR, *Tangent projection equations and general variational inequalities*, J. Math. Anal. Appl., **258** (2001), 755–762.

(Received September 26, 2008)

Abdellah Bnouhachem
 School of Management Science and Engineering
 Nanjing University
 Nanjing, 210093
 P.R. China
 and
 Ibn Zohr University
 ENSA, BP:1136
 Agadir, Morocco

Muhammad Aslam Noor
 Mathematics Department
 COMSATS Institute of Information Technology
 Islamabad, Pakistan
 and
 Mathematics Department, College of Science
 King Saud University
 Riyadh, Saudi Arabia
 e-mail: noormaslam@hotmail.com

Mohamed Khalfouï
 Ecole Supérieure de Technologie de Salé
 Mohamed 5 University of Science
 Agdal-Rabat, Morocco

Sheng Zhaohan
 School of Management Science and Engineering
 Nanjing University
 Nanjing, 210093
 P.R. China