

UPPER BOUNDS ON MULTIPLE GENERALIZED MATHIEU SERIES

ŽIVORAD TOMOVSKI, DELČO LEŠKOVSKI AND TIBOR K. POGÁNY

(Communicated by A. Čížmešija)

Abstract. The main aim of this short note is to obtain new very general upper bounds for multiple generalized Mathieu series considering the related integral representation obtained recently by Pogány and Tomovski [10], by means of the multiple Hardy–Hilbert type integral inequality given in [1].

1. Introduction and preparation

The series

$$S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} \quad r > 0,$$

was introduced by Émile Leonard Mathieu in 1890 in his book [7] devoted to mathematical physics investigations of the elasticity of rigid bodies. Several extensions and unifications such as generalized Mathieu, \mathbf{a} –, $(\mathbf{a}, \boldsymbol{\lambda})$ –series and their alternating variants have been considered in getting integral representation results are obtained recently by Cerone, Pogány, Srivastava, Tomovski and their coworkers [2, 3, 8, 9, 12, 13, 14, 17], and related bounding inequalities [4, 10, 11, 12, 14, 15, 16]. However, to the best knowledge of the authors the multiple Mathieu series has been considered only in two publications, in ones by Pogány and Tomovski [10] and by Draščić Ban [4], where Draščić Ban focused to the so-called multiple Mathieu’s $(\mathbf{a}, \boldsymbol{\lambda})$ –series. In both papers certain sharp bounding inequalities have been obtained in the sense that the well-known sharp bilateral bound $0 \leq \{u\} < 1$ was employed, where $\{u\}$ stands for the fractional part of some real u , see [10] and [4], respectively.

We consider here the *multiple generalized Mathieu series* [10, Eq. (2)] defined by

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) = \sum_{\mathbf{n} \in \mathbb{N}^m} \frac{2\mathbf{n}^{|\mathbf{s}|}}{\langle \mathbf{n}^{\mathbf{v}}, \mathbf{n}^{\mathbf{v}} \rangle + r^2)^{p+1}},$$

where $\mathbf{n}^{\mathbf{v}} := (n_1^{v_1}, \dots, n_m^{v_m})$; $\mathbf{n}^{\boldsymbol{\alpha}|\mathbf{s}|} := n_1^{\alpha_1 s_1} \dots n_m^{\alpha_m s_m}$; \mathbf{s}, \mathbf{v} have positive coordinates, i.e. $s_\ell, v_\ell > 0$, $\ell = \overline{1, m}$ and $\langle \mathbf{a}, \mathbf{b} \rangle$ stands for the inner product in \mathbb{R}_+^m .

Mathematics subject classification (2010): Primary: 26D15, 40B05; secondary: 26B40, 33E20.

Keywords and phrases: Multiple generalized Mathieu series, generalized Hardy–Hilbert integral inequality, non-conjugated Hölder parameters.

Let us introduce few necessary notations and conventions. Denote here, and in what follows, $[x]$ the integer part of some $x \in \mathbb{R}$ and $(a)_m := a(a+1)\cdots(a+m-1)$, $a \in \mathbb{C}$, $m \in \mathbb{N} = \{1, 2, \dots\}$ the Pochhammer symbol (or *shifted factorial*).

The following integral representation formula has been derived by Pogány and Tomovski [10].

LEMMA 1. [10, Theorem 1] *Let $r, p+1 > 0$, let the multiindices \mathbf{s}, \mathbf{v} be positive and taken so, that the multiple generalized Mathieu series $\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v})$ converges. Then we have*

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) = (p+1)_m \int_{[1, \infty)^m} \frac{\prod_{j=1}^m \left(\frac{[t_j^{1/(2v_j)}]^{s_j+1}}{s_j+1} + s_j \int_0^{[t_j^{1/(2v_j)}]} \{u\} u^{s_j-1} du \right)}{(r^2 + t_1 + \dots + t_m)^{p+m+1}} d\mathbf{t}, \tag{1}$$

where $d\mathbf{t} := dt_1 \cdots dt_m$.

Let $p_j > 1$, $j = 1, \dots, m$ be parameters such that $\sum_{j=1}^m p_j^{-1} \geq 1$, and let p'_j be the conjugated Hölder parameter related to p_j , that is $p_j^{-1} + (p'_j)^{-1} = 1$, $j = 1, \dots, m$. By the standard notation

$$\lambda = \frac{1}{m-1} \sum_{j=1}^m \frac{1}{p'_j} = \sum_{j=1}^m \frac{1}{q_j},$$

where $q_j^{-1} := \lambda - (p'_j)^{-1}$, $j = 1, \dots, m$. Obviously $0 < \lambda \leq 1$ and for $\lambda = 1$ the involved parameters correspond to the conjugated case.

We will also need the following, *here necessarily precised*, Hardy–Hilbert type inequality result by Brnetić *et al.* [1], achieved for the non–conjugated Hölder parameters case.

LEMMA 2. [1, Theorem 1, Eq. (7)] *Let $m \geq 2$ be an integer, $\lambda, p_j, p'_j; q_j > 0$, $j = 1, \dots, m$ be the real numbers previously defined. Let the domain of integration $\Omega \subseteq \mathbb{R}_+$, and assume that the kernel function $K: \Omega^m \mapsto \mathbb{R}_+$ and $\mu_j, j = 1, \dots, m$ are nonnegative, σ –finite measures. If the weight functions $\phi_{ij} \geq 0$, $i, j = 1, \dots, m$ satisfy condition $\prod_{i,j=1}^m \phi_{ij}(x_j) = 1$, then the following inequality holds*

$$\int_{\Omega^m} K^\lambda(\mathbf{x}) \prod_{j=1}^m f_j(x_j) d\mu_j(x_j) \leq \prod_{j=1}^m \left(\int_{\Omega} \left(\phi_{jj}(x_j) F_j(x_j) f_j(x_j) \right)^{p_j} d\mu_j(x_j) \right)^{1/p_j} \tag{2}$$

where $f_j: \Omega \mapsto \mathbb{R}_+$, $j = 1, \dots, m$; and $\mathbf{x} := (x_1, \dots, x_m)$, while

$$F_i(x_i) = \left(\int_{\Omega^{m-1}} K(\mathbf{x}) \prod_{\substack{j=1 \\ j \neq i}}^m \phi_{ij}^{q_i}(x_j) d\mu_j(x_j) \right)^{1/q_i};$$

provided both sides of (2) there exist.

REMARK. For the sake of completeness we have to mention that Hardy–Hilbert integral inequalities play significant roles in deriving bounding inequalities for alternating Mathieu–type series [16]; they were also important tool in estimating generalized hypergeometric functions ${}_mF_n$ [6].

2. Main results

Our aim is to apply Lemma 2 to the integral representation (1) to derive new upper bounds for $\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v})$. Then, suitably specifying the kernel function K we derive further more elegant corollaries of the main theorem.

THEOREM 1. Let $m \geq 2$ be an integer, $\lambda, p_j, p'_j; q_j > 0, j = 1, \dots, m$ be the real numbers defined in the preambula of the Lemma 2, and be $A_{ij} + q_i^{-1} > 0, i \neq j; i, j = 1, \dots, m$ such, that $\sum_{i=1}^m A_{ij} = 0$ and

$$A_{ii} - \alpha_i > \frac{(m - 1)\lambda - m - p - 1}{\lambda q_i} \quad \alpha_i := \sum_{j=1}^m A_{ij}, \quad i = 1, \dots, m.$$

Then the following inequality holds

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) \leq C \cdot \prod_{i=1}^m \left(\int_{\mathbb{R}_+} x_i^{A_{ii} p_i} (r^2 + m + x_i)^{\frac{p_i}{\lambda q_i} ((m-1)\lambda - m - p - 1) + p_i(\alpha_i - A_{ii})} f_i^{p_i}(x_i) dx_i \right)^{1/p_i}, \tag{3}$$

where

$$C = \frac{2(p+1)_m}{\Gamma^\lambda \left(\frac{p+m+1}{\lambda} \right)} \prod_{i=1}^m \Gamma^{1/q_i} \left(\frac{p+m+1}{\lambda} - m + 1 - q_i(\alpha_i - A_{ii}) \right) \cdot \prod_{\substack{i,j=1 \\ i \neq j}}^m \Gamma^{1/q_i} (q_i A_{ij} + 1),$$

and

$$f_j(x_j) = \frac{[(x_j + 1)^{1/(2v_j)}]^{s_j+1}}{s_j + 1} + s_j \int_0^{[(x_j+1)^{1/(2v_j)}]} \{u\} u^{s_j-1} du, \quad j = 1, \dots, m. \tag{4}$$

Proof. First, by substituting $x_j + 1 \mapsto t_j, j = 1, \dots, m$ the integral formula (1) becomes

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) = 2(p + 1)_m \int_{\mathbb{R}_+^m} \frac{\prod_{j=1}^m f_j(x_j)}{(r^2 + m + x_1 + \dots + x_m)^{p+m+1}} d\mathbf{x}, \tag{5}$$

where $f_j(x_j)$ is defined in (4). Choose the kernel function

$$K(\mathbf{x}) = (r^2 + m + x_1 + \dots + x_m)^{-(p+m+1)/\lambda},$$

and define $\phi_{ij}(x_j) = x_j^{A_{ij}}$ on $\Omega := \mathbb{R}_+$. Then the normalizing condition of Lemma 2 is satisfied. Indeed

$$\prod_{i,j=1}^m \phi_{ij}(x_j) = \prod_{i,j=1}^m x_j^{A_{ij}} = \prod_{j=1}^m x_j^{\sum_{i=1}^m A_{ij}} = 1,$$

by virtue of the assumed $\sum_{i=1}^m A_{ij} = 0, j = 1, \dots, m$. Now, we conclude

$$F_i(x_i) = \left(\int_{\mathbb{R}_+^{m-1}} K(\mathbf{x}) \prod_{\substack{j=1 \\ j \neq i}}^m x_j^{q_i A_{ij}} \cdot \frac{d\mathbf{x}}{dx_i} \right)^{1/q_i}.$$

By substitution $x_k(r^2 + m + x_i)^{-1} \mapsto u_k, k = 1, \dots, m, k \neq i$, we get

$$F_i(x_i) = (r^2 + m + x_i)^{\sum_{j=1, j \neq i}^m A_{ij} + \frac{m-1}{q_i} - \frac{p+m+1}{\lambda q_i}} \left(\int_{\mathbb{R}_+^{m-1}} \frac{\prod_{\substack{j=1 \\ j \neq i}}^m u_j^{q_i A_{ij}} \cdot \frac{du}{du_i}}{\left(1 + \sum_{\substack{k=1 \\ k \neq i}}^m u_k\right)^{(p+m+1)/\lambda}} \right)^{1/q_i}.$$

Employing the familiar formula for the Gamma–function integral [18, Lemma 5.1]

$$\int_{\mathbb{R}_+^{k-1}} \frac{\prod_{j=1}^{k-1} z_j^{a_j-1}}{\left(1 + \sum_{j=1}^{k-1} z_j\right)^{\sum_{j=1}^k a_j}} d\mathbf{z} = \frac{\prod_{j=1}^k \Gamma(a_j)}{\Gamma\left(\sum_{j=1}^k a_j\right)} \quad \min_{1 \leq j \leq k} \Re\{a_j\} > 0,$$

we deduce

$$F_i^{q_i}(x_i) = \frac{(r^2 + m + x_i)^{\left(m-1 - \frac{p+m+1}{\lambda}\right) + q_i(\alpha_i - A_{ii})} \prod_{\substack{j=1 \\ j \neq i}}^m \Gamma(q_i A_{ij} + 1)}{\Gamma\left(\frac{p+m+1}{\lambda}\right)} \times \Gamma\left(\frac{p+m+1}{\lambda} - m + 1 - q_i(\alpha_i - A_{ii})\right),$$

such that ensures the asserted upper bound (3). \square

Choosing $A_{ij} = 0, i, j = 1, \dots, m$ in the Theorem 1 we get the following result.

COROLLARY 1. *Let $m \geq 2$ be an integer, $\lambda, p_j, p'_j; q_j > 0, j = 1, \dots, m$ be the real numbers defined in Theorem 1. Then the following inequality holds*

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) \leq C_1 \cdot \prod_{i=1}^m \left(\int_{\mathbb{R}_+} (r^2 + m + x_i)^{\frac{p_i}{q_i} \left(m-1 - \frac{m-p-1}{\lambda}\right)} f_i^{p_i}(x_i) dx_i \right)^{1/p_i},$$

where

$$C_1 = \frac{2(p+1)_m}{\left(\frac{p+m+1}{\lambda} - m + 1\right)_{m-1}^\lambda},$$

and functions $f_i(x_i)$ are given by (4).

Now, if we put $A_{ii} = A_i$, $A_{i,i+1} = -A_{i+1}$ while $A_{ij} = 0$, $|i - j| > 1$ and the indices are taken modulo m , then we obtain the next result.

COROLLARY 2. *Let $m \geq 2$ be an integer, λ , p_j , p'_j ; $q_j > 0$, $j = 1, \dots, m$ be the real numbers defined in the Theorem 1. Then, for all*

$$p + (1 - \lambda)m > 2\lambda - 1, \quad A_i \in \left(m - \frac{p+m+1}{\lambda} - \frac{1}{q_i}, \frac{1}{q_i}\right)$$

and $A_{m,m+1} = -A_1$, the following inequality holds

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) \leq C_2 \prod_{i=1}^m \left(\int_{\mathbb{R}_+} x_i^{A_i p_i} (r^2 + m + x_i)^{\frac{p_i}{q_i} (m-1 - \frac{m+p+1}{\lambda}) - p_i A_{i+1}} f_i^{p_i}(x_i) dx_i \right)^{1/p_i},$$

where

$$C_2 = \frac{2(p+1)_m}{\Gamma^\lambda \left(\frac{p+m+1}{\lambda}\right)} \prod_{i=1}^m \left(\Gamma(1 - q_i A_{i+1}) \Gamma\left(\frac{p+m+1}{\lambda} - m + 1 + q_i A_{i+1}\right) \right)^{1/q_i},$$

and functions $f_i(x_i)$ remain the same as in (4).

Finally, we give another variant of upper bounds choosing weight functions ϕ_{ij} to be exponential instead of the power weight function set such that results in Theorem 1.

THEOREM 2. *Let $m \geq 2$, and λ , p_j , p'_j ; $q_j > 0$, $j = 1, \dots, m$ satisfy the non-conjugated Hölder parameters constraints. Assume that $A_{ij} \in \mathbb{R}$, $i, j = 1, \dots, m$ such that $\sum_{i=1}^m A_{ij} = 0$ and*

$$0 < q_1 A_{i1} < q_2 A_{i2} < \dots < q_m A_{im}, \quad i = 1, \dots, m.$$

Then the following inequality holds true:

$$\mathcal{S}_p^m(r, \mathbf{s}, \mathbf{v}) \leq 2(p+1)_m \cdot \prod_{k=1}^m \left(\int_{\mathbb{R}_+} e^{-A_{kk} p_k x_k} F_k^{p_k}(x_k) f_k^{p_k}(x_k) dx_k \right)^{1/p_k}, \quad (6)$$

where

$$F_k(x_k) = \left(\frac{\sum_{\substack{j=1 \\ j \neq k}}^m \frac{(q_j A_{kj})^{\frac{p+m+1-1}{\lambda}}}{\prod_{\substack{n=1 \\ n \neq j,k}}^m (q_j A_{kj} - q_n A_{kn})} e^{(r^2+m+x_k)q_j A_{kj}}}{\prod_{\substack{n=1 \\ n \neq j,k}}^m (q_j A_{kj} - q_n A_{kn})} \Gamma\left(1 - \frac{p+m+1}{\lambda}, (r^2+m+x_k)q_j A_{kj}\right) \right)^{1/q_k},$$

and $f_k(x_k)$, $k = 1, \dots, m$ remain the same (4).

Proof. Let us begin the proving procedure in the same manner as for Theorem 1, that is, considering integral expression (5). The kernel function remains also the same.

Now, define $\phi_{kj}(x_j) = e^{-A_{kj}x_j}$, $k, j = 1, \dots, m$ on $\Omega := \mathbb{R}_+$. The normalizing constraint of the Lemma 2 is obviously satisfied since $\sum_{i=1}^m A_{ki} = 0$, $j = 1, \dots, m$. Therefore we have

$$\begin{aligned} F_k^{qk}(x_k) &= \int_{\mathbb{R}_+^{m-1}} K(\mathbf{x}) \exp \left\{ - \sum_{\substack{j=1 \\ j \neq k}}^m q_k A_{kj} x_j \right\} \cdot \frac{d\mathbf{x}}{dx_k} \\ &= \frac{1}{\Gamma\left(\frac{p+m+1}{\lambda}\right)} \int_{\mathbb{R}_+} t^{\frac{p+m+1}{\lambda}-1} e^{-(r^2+m+x_k)t} \left(\int_{\mathbb{R}_+^{m-1}} e^{-\sum_{\substack{j=1 \\ j \neq k}}^m (t+q_k A_{kj})x_j} \frac{d\mathbf{x}}{dx_k} \right) dt \\ &= \frac{1}{\Gamma\left(\frac{p+m+1}{\lambda}\right)} \int_{\mathbb{R}_+} \frac{t^{\frac{p+m+1}{\lambda}-1}}{\prod_{\substack{j=1 \\ j \neq k}}^m (t + q_k A_{kj})} e^{-(r^2+m+x_k)t} dt. \end{aligned}$$

The partial fraction decomposition of the rational part of the integrand gives

$$F_k^{qk}(x_k) = \frac{1}{\Gamma\left(\frac{p+m+1}{\lambda}\right)} \sum_{\substack{j=1 \\ j \neq k}}^m \frac{\int_{\mathbb{R}_+} \frac{t^{\frac{p+m+1}{\lambda}-1}}{t + q_j A_{kj}} e^{-(r^2+m+x_k)t} dt}{\prod_{\substack{n=1 \\ n \neq j,k}}^m (q_n A_{kn} - q_j A_{kj})}.$$

Employing the formula [5, 3.383 10.]

$$\begin{aligned} \int_{\mathbb{R}_+} \frac{x^{v-1} e^{-\mu x}}{x + \beta} dx &= \beta^{v-1} e^{\beta \mu} \Gamma(v) \Gamma(1-v, \beta \mu), \\ |\arg(\beta)| < \pi, \quad \min(\Re\{\mu\}, \Re\{v\}) > 0, \end{aligned}$$

where $\Gamma(s, z) = \int_z^\infty x^{s-1} e^{-sx} dx$ is the familiar upper incomplete Gamma function, we deduce

$$F_k^{qk}(x_k) = \sum_{\substack{j=1 \\ j \neq k}}^m \frac{(q_j A_{kj})^{\frac{p+m+1}{\lambda}-1} e^{(r^2+m+x_k)q_j A_{kj}}}{\prod_{\substack{n=1 \\ n \neq j,k}}^m (q_n A_{kn} - q_j A_{kj})} \Gamma\left(1 - \frac{p+m+1}{\lambda}, (r^2+m+x_k)q_j A_{kj}\right).$$

By virtue of Lemma 2, taking $\mu_k, k = 1, \dots, m$ to be ordinary Lebesgue measures, we immediately arrive at the asserted inequality (6). \square

REFERENCES

- [1] I. BRNETIĆ, M. KRNIĆ M. AND J. PEČARIĆ, *Multiple Hilbert and Hardy–Hilbert inequalities with non–conjugated parameters*, Bull. Austral. Math. Soc. **71** (2005), 447–457.
- [2] P. CERONE AND C.T. LENARD, *On integral forms of generalized Mathieu series*, JIPAM J. Inequal. Pure Appl. Math. **4**, 5 (2003), Art. 100.
- [3] B. DRAŠČIĆ AND T.K. POGÁNY, *On integral representation of the first kind Bessel function*, J. Math. Anal. Appl. **308**, 2 (2005), 775–780.
- [4] B. DRAŠČIĆ BAN, *On multiple Mathieu $(\mathbf{a}, \boldsymbol{\lambda})$ –series*, Miskolc Math. Notes **11**, 1 (2010), 53–62.
- [5] I.S. GRADSHTEIN AND I.M. RYZHIK, *Table of Integrals, Series and Products*, 4.Ed. supervised by YU. V. GERONIMUS AND M. YU. TSEITLIN, Fizmatgiz, Moscow, 1963. (In Russian)
- [6] M. KRNIĆ, Ž. TOMOVSKI AND J. PEČARIĆ, *Hilbert inequalities related to generalized hypergeometric functions ${}_mF_n$* , Math. Balkanica (N.S.) **22**, 3–4 (2008), 307–322.
- [7] É.L. MATHIEU, *Traité de Physique Mathématique VI–VII: Théorie de l'élasticité des corps solides*, Gauthier-Villars, Paris, 1890.
- [8] T.K. POGÁNY, *Integral representation of a series which includes the Mathieu \mathbf{a} -series*, J. Math. Anal. Appl. **296**, 1 (2004), 309–313.
- [9] T.K. POGÁNY, *Integral representation of Mathieu $(\mathbf{a}, \boldsymbol{\lambda})$ -series*, Integral Transform Spec. Funct. **16**, 8 (2005), 685–689.
- [10] T.K. POGÁNY AND Ž. TOMOVSKI, *On multiple generalized Mathieu series*, Integral Transforms Spec. Funct. **17**, 4 (2006), 285–293.
- [11] T.K. POGÁNY AND Ž. TOMOVSKI, *Bounds improvement for alternating Mathieu type series*, J. Math. Inequal. **4**, 3 (2010), 315–324.
- [12] T.K. POGÁNY, H.M. SRIVASTAVA AND Ž. TOMOVSKI, *Some families of Mathieu \mathbf{a} -series and alternating Mathieu \mathbf{a} -series*, Appl. Math. Comput. **173**, 1 (2006), 69–108.
- [13] H.M. SRIVASTAVA AND Ž. TOMOVSKI, *Some problems and solutions involving Mathieu's series and its generalizations*, JIPAM J. Inequal. Pure Appl. Math. **5**, 2 (2004), Art.45.
- [14] Ž. TOMOVSKI, *New double inequalities for Mathieu type series*, Publikacije Elektrotehn. Fak. Ser. Mat. Univ. Beograd **15** (2004), 79–83.
- [15] Ž. TOMOVSKI AND D. LEŠKOVSKI, *Refinements and sharpness of some inequalities for Mathieu type series*, Math. Maced. **6** (2008), 61–71.
- [16] Ž. TOMOVSKI AND T.K. POGÁNY, *New upper bounds for Mathieu type series*, Banach J. Math. Anal. **3**, 2 (2008), 9–15.
- [17] Ž. TOMOVSKI AND K. TRENČEVSKI, *On an open problem of Bai-Ni Guo and Feng Qi*, JIPAM J. Inequal. Pure Appl. Math. **4**, 2 (2003), Art. 29.
- [18] B.–C. YANG AND T.M. RASSIAS, *On the way of weight coefficients and research for the Hilbert–type inequalities*, Math. Inequal. Appl. **6** (2003), 625–658.

(Received February 16, 2011)

Živorad Tomovski
Institute of Mathematics
St. Cyril and Methodius University
MK-1000 Skopje, Macedonia
e-mail: zivoradt@yahoo.com

Delčo Leškovski
Faculty of Technical Sciences
International Balkan University
MK-1000 Skopje, Macedonia
e-mail: dleskovski@gmail.com

Tibor K. Pogány
Faculty of Maritime Studies
University of Rijeka
HR-51000 Rijeka, Croatia
e-mail: poganj@pfri.hr