

BOUNDEDNESS ON MORREY SPACE FOR TOEPLITZ TYPE OPERATOR ASSOCIATED TO SINGULAR INTEGRAL OPERATOR WITH VARIABLE CALDERÓN–ZYGmund KERNEL

CHUANGXIA HUANG, SHENG GUO AND LANZHE LIU

(Communicated by N. Elezović)

Abstract. In this paper, the boundedness of the Toeplitz type operators associated to the singular integral operator with variable Calderón–Zygmund kernel on Morrey spaces is obtained. For this purpose, some M^k -type sharp maximal function inequalities for the operators are proved.

1. Introduction and Preliminaries

As the development of singular integral operators (see [7], [21]), their commutators have been well studied. In [4], [19], [20], the authors proved that the commutators generated by the singular integral operators and BMO functions are bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$. Chanillo (see [2]) proved a similar result when singular integral operators are replaced by the fractional integral operators. In [1], Calderón and Zygmund introduced some singular integral operators with variable kernel and discussed their boundedness. In [11–13], [22], the authors obtained the boundedness for the commutators generated by the singular integral operators with variable kernel and BMO functions. In [15], the authors proved the boundedness for the multilinear oscillatory singular integral operators generated by the operators and BMO functions. In [8], [9], [14], some Toeplitz type operators associated to the singular integral operators and strongly singular integral operators are introduced, and the boundedness for the operators generated by BMO and Lipschitz functions is obtained. Motivated by these, in this paper, we will study the Toeplitz type operator generated by the singular integral operator with variable Calderón–Zygmund kernel.

First, let us introduce some notations. Throughout this paper, Q will denote a cube of \mathbb{R}^n with sides parallel to the axes. For any locally integrable function f , the sharp maximal function of f is defined by

$$f^\#(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y) - f_Q| dy,$$

Mathematics subject classification (2010): 42B20, 42B25.

Keywords and phrases: Toeplitz type operator, singular integral operator, variable Calderón–Zygmund kernel, Sharp maximal function, BMO , Morrey space.

where, and in what follows, $f_Q = |Q|^{-1} \int_Q f(x) dx$. It is well-known that (see [7], [21])

$$f^\#(x) \approx \sup_{Q \ni x} \inf_{c \in \mathbb{C}} \frac{1}{|Q|} \int_Q |f(y) - c| dy.$$

We say that f belongs to $BMO(\mathbb{R}^n)$ if $f^\#$ belongs to $L^\infty(\mathbb{R}^n)$ and define $\|f\|_{BMO} = \|f^\#\|_{L^\infty}$. It is known that (see [21])

$$\|f - f_{2^k Q}\|_{BMO} \leq Ck \|f\|_{BMO}.$$

For $0 < r < \infty$, we denote $f_r^\#$ by

$$f_r^\#(x) = [(|f|^r)^\#(x)]^{1/r}.$$

Let

$$M(f)(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)| dy.$$

For $\eta > 0$, let $M_\eta(f) = M(|f|^\eta)^{1/\eta}$. For $k \in \mathbb{N}$, we denote by M^k the operator M iterated k times, i.e., $M^1(f) = M(f)$ and

$$M^k(f) = M(M^{k-1}(f)) \text{ when } k \geq 2.$$

For $0 < \eta < n$ and $1 \leq r < \infty$, set

$$M_{\eta,r}(f)(x) = \sup_{Q \ni x} \left(\frac{1}{|Q|^{1-r\eta/n}} \int_Q |f(y)|^r dy \right)^{1/r}.$$

Let Φ be a Young function and $\tilde{\Phi}$ be the complementary associated to Φ , we denote the Φ -average for a function f by

$$\|f\|_{\Phi,Q} = \inf \left\{ \lambda > 0 : \frac{1}{|Q|} \int_Q \Phi \left(\frac{|f(y)|}{\lambda} \right) dy \leq 1 \right\}$$

and the maximal function associated to Φ by

$$M_\Phi(f)(x) = \sup_{x \in Q} \|f\|_{\Phi,Q}.$$

The Young functions which we use in this paper are $\Phi(t) = t(1 + \log t)$ and $\tilde{\Phi}(t) = \exp(t)$, the corresponding average and maximal functions are denoted by $\|\cdot\|_{L(\log L),Q}$, $M_{L(\log L)}$ and $\|\cdot\|_{\exp L,Q}$, $M_{\exp L}$. Following [19], we know the generalized Hölder's inequality and the following inequalities hold:

$$\frac{1}{|Q|} \int_Q |f(y)g(y)| dy \leq \|f\|_{\Phi,Q} \|g\|_{\tilde{\Phi},Q},$$

$$\|f\|_{L(\log L),Q} \leq M_{L(\log L)}(f) \leq CM^2(f),$$

$$\|f - f_{2jQ}\|_{\text{exp}L,Q} \leq Cj\|f\|_{BMO}.$$

DEFINITION 1. Let φ be a positive, increasing function on R^+ and there exists a constant $D > 0$ such that

$$\varphi(2t) \leq D\varphi(t) \text{ for } t \geq 0.$$

Let f be a locally integrable function on R^n . Set, for $1 \leq p < \infty$,

$$\|f\|_{L^{p,\varphi}} = \sup_{x \in R^n, d > 0} \left(\frac{1}{\varphi(d)} \int_{Q(x,d)} |f(y)|^p dy \right)^{1/p},$$

where $Q(x, d) = \{y \in R^n : |x - y| < d\}$. The generalized Morrey space is defined by

$$L^{p,\varphi}(R^n) = \{f \in L^1_{loc}(R^n) : \|f\|_{L^{p,\varphi}} < \infty\}.$$

If $\varphi(d) = d^\delta$, $\delta > 0$, then $L^{p,\varphi}(R^n) = L^{p,\delta}(R^n)$, which is the classical Morrey spaces (see [17], [18]). If $\varphi(d) = 1$, then $L^{p,\varphi}(R^n) = L^p(R^n)$, which is the Lebesgue spaces (see [7]).

As the Morrey space may be considered as an extension of the Lebesgue space, it is natural and important to study the boundedness of the operator on the Morrey spaces (see [5], [6], [10], [16]).

In this paper, we will study some singular integral operators as follows (see [1]).

DEFINITION 2. Let $K(x) = \Omega(x)/|x|^n : R^n \setminus \{0\} \rightarrow R$. K is said to be a Calderón-Zygmund kernel if

- (a) $\Omega \in C^\infty(R^n \setminus \{0\})$;
- (b) Ω is homogeneous of degree zero;
- (c) $\int_\Sigma \Omega(x)x^\alpha d\sigma(x) = 0$ for all multi-indices $\alpha \in (N \cup \{0\})^n$ with $|\alpha| = N$, where $\Sigma = \{x \in R^n : |x| = 1\}$ is the unit sphere of R^n .

DEFINITION 3. Let $K(x, y) = \Omega(x, y)/|y|^n : R^n \times (R^n \setminus \{0\}) \rightarrow R$. K is said to be a variable Calderón-Zygmund kernel if

- (d) $K(x, \cdot)$ is a Calderón-Zygmund kernel for a.e. $x \in R^n$;
- (e) $\max_{|\gamma| \leq 2n} \left\| \frac{\partial^{|\gamma|}}{\partial y^\gamma} \Omega(x, y) \right\|_{L^\infty(R^n \times \Sigma)} = L < \infty$.

Moreover, let b be a locally integrable function on R^n and T be the singular integral operator with variable Calderón-Zygmund kernel as

$$T(f)(x) = \int_{R^n} K(x, x - y)f(y)dy,$$

where $K(x, x - y) = \frac{\Omega(x, x - y)}{|x - y|^n}$ and that $\Omega(x, y)/|y|^n$ is a variable Calderón-Zygmund kernel. The Toeplitz type operators associated to T is defined by

$$T_b = \sum_{k=1}^m T^{k,1} M_b T^{k,2}$$

and

$$S_b = \sum_{k=1}^m (T^{k,3} M_b I_\alpha T^{k,4} + T^{k,5} I_\alpha M_b T^{k,6}),$$

where $T^{k,1}$ and $T^{k,3}$ are the singular integral operator T with variable Calderón-Zygmund kernel or $\pm I$ (the identity operator), $T^{k,2}$, $T^{k,4}$ and $T^{k,6}$ are the bounded linear operators on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$, $T^{k,5} = \pm I$, $k = 1, \dots, m$, $M_b(f) = bf$ and I_α is the fractional integral operator ($0 < \alpha < n$) (see [2]).

Note that the commutator $[b, T](f) = bT(f) - T(bf)$ is a particular operator of the Toeplitz type operators T_b and S_b . The Toeplitz type operators T_b and S_b are the non-trivial generalizations of the commutator. It is well known that commutators are of great interest in harmonic analysis and have been widely studied by many authors (see [19], [20]). The main purpose of this paper is to prove the M^k -type sharp maximal inequalities for the Toeplitz type operator T_b . As the application, we obtain the boundedness on the Morrey space for the Toeplitz type operators T_b and S_b .

2. Theorems and Lemmas

We shall prove the following theorems.

THEOREM 1. *Let T be the singular integral operator as Definition 3, $0 < \eta < 1$, $1 < s < \infty$ and $b \in BMO(\mathbb{R}^n)$. If $T_1(g) = 0$ for any $g \in L^r(\mathbb{R}^n)$ ($1 < r < \infty$), then there exists a constant $C > 0$ such that, for any $f \in C_0^\infty(\mathbb{R}^n)$ and $\tilde{x} \in \mathbb{R}^n$,*

$$(T_b(f))_\eta^\#(\tilde{x}) \leq C \|b\|_{BMO} \sum_{k=1}^m M^2(T^{k,2}(f))(\tilde{x}).$$

THEOREM 2. *Let T be the singular integral operator as Definition 3, $0 < \eta < 1$, $1 < s < \infty$ and $b \in BMO(\mathbb{R}^n)$. If $S_1(g) = 0$ for any $g \in L^r(\mathbb{R}^n)$ ($1 < r < \infty$), then there exists a constant $C > 0$ such that, for any $f \in C_0^\infty(\mathbb{R}^n)$ and $\tilde{x} \in \mathbb{R}^n$,*

$$(S_b(f))_\eta^\#(\tilde{x}) \leq C \|b\|_{BMO} \sum_{k=1}^m (M^2(I_\alpha T^{k,4}(f))(\tilde{x}) + M_{\alpha,s}(T^{k,6}(f))(\tilde{x})).$$

THEOREM 3. *Let T be the singular integral operator as Definition 3, $1 < p < \infty$, $0 < D < 2^n$ and $b \in BMO(\mathbb{R}^n)$. If $T_1(g) = 0$ for any $g \in L^r(\mathbb{R}^n)$ ($1 < r < \infty$), then T_b is bounded on $L^{p,\varphi}(\mathbb{R}^n)$.*

THEOREM 4. *Let T be the singular integral operator as Definition 3, $0 < D < 2^n$, $1 < p < n/\alpha$, $1/q = 1/p - \alpha/n$ and $b \in BMO(\mathbb{R}^n)$. If $S_1(g) = 0$ for any $g \in L^r(\mathbb{R}^n)$ ($1 < r < \infty$), then S_b is bounded from $L^{p,\varphi}(\mathbb{R}^n)$ to $L^{q,\varphi}(\mathbb{R}^n)$.*

COROLLARY. *Let $[b, T](f) = bT(f) - T(bf)$ be the commutator generated by the singular integral operator T as Definition 3 and b . Then Theorems 1–4 hold for $[b, T]$.*

To prove the theorems, we need the following lemmas.

LEMMA 1. ([7, p. 485]) *Let $0 < p < q < \infty$ and for any function $f \geq 0$. We define that, for $1/r = 1/p - 1/q$,*

$$\|f\|_{WL^q} = \sup_{\lambda > 0} \lambda \{x \in \mathbb{R}^n : f(x) > \lambda\}^{1/q}, \quad N_{p,q}(f) = \sup_Q \|f\chi_Q\|_{L^p} / \|\chi_Q\|_{L^r},$$

where the sup is taken for all measurable sets Q with $0 < |Q| < \infty$. Then

$$\|f\|_{WL^q} \leq N_{p,q}(f) \leq (q/(q-p))^{1/p} \|f\|_{WL^q}.$$

LEMMA 2. (see [1]) Let T be the singular integral operator as Definition 3. Then T is bounded on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$, and weak (L^1, L^1) bounded.

LEMMA 3. (see [19]) We have

$$\frac{1}{|Q|} \int_Q |f(x)g(x)| dx \leq \|f\|_{\exp L, Q} \|g\|_{L(\log L), Q}.$$

LEMMA 4. (see [2], [7]) Let $0 < \alpha < n$, $1 \leq s < p < n/\alpha$ and $1/q = 1/p - \alpha/n$. Then

$$\|M_{\alpha,s}(f)\|_{L^q} \leq C \|f\|_{L^p}$$

and

$$\|I_\alpha(f)\|_{L^q} \leq C \|f\|_{L^p}.$$

LEMMA 5. (see [7]) Let $0 < p, \eta < \infty$. Then, for any smooth function f for which the left-hand side is finite,

$$\int_{\mathbb{R}^n} M_\eta(f)(x)^p dx \leq C \int_{\mathbb{R}^n} f_\eta^\#(f)(x)^p dx.$$

LEMMA 6. Let $0 < p, \eta < \infty$ and $0 < D < 2^n$. Then, for any smooth function f for which the left-hand side is finite,

$$\|M_\eta(f)\|_{L^{p,\varphi}} \leq C \|f_\eta^\#(f)\|_{L^{p,\varphi}}.$$

Proof. For any cube $Q = Q(x_0, d)$ in \mathbb{R}^n , we know $M(\chi_Q) \in A_1$ for any cube $Q = Q(x, d)$ by [3]. Noticing that $M(\chi_Q) \leq 1$ and $M(\chi_Q)(x) \leq d^n / (|x - x_0| - d)^n$ if $x \in Q^c$, by Lemma 5, we have, for $f \in L^{p,\varphi}(\mathbb{R}^n)$,

$$\begin{aligned} & \int_Q M_\eta(f)(x)^p dx = \int_{\mathbb{R}^n} M_\eta(f)(x)^p \chi_Q(x) dx \\ & \leq \int_{\mathbb{R}^n} M_\eta(f)(x)^p M(\chi_Q)(x) dx \\ & \leq C \int_{\mathbb{R}^n} f_\eta^\#(x)^p M(\chi_Q)(x) dx \\ & = C \left(\int_Q f_\eta^\#(x)^p M(\chi_Q)(x) dx + \sum_{k=0}^\infty \int_{2^{k+1}Q \setminus 2^kQ} f_\eta^\#(x)^p M(\chi_Q)(x) dx \right) \\ & \leq C \left(\int_Q f_\eta^\#(x)^p dx + \sum_{k=0}^\infty \int_{2^{k+1}Q \setminus 2^kQ} f_\eta^\#(x)^p \frac{|Q|}{|2^{k+1}Q|} dx \right) \end{aligned}$$

$$\begin{aligned}
 &\leq C \left(\int_Q f_\eta^\#(x)^p dx + \sum_{k=0}^\infty \int_{2^{k+1}Q} f_\eta^\#(x)^p 2^{-kn} dy \right) \\
 &\leq C \|f_\eta^\#\|_{L^{p,\varphi}}^p \sum_{k=0}^\infty 2^{-kn} \varphi(2^{k+1}d) \\
 &\leq C \|f_\eta^\#\|_{L^{p,\varphi}}^p \sum_{k=0}^\infty (2^{-n}D)^k \varphi(d) \\
 &\leq C \|f_\eta^\#\|_{L^{p,\varphi}}^p \varphi(d),
 \end{aligned}$$

thus

$$\left(\frac{1}{\varphi(d)} \int_Q M_\eta(f)(x)^p dx \right)^{1/p} \leq C \left(\frac{1}{\varphi(d)} \int_Q f_\eta^\#(x)^p dx \right)^{1/p}$$

and

$$\|M_\eta(f)\|_{L^{p,\varphi}} \leq C \|f_\eta^\#\|_{L^{p,\varphi}}.$$

This finishes the proof. \square

LEMMA 7. Let $0 < \alpha < n$, $0 < D < 2^n$, $1 \leq s < p < n/\alpha$ and $1/q = 1/p - \alpha/n$. Then

$$\|M_{\alpha,s}(f)\|_{L^{q,\varphi}} \leq C \|f\|_{L^{p,\varphi}}$$

and

$$\|I_\alpha(f)\|_{L^{q,\varphi}} \leq C \|f\|_{L^{p,\varphi}}.$$

LEMMA 8. Let T be the bounded linear operators on $L^r(\mathbb{R}^n)$ for any $1 < r < \infty$. Then, for $1 < p < \infty$ and $0 < D < 2^n$,

$$\|T(f)\|_{L^{p,\varphi}} \leq C \|f\|_{L^{p,\varphi}}.$$

The proofs of two Lemmas are similar to that of Lemma 6 by Lemma 4, we omit the details.

3. Proofs of Theorems

Proof of Theorem 1. It suffices to prove for $f \in C_0^\infty(\mathbb{R}^n)$ and some constant C_0 , the following inequality holds:

$$\left(\frac{1}{|Q|} \int_Q |T_b(f)(x) - C_0|^\eta dx \right)^{1/\eta} \leq C \|b\|_{BMO} \sum_{k=1}^m M^2(T^{k,2}(f))(\bar{x}).$$

Without loss of generality, we may assume $T^{k,1}$ are T ($k = 1, \dots, m$). Fix a cube $Q = Q(x_0, d)$ and $\bar{x} \in Q$. We write, by $T_1(g) = 0$,

$$T_b(f)(x) = T_{b-b_{2Q}}(f)(x) = T_{(b-b_{2Q})\chi_{2Q}}(f)(x) + T_{(b-b_{2Q})\chi_{(2Q)^c}}(f)(x) = f_1(x) + f_2(x).$$

Then

$$\begin{aligned} & \left(\frac{1}{|Q|} \int_Q |T_b(f)(x) - f_2(x_0)|^\eta dx \right)^{1/\eta} \\ & \leq C \left(\frac{1}{|Q|} \int_Q |f_1(x)|^\eta dx \right)^{1/\eta} + C \left(\frac{1}{|Q|} \int_Q |f_2(x) - f_2(x_0)|^\eta dx \right)^{1/\eta} = I + II. \end{aligned}$$

For I , by Lemma 1, 2 and 3, we obtain

$$\begin{aligned} & \left(\frac{1}{|Q|} \int_Q |T^{k,1} M_{(b-b_{2Q})\chi_{2Q}} T^{k,2}(f)(x)|^\eta dx \right)^{1/\eta} \\ & \leq |Q|^{-1} \frac{\|T^{k,1} M_{(b-b_{2Q})\chi_{2Q}} T^{k,2}(f)\chi_Q\|_{L^\eta}}{|Q|^{1/\eta-1}} \\ & \leq C|Q|^{-1} \|T^{k,1} M_{(b-b_{2Q})\chi_{2Q}} T^{k,2}(f)\|_{W^{L^1}} \\ & \leq C|Q|^{-1} \|M_{(b-b_{2Q})\chi_{2Q}} T^{k,2}(f)\|_{L^1} \\ & \leq C|Q|^{-1} \int_{2Q} |b(x) - b_{2Q}| |T^{k,2}(f)(x)| dx \\ & \leq C \|b - b_{2Q}\|_{\text{exp}L, 2Q} \|T^{k,2}(f)\|_{L(\log L), 2Q} \\ & \leq C \|b\|_{BMO} M^2(T^{k,2}(f))(\tilde{x}), \end{aligned}$$

thus

$$\begin{aligned} I & \leq \sum_{k=1}^m \left(\frac{C}{|Q|} \int_Q |T^{k,1} M_{(b-b_{2Q})\chi_{2Q}} T^{k,2}(f)(x)|^\eta dx \right)^{1/\eta} \\ & \leq C \|b\|_{BMO} \sum_{k=1}^m M^2(T^{k,2}(f))(\tilde{x}). \end{aligned}$$

For II , by [1], we know that

$$T(f)(x) = \sum_{u=1}^\infty \sum_{v=1}^{g_u} a_{uv}(x) \int_{R^n} \frac{Y_{uv}(x-y)}{|x-y|^{n+m}} f(y) dy,$$

where $g_u \leq Cu^{n-2}$, $\|a_{uv}\|_{L^\infty} \leq Cu^{-2n}$, $|Y_{uv}(x-y)| \leq Cu^{n/2-1}$ and

$$\left| \frac{Y_{uv}(x-y)}{|x-y|^n} - \frac{Y_{uv}(x_0-y)}{|x_0-y|^n} \right| \leq Cu^{n/2} |x-x_0|/|x_0-y|^{n+1}$$

for $|x-y| > 2|x_0-x| > 0$. Then, we get, for $x \in Q$,

$$\begin{aligned} & |T^{k,1} M_{(b-b_{2Q})\chi_{(2Q)^c}}(f)(x) - T^{k,1} M_{(b-b_{2Q})\chi_{(2Q)^c}} T^{k,2}(f)(x_0)| \\ & \leq \int_{(2Q)^c} |b(y) - b_{2Q}| |K(x, x-y) - K(x_0, x_0-y)| |T^{k,2}(f)(y)| dy \end{aligned}$$

$$\begin{aligned}
 &= \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| |K(x, x-y) - K(x_0, x_0-y)| |T^{k,2}(f)(y)| dy \\
 &\leq C \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| \sum_{u=1}^{\infty} \sum_{v=1}^{g_u} |a_{uv}(x)| \\
 &\quad \times \left| \frac{Y_{uv}(x-y)}{|x-y|^n} - \frac{Y_{uv}(x_0-y)}{|x_0-y|^n} \right| |T^{k,2}(f)(y)| dy \\
 &\leq C \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| \frac{|x-x_0|}{|x_0-y|^{n+1}} |T^{k,2}(f)(y)| dy \\
 &\leq C \sum_{j=1}^{\infty} \frac{d}{(2^{j+1}d)^{n+1}} \int_{2^{j+1}Q} |b(y) - b_{2Q}| |T^{k,2}(f)(y)| dy \\
 &\leq C \sum_{j=1}^{\infty} 2^{-j} \frac{1}{|2^{j+1}Q|} \int_{2^{j+1}Q} |b(y) - b_{2Q}| |T^{k,2}(f)(y)| dy \\
 &\leq C \sum_{j=1}^{\infty} 2^{-j} \|b - b_{2Q}\|_{\text{exp}L, 2^{j+1}Q} \|T^{k,2}(f)\|_{L(\log L), 2^{j+1}Q} \\
 &\leq C \sum_{j=1}^{\infty} j 2^{-j} \|b\|_{BMO} M^2(T^{k,2}(f))(\tilde{x}) \\
 &\leq C \|b\|_{BMO} M^2(T^{k,2}(f))(\tilde{x}),
 \end{aligned}$$

thus

$$\begin{aligned}
 II &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^m |T^{k,1} M_{(b-b_{2Q})\chi_{(2Q)^c}} T^{k,2}(f)(x) - T^{k,1} M_{(b-b_{2Q})\chi_{(2Q)^c}} T^{k,2}(f)(x_0)| dx \\
 &\leq C \|b\|_{BMO} \sum_{k=1}^m M^2(T^{k,2}(f))(\tilde{x}).
 \end{aligned}$$

This completes the proof of Theorem 1. \square

Proof of Theorem 2. It suffices to prove for $f \in C_0^\infty(R^n)$ and some constant C_0 , the following inequality holds:

$$\left(\frac{1}{|Q|} \int_Q |S_b(f)(x) - C_0|^\eta dx \right)^{1/\eta} \leq C \|b\|_{BMO} \sum_{k=1}^m (M^2(I_\alpha T^{k,4}(f))(\tilde{x}) + M_{\alpha,s}(T^{k,6}(f))(\tilde{x})).$$

Without loss of generality, we may assume $T^{k,3}$ are $T(k = 1, \dots, m)$. Fix a cube $Q = Q(x_0, d)$ and $\tilde{x} \in Q$. Write, by $T_1(g) = 0$,

$$\begin{aligned}
 S_b(f)(x) &= \sum_{k=1}^m T^{k,3} M_b I_\alpha T^{k,4}(f)(x) + \sum_{k=1}^m T^{k,5} I_\alpha M_b T^{k,6}(f)(x) \\
 &= A_b(x) + B_b(x) = A_{b-b_Q}(x) + B_{b-b_Q}(x),
 \end{aligned}$$

where

$$\begin{aligned} A_{b-b_Q}(x) &= \sum_{k=1}^m T^{k,3} M_{(b-b_Q)\chi_{2Q}} I_\alpha T^{k,4}(f)(x) + \sum_{k=1}^m T^{k,3} M_{(b-b_Q)\chi_{(2Q)^c}} I_\alpha T^{k,4}(f)(x) \\ &= A_1(x) + A_2(x) \end{aligned}$$

and

$$\begin{aligned} B_{b-b_Q}(x) &= \sum_{k=1}^m T^{k,5} I_\alpha M_{(b-b_Q)\chi_{2Q}} T^{k,6}(f)(x) + \sum_{k=1}^m T^{k,5} I_\alpha M_{(b-b_Q)\chi_{(2Q)^c}} T^{k,6}(f)(x) \\ &= B_1(x) + B_2(x). \end{aligned}$$

Then

$$\begin{aligned} &\left(\frac{1}{|Q|} \int_Q |A_{b-b_Q}(f)(x) - A_2(x_0)|^\eta dx \right)^{1/\eta} \\ &\leq C \left(\frac{1}{|Q|} \int_Q |A_1(x)|^\eta dx \right)^{1/\eta} + C \left(\frac{1}{|Q|} \int_Q |A_2(x) - A_2(x_0)|^\eta dx \right)^{1/\eta} \\ &= I_1 + I_2 \end{aligned}$$

and

$$\begin{aligned} &\left(\frac{1}{|Q|} \int_Q |B_{b-b_Q}(f)(x) - B_2(x_0)|^\eta dx \right)^{1/\eta} \\ &\leq C \left(\frac{1}{|Q|} \int_Q |B_1(x)|^\eta dx \right)^{1/\eta} + C \left(\frac{1}{|Q|} \int_Q |B_2(x) - B_2(x_0)|^\eta dx \right)^{1/\eta} \\ &= I_3 + I_4. \end{aligned}$$

For I_1 and I_2 , by using the same argument as in the proof of Theorem 1, we get

$$\begin{aligned} I_1 &\leq C \sum_{k=1}^m \left(\frac{1}{|Q|} \int_{R^n} |T^{k,3} M_{(b-b_Q)\chi_{2Q}} I_\alpha T^{k,4}(f)(x)|^\eta dx \right)^{1/\eta} \\ &\leq \sum_{k=1}^m |Q|^{-1} \frac{\|T^{k,3} M_{(b-b_Q)\chi_{2Q}} I_\alpha T^{k,4}(f)\chi_Q\|_{L^\eta}}{|Q|^{1/\eta-1}} \\ &\leq C \sum_{k=1}^m |Q|^{-1} \|T^{k,3} M_{(b-b_Q)\chi_{2Q}} I_\alpha T^{k,4}(f)\|_{W L^1} \\ &\leq C \sum_{k=1}^m |Q|^{-1} \|M_{(b-b_Q)\chi_{2Q}} I_\alpha T^{k,4}(f)\|_{L^1} \\ &\leq C \sum_{k=1}^m |Q|^{-1} \int_{2Q} |b(x) - b_{2Q}| |I_\alpha T^{k,4}(f)(x)| dx \\ &\leq C \sum_{k=1}^m \|b - b_{2Q}\|_{\text{exp}L, 2Q} \|I_\alpha T^{k,4}(f)\|_{L(\log L), 2Q} \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m M^2(I_\alpha T^{k,4}(f))(\tilde{x}), \end{aligned}$$

$$\begin{aligned}
 I_2 &\leq C \sum_{k=1}^m \frac{1}{|Q|} \int_Q \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| |K(x, x-y) - K(x_0, x_0-y)| \\
 &\quad \times |I_{\alpha} T^{k,4}(f)(y)| dy dx \\
 &\leq C \sum_{k=1}^m \frac{1}{|Q|} \int_Q \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| \frac{|x-x_0|}{|x_0-y|^{n+1}} |I_{\alpha} T^{k,4}(f)(y)| dy dx \\
 &\leq C \sum_{k=1}^m \sum_{j=1}^{\infty} 2^{-j} \frac{1}{|2^{j+1}Q|} \int_{2^{j+1}Q} |b(y) - b_{2Q}| |I_{\alpha} T^{k,4}(f)(y)| dy \\
 &\leq C \sum_{k=1}^m \sum_{j=1}^{\infty} 2^{-j} \|b - b_{2Q}\|_{\text{exp}L, 2^{j+1}Q} \|I_{\alpha} T^{k,4}(f)\|_{L(\log L), 2^{j+1}Q} \\
 &\leq C \sum_{k=1}^m \sum_{j=1}^{\infty} j 2^{-j} \|b\|_{BMO} M^2(I_{\alpha} T^{k,4}(f))(\tilde{x}) \\
 &\leq C \|b\|_{BMO} \sum_{k=1}^m M^2(I_{\alpha} T^{k,4}(f))(\tilde{x}).
 \end{aligned}$$

For I_3 , by Lemma 4, we get, for $1 < p < \min(s, n/\alpha)$ and $1/q = 1/p - \alpha/n$,

$$\begin{aligned}
 I_3 &\leq C \sum_{k=1}^m \left(\frac{1}{|Q|} \int_{R^n} |I_{\alpha} M_{(b-b_Q)\chi_{2Q}} T^{k,6}(f)(x)|^q dx \right)^{1/q} \\
 &\leq C \sum_{k=1}^m |Q|^{-1/q} \left[\int_{2Q} (|b(x) - b_Q| |T^{k,6}(f)(x)|)^p dx \right]^{1/p} \\
 &\leq C \sum_{k=1}^m \left(\frac{1}{|2Q|} \int_{2Q} |b(x) - b_Q|^{ps/(s-p)} dx \right)^{(s-p)/ps} \left(\frac{1}{|2Q|^{1-s\alpha/n}} \int_{2Q} |T^{k,6}(f)(x)|^s dx \right)^{1/s} \\
 &\leq C \|b\|_{BMO} \sum_{k=1}^m M_{\alpha, s}(T^{k,6}(f))(\tilde{x}).
 \end{aligned}$$

For I_4 , we get

$$\begin{aligned}
 I_4 &\leq |Q|^{-1} \sum_{k=1}^m \int_Q \int_{(2Q)^c} |b(y) - b_{2Q}| \left| \frac{1}{|x-y|^{n-\alpha}} - \frac{1}{|x_0-y|^{n-\alpha}} \right| |T^{k,6}(f)(y)| dy dx \\
 &\leq C \sum_{k=1}^m \sum_{j=1}^{\infty} \int_{2^j d \leq |y-x_0| < 2^{j+1} d} |b(y) - b_{2Q}| \frac{d}{|x_0-y|^{n-\alpha+1}} |T^{k,6}(f)(y)| dy \\
 &\leq C \sum_{k=1}^m \sum_{j=1}^{\infty} d(2^j d)^{-n+\alpha-1} (2^j d)^{n(1-1/s)} (2^j d)^{n/s-\alpha} \\
 &\quad \times \left(\frac{1}{|2^{j+1}Q|} \int_{2^{j+1}Q} |b(y) - b_Q|^{s'} dy \right)^{1/s'} \left(\frac{1}{|2^{j+1}Q|^{1-s\alpha/n}} \int_{2^{j+1}Q} |T^{k,6}(f)(y)|^s dy \right)^{1/s}
 \end{aligned}$$

$$\begin{aligned} &\leq C \|b\|_{BMO} \sum_{k=1}^m M_{\alpha,s}(T^{k,6}(f))(\tilde{x}) \sum_{j=1}^{\infty} j 2^{-j} \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m M_{\alpha,s}(T^{k,6}(f))(\tilde{x}). \end{aligned}$$

This completes the proof of Theorem 2. \square

Proof of Theorem 3. By Theorem 1 and Lemmas 7–8, we have

$$\begin{aligned} \|T_b(f)\|_{L^{p,\varphi}} &\leq \|M_{\eta}(T_b(f))\|_{L^{p,\varphi}} \leq C \|(T_b(f))_{\eta}^{\#}\|_{L^{p,\varphi}} \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m \|M^2(T^{k,2}(f))\|_{L^{p,\varphi}} \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m \|T^{k,2}(f)\|_{L^{p,\varphi}} \\ &\leq C \|b\|_{BMO} \|f\|_{L^{p,\varphi}}. \end{aligned}$$

This completes the proof of Theorem 3. \square

Proof of Theorem 4. Choose $1 < s < p$ in Theorem 2, then we have, by Lemmas 7 and 8,

$$\begin{aligned} \|S_b(f)\|_{L^{q,\varphi}} &\leq \|M_{\eta}(S_b(f))\|_{L^{q,\varphi}} \leq C \|(S_b(f))_{\eta}^{\#}\|_{L^{q,\varphi}} \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m (\|M^2(I_{\alpha}T^{k,4}(f))\|_{L^{q,\varphi}} + \|M_{\alpha,s}(T^{k,6}(f))\|_{L^{q,\varphi}}) \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m (\|I_{\alpha}T^{k,4}(f)\|_{L^{q,\varphi}} + \|T^{k,6}(f)\|_{L^{p,\varphi}}) \\ &\leq C \|b\|_{BMO} \sum_{k=1}^m (\|T^{k,4}(f)\|_{L^{p,\varphi}} + \|f\|_{L^{p,\varphi}}) \\ &\leq C \|b\|_{BMO} \|f\|_{L^{p,\varphi}}. \end{aligned}$$

This completes the proof of Theorem 4. \square

REFERENCES

- [1] A. P. CALDERÓN AND A. ZYGMUND, *On singular integrals with variable kernels*, Appl. Anal., **7** (1978), 221–238.
- [2] S. CHANILLO, *A note on commutators*, Indiana Univ. Math. J., **31** (1982), 7–16.
- [3] R. COIFMAN AND R. ROCHBERG, *Another characterization of BMO*, Proc. Amer. Math. Soc., **79**, 249–254 (1980).
- [4] R. R. COIFMAN, R. ROCHBERG AND G. WEISS, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math., **103** (1976), 611–635.
- [5] G. DI FAZIO AND M. A. RAGUSA, *Commutators and Morrey spaces*, Boll. Un. Mat. Ital., **5-A** (7) (1991), 323–332.

- [6] G. DI FAZIO AND M. A. RAGUSA, *Interior estimates in Morrey spaces for strong solutions to nondivergence form equations with discontinuous coefficients*, J. Func. Anal., **112** (1993), 241–256.
- [7] J. GARCIA-CUERVA AND J. L. RUBIO DE FRANCIA, *Weighted norm inequalities and related topics*, North-Holland Math., 16, Amsterdam, 1985.
- [8] S. KRANTZ AND S. LI, *Boundedness and compactness of integral operators on spaces of homogeneous type and applications*, J. of Math. Anal. Appl., **258** (2001), 629–641.
- [9] Y. LIN AND S. Z. LU, *Toeplitz type operators associated to strongly singular integral operator*, Sci. in China (ser. A), **36** (2006), 615–630.
- [10] L. Z. LIU, *Interior estimates in Morrey spaces for solutions of elliptic equations and weighted boundedness for commutators of singular integral operators*, Acta Math. Scientia, **25** (B) (1) (2005), 89–94.
- [11] L. Z. LIU, *The continuity for multilinear singular integral operators with variable Calderón-Zygmund kernel on Hardy and Herz spaces*, Siberia Electronic Math. Reports, **2** (2005), 156–166.
- [12] L. Z. LIU, *Good λ estimate for multilinear singular integral operators with variable Calderón-Zygmund kernel*, Kragujevac J. of Math., **27** (2005), 19–30.
- [13] L. Z. LIU, *Weighted estimates of multilinear singular integral operators with variable Calderón-Zygmund kernel for the extreme cases*, Vietnam J. of Math., **34** (1) (2006), 51–61.
- [14] S. Z. LU AND H. X. MO, *Toeplitz type operators on Lebesgue spaces*, Acta Math. Scientia, **29** (B) (1) (2009), 140–150.
- [15] S. Z. LU, D. C. YANG AND Z. S. ZHOU, *Oscillatory singular integral operators with Calderón-Zygmund kernels*, Southeast Asian Bull. of Math., **23** (1999), 457–470.
- [16] T. MIZUHARA, *Boundedness of some classical operators on generalized Morrey spaces*, in “Harmonic Analysis”, Proceedings of a conference held in Sendai, Japan, 1990, 183–189.
- [17] J. PEETRE, *On convolution operators leaving $L^{p,\lambda}$ -spaces invariant*, Ann. Mat. Pura. Appl., **72** (1966), 295–304.
- [18] J. PEETRE, *On the theory of $L^{p,\lambda}$ -spaces*, J. Func. Anal., **4** (1969), 71–87.
- [19] C. PÉREZ, *Endpoint estimate for commutators of singular integral operators*, J. Func. Anal., **128** (1995), 163–185.
- [20] C. PÉREZ AND R. TRUJILLO-GONZALEZ, *Sharp weighted estimates for multilinear commutators*, J. London Math. Soc., **65** (2002), 672–692.
- [21] E. M. STEIN, *Harmonic analysis: real variable methods, orthogonality and oscillatory integrals*, Princeton Univ. Press, Princeton NJ, 1993.
- [22] H. XU AND L. Z. LIU, *Weighted boundedness for multilinear singular integral operator with variable Calderón-Zygmund kernel*, African Diaspora J. of Math., **6** (1) (2008), 1–12.

(Received July 30, 2012)

Chuangxia Huang
Department of Mathematics
Changsha University of Science and Technology
Changsha 410077, P. R. of China

Sheng Guo
Department of Mathematics
Changsha University of Science and Technology
Changsha 410077, P. R. of China

Lanzhe Liu
Department of Mathematics
Changsha University of Science and Technology
Changsha 410077, P. R. of China
e-mail: lanzheliu@163.com