

SOME REMARKS ON (s, m) -CONVEXITY IN THE SECOND SENSE

NOHA EFTEKHARI

(Communicated by G. Toader)

Abstract. The aim of this work is to establish several inequalities for functions whose first derivative in absolute value are (s, m) -convex in the second sense. Some estimates to the left hand side of the Hermite-Hadamard type inequality for (s, m) -convex functions in the second sense are given.

1. Introduction

Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function, the following double inequality is well known as the Hermite-Hadamard integral inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}.$$

In 1984 [5], G. Toader defined the class of m -convex functions. This class is defined in the following way: a function $f : [0, b] \rightarrow \mathbb{R}$ is said to be m -convex, where $m \in [0, 1]$, if for every $x, y \in [0, b]$ and $t \in [0, 1]$ we have

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y).$$

In 1993 [4], V. Mihesan introduced the class of (α, m) -convex functions as the following: a function $f : [0, b] \rightarrow \mathbb{R}$ is said to be (α, m) -convex, where $(\alpha, m) \in [0, 1]^2$, if for every $x, y \in [0, b]$ and $t \in [0, 1]$ we have

$$f(tx + m(1-t)y) \leq t^\alpha f(x) + m(1-t^\alpha)f(y).$$

For recent results and generalizations and new inequalities concerning (α, m) -convex functions, see [2, 6].

In 1978 [3], W. W. Breckner introduced the class of s -convex functions in the second sense, in the following way: a function $f : [0, \infty) \rightarrow \mathbb{R}$ is said to be s -convex in the second sense if for all $x, y \in [0, \infty)$, $t \in [0, 1]$ and for some fixed $s \in (0, 1]$, we have

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y).$$

Now, we combine two definitions of m -convexity and s -convexity in the second sense and obtain the class of (s, m) -convex functions in the second sense as the following.

Mathematics subject classification (2010): 26D10, 39B62.

Keywords and phrases: Hermite-Hadamard inequality, (s, m) -convex function in the second sense, convexity, Hölder's inequality.

DEFINITION 1.1. A function $f : [0, b] \rightarrow \mathbb{R}$ is said to be (s, m) -convex in the second sense, where $(s, m) \in (0, 1]^2$, if for every $x, y \in [0, b]$ and $t \in [0, 1]$ we have

$$f(tx + m(1 - t)y) \leq t^s f(x) + m(1 - t)^s f(y).$$

The aim of this work is to establish some upper bounds to the left hand side of the Hermite-Hadamard type inequality for (s, m) -convex functions in the second sense.

2. Main results

In order to prove new theorems, we need Lemma 1 in [1] by setting $x = \frac{a+b}{2}$, which implies the following lemma.

LEMMA 2.1. Let $f : I \rightarrow \mathbb{R}$, $I \subset \mathbb{R}$ be a differentiable function on I° . If $f' \in L^1[a, b]$, where $a, b \in I$ with $a < b$, then the following equality holds:

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \\ &= \frac{b-a}{4} \int_0^1 t \left[f'\left(t\frac{a+b}{2} + (1-t)a\right) - f'\left(t\frac{a+b}{2} + (1-t)b\right) \right] dt. \end{aligned}$$

The following results may be stated:

THEOREM 2.1. Let $f : [a, b] \rightarrow \mathbb{R}$, $[a, b] \subset [0, \infty)$ be a differentiable function on (a, b) such that $f' \in L^1[a, b]$. If $|f'|$ is (s, m) -convex in the second sense on $[a, b]$ for $(s, m) \in (0, 1]^2$, then the following inequality holds:

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{4(s+2)} \left[2 \left| f'\left(\frac{a+b}{2}\right) \right| + \frac{m}{s+1} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right]. \end{aligned}$$

Proof. By Lemma 2.1, it follows that

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{4} \int_0^1 t \left[\left| f'\left(t\frac{a+b}{2} + (1-t)a\right) \right| + \left| f'\left(t\frac{a+b}{2} + (1-t)b\right) \right| \right] dt. \end{aligned} \tag{1}$$

Since $|f'|$ is (s, m) -convex in the second sense on $[a, b]$, for any $t \in [0, 1]$, we obtain

$$\left| f'\left(t\frac{a+b}{2} + (1-t)a\right) \right| \leq t^s \left| f'\left(\frac{a+b}{2}\right) \right| + m(1-t)^s \left| f'\left(\frac{a}{m}\right) \right|, \tag{2}$$

$$\left| f'\left(t\frac{a+b}{2} + (1-t)b\right) \right| \leq t^s \left| f'\left(\frac{a+b}{2}\right) \right| + m(1-t)^s \left| f'\left(\frac{b}{m}\right) \right|. \tag{3}$$

Now (1), (2) and (3) imply

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{4} \int_0^1 \left[2t^{s+1} \left| f'\left(\frac{a+b}{2}\right) \right| + mt(1-t)^s \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right] dt \\ & = \left(\frac{b-a}{4}\right) \left[\frac{2}{s+2} \left| f'\left(\frac{a+b}{2}\right) \right| + \frac{m}{(s+1)(s+2)} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right] \\ & = \left(\frac{b-a}{4(s+2)}\right) \left[2 \left| f'\left(\frac{a+b}{2}\right) \right| + \frac{m}{s+1} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right]. \quad \square \end{aligned}$$

THEOREM 2.2. *Suppose that all the assumptions of Theorem 2.1 are satisfied. Then the following inequality holds:*

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{b-a}{2^{s+2}(s+2)} \left[\left| f'(a) \right| + \left| f'(b) \right| + \frac{m(2^{s+2} - s + 3)}{s+1} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right]. \end{aligned}$$

Proof. Since $t\left(\frac{a+b}{2}\right) + (1-t)a = \frac{t}{2}b + (1-\frac{t}{2})a$ and $|f'|$ is (s, m) -convex in the second sense on $[a, b]$, then for any $t \in [0, 1]$, one can get the following inequalities:

$$\left| f'\left(t\frac{a+b}{2} + (1-t)a\right) \right| \leq \left(\frac{t}{2}\right)^s |f'(b)| + m\left(1-\frac{t}{2}\right)^s \left| f'\left(\frac{a}{m}\right) \right|, \quad (4)$$

$$\left| f'\left(t\frac{a+b}{2} + (1-t)b\right) \right| \leq \left(\frac{t}{2}\right)^s |f'(a)| + m\left(1-\frac{t}{2}\right)^s \left| f'\left(\frac{b}{m}\right) \right|, \quad (5)$$

so (1), (4) and (5) imply

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{b-a}{4} \int_0^1 \left[\frac{t^{s+1}}{2^s} (|f'(a)| + |f'(b)|) + mt\left(1-\frac{t}{2}\right)^s \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right] dt \\ & = \frac{b-a}{2^{s+2}(s+2)} \left[|f'(a)| + |f'(b)| + \frac{m(2^{s+2} - s + 3)}{s+1} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right]. \quad \square \end{aligned}$$

THEOREM 2.3. *Let $f : [a, b] \rightarrow \mathbb{R}$, $[a, b] \subset [0, \infty)$ be a differentiable function on (a, b) such that $f' \in L^1[a, b]$. If $|f'|^q$ is (s, m) -convex in the second sense on $[a, b]$ for*

$(s, m) \in (0, 1]^2$ and $q > 1$, then the following inequalities hold:

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \left(\frac{b-a}{4}\right) \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{s+1}\right)^{\frac{1}{q}} \left[2 \left| f'\left(\frac{a+b}{2}\right) \right| + m^{\frac{1}{q}} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right], \end{aligned} \tag{6}$$

and

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \left(\frac{b-a}{8}\right) \left(\frac{2}{s+2}\right)^{\frac{1}{q}} \left[2 \left| f'\left(\frac{a+b}{2}\right) \right| + \left(\frac{m}{s+1}\right)^{\frac{1}{q}} \left(\left| f'\left(\frac{a}{m}\right) \right| + \left| f'\left(\frac{b}{m}\right) \right| \right) \right]. \end{aligned} \tag{7}$$

Proof. By applying (2), (3) for $|f'|^q$ and by using the Hölder’s inequality for $q > 1$ and $p = \frac{q}{q-1}$, it follows that

$$\begin{aligned} & \int_0^1 t \left| f'\left(t\left(\frac{a+b}{2}\right) + (1-t)a\right) \right| dt \\ & \leq \left(\int_0^1 t^p dt\right)^{\frac{1}{p}} \left(\int_0^1 \left| f'\left(t\left(\frac{a+b}{2}\right) + (1-t)a\right) \right|^q dt\right)^{\frac{1}{q}} \\ & \leq \left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left[\int_0^1 \left(t^s \left| f'\left(\frac{a+b}{2}\right) \right|^q + m(1-t)^s \left| f'\left(\frac{a}{m}\right) \right|^q \right) dt \right]^{\frac{1}{q}} \\ & = \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{s+1}\right)^{\frac{1}{q}} \left[\left| f'\left(\frac{a+b}{2}\right) \right|^q + m \left| f'\left(\frac{a}{m}\right) \right|^q \right]^{\frac{1}{q}}, \end{aligned} \tag{8}$$

and similarly

$$\begin{aligned} & \int_0^1 t \left| f'\left(t\left(\frac{a+b}{2}\right) + (1-t)b\right) \right| dt \\ & \leq \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{s+1}\right)^{\frac{1}{q}} \left[\left| f'\left(\frac{a+b}{2}\right) \right|^q + m \left| f'\left(\frac{b}{m}\right) \right|^q \right]^{\frac{1}{q}}. \end{aligned} \tag{9}$$

Therefore, the inequalities (1), (8), (9) and the following relation imply (6),

$$\sum_{i=1}^n (a_i + b_i)^r \leq \sum_{i=1}^n a_i^r + \sum_{i=1}^n b_i^r, \tag{10}$$

for $0 < r < 1$, $a_1, a_2, \dots, a_n \geq 0$ and $b_1, b_2, \dots, b_n \geq 0$.

Now, if we use the Hölder's inequality in the following way and by applying the inequalities (1), (10), then it follows (7),

$$\begin{aligned}
 & \int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right| dt = \int_0^1 t^{1-\frac{1}{q}} t^{\frac{1}{q}} \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right| dt \\
 & \leq \left(\int_0^1 t dt \right)^{\frac{1}{p}} \left(\int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right|^q dt \right)^{\frac{1}{q}} \\
 & \leq \left(\frac{1}{2} \right)^{\frac{1}{p}} \left[\int_0^1 \left(t^{s+1} \left| f' \left(\frac{a+b}{2} \right) \right|^q + mt(1-t)^s \left| f' \left(\frac{a}{m} \right) \right|^q \right) dt \right]^{\frac{1}{q}} \\
 & = \left(\frac{1}{2} \right)^{\frac{q-1}{q}} \left(\frac{1}{s+2} \right)^{\frac{1}{q}} \left[\left| f' \left(\frac{a+b}{2} \right) \right|^q + \frac{m}{s+1} \left| f' \left(\frac{a}{m} \right) \right|^q \right]^{\frac{1}{q}} \\
 & \leq \left(\frac{1}{2} \right) \left(\frac{2}{s+2} \right)^{\frac{1}{q}} \left[\left| f' \left(\frac{a+b}{2} \right) \right| + \left(\frac{m}{s+1} \right)^{\frac{1}{q}} \left| f' \left(\frac{a}{m} \right) \right| \right]. \quad \square
 \end{aligned}$$

THEOREM 2.4. *Suppose that all the assumptions of Theorem 2.3 are satisfied. Then the following inequalities hold:*

$$\begin{aligned}
 & \left| f \left(\frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left(\frac{b-a}{4} \right) \left(\frac{q-1}{2q-1} \right)^{\frac{q-1}{q}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\frac{1}{2} \right)^{\frac{s}{q}} \\
 & \times \left[|f'(a)| + |f'(b)| + (m(2^{s+1}-1))^{\frac{1}{q}} \left(\left| f' \left(\frac{a}{m} \right) \right| + \left| f' \left(\frac{b}{m} \right) \right| \right) \right], \quad (11)
 \end{aligned}$$

and

$$\begin{aligned}
 & \left| f \left(\frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left(\frac{b-a}{8} \right) \left(\frac{2}{s+2} \right)^{\frac{1}{q}} \left(\frac{1}{2} \right)^{\frac{s}{q}} \\
 & \times \left[|f'(a)| + |f'(b)| + \left(\frac{m(2^{s+2}-s+3)}{s+1} \right)^{\frac{1}{q}} \left(\left| f' \left(\frac{a}{m} \right) \right| + \left| f' \left(\frac{b}{m} \right) \right| \right) \right]. \quad (12)
 \end{aligned}$$

Proof. By applying (4), (5) for $|f'|^q$ and by using the Hölder's inequality for $q > 1$ and $p = \frac{q}{q-1}$, we obtain

$$\begin{aligned}
 & \int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right| dt \\
 & \leq \left(\int_0^1 t^p dt \right)^{\frac{1}{p}} \left(\int_0^1 \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right|^q dt \right)^{\frac{1}{q}}
 \end{aligned}$$

$$\begin{aligned} &\leq \left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left[\int_0^1 \left(\left(\frac{t}{2}\right)^s |f'(b)|^q + m \left(1 - \frac{t}{2}\right)^s \left|f'\left(\frac{a}{m}\right)\right|^q \right) dt \right]^{\frac{1}{q}} \\ &= \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{2}\right)^{\frac{s}{q}} \left[\frac{|f'(b)|^q}{s+1} + m \frac{2^{s+1}-1}{s+1} \left|f'\left(\frac{a}{m}\right)\right|^q \right]^{\frac{1}{q}}, \end{aligned} \tag{13}$$

and similarly

$$\begin{aligned} &\int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)b \right) \right| dt \\ &\leq \left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}} \left(\frac{1}{2}\right)^{\frac{s}{q}} \left[\frac{|f'(a)|^q}{s+1} + m \frac{2^{s+1}-1}{s+1} \left|f'\left(\frac{b}{m}\right)\right|^q \right]^{\frac{1}{q}}. \end{aligned} \tag{14}$$

Therefore, the inequalities (1), (10),(13), (14) imply (11).

Now, the inequalities (1), (10) and (4), (5) for $|f'|^q$ and by using the Hölder’s inequality in the following way, imply (12)

$$\begin{aligned} &\int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right| dt = \int_0^1 t^{1-\frac{1}{q}} t^{\frac{1}{q}} \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right| dt \\ &\leq \left(\int_0^1 t dt \right)^{\frac{1}{p}} \left(\int_0^1 t \left| f' \left(t \left(\frac{a+b}{2} \right) + (1-t)a \right) \right|^q dt \right)^{\frac{1}{q}} \\ &\leq \left(\frac{1}{2}\right)^{\frac{1}{p}} \left[\int_0^1 \left(t \left(\frac{t}{2}\right)^s |f'(b)|^q + mt \left(1 - \frac{t}{2}\right)^s \left|f'\left(\frac{a}{m}\right)\right|^q \right) dt \right]^{\frac{1}{q}} \\ &= \left(\frac{1}{2}\right)^{\frac{q-1}{q}} \left(\frac{1}{2}\right)^{\frac{s}{q}} \left[\frac{|f'(b)|^q}{s+2} + \frac{m(2^{s+2}-s+3)}{(s+1)(s+2)} \left|f'\left(\frac{a}{m}\right)\right|^q \right]^{\frac{1}{q}} \\ &\leq \left(\frac{1}{2}\right)^{\frac{s+q-1}{q}} \left(\frac{1}{s+2}\right)^{\frac{1}{q}} \left[|f'(b)| + \left(\frac{m(2^{s+2}-s+3)}{s+1}\right)^{\frac{1}{q}} \left|f'\left(\frac{a}{m}\right)\right| \right]. \quad \square \end{aligned}$$

3. Applications to special means

We consider the means for arbitrary positive numbers α, β ($\alpha \neq \beta$) as follows:

(1) The arithmetic mean: $A(\alpha, \beta) = \frac{\alpha + \beta}{2},$

(2) The generalized log-mean:

$$L_p(\alpha, \beta) = \left[\frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{\frac{1}{p}}, \quad p \in \mathbb{R} \setminus \{-1, 0\}.$$

Now by using the results of Section 2, we give some applications to special means of real numbers.

PROPOSITION 3.1. *Let $a, b \in (0, \infty)$, $a < b$ and $s \in (0, 1)$. Then it follows that*

$$|A^s(a, b) - L_s^s(a, b)| \leq \frac{(b-a)s}{2(s+2)} \left[\frac{1}{s+1} A(a^{s-1}, b^{s-1}) + A^{s-1}(a, b) \right],$$

and

$$|A^s(a, b) - L_s^s(a, b)| \leq \frac{(b-a)(2^s+1)s}{2^{s-1}(s+1)(s+2)} A(a^{s-1}, b^{s-1}).$$

Proof. The assertions follow from Theorem 2.1 and Theorem 2.2, for $m = 1$ applied to the s -convex function $f(x) = x^s$.

PROPOSITION 3.2. *Let $a, b \in (0, \infty)$, $a < b$ and $s \in (0, 1)$. Then for all $q > 1$, it follows that*

$$|A^s(a, b) - L_s^s(a, b)| \leq s \left(\frac{b-a}{4} \right) \left(\frac{2}{s+2} \right)^{\frac{1}{q}} \left[A^{s-1}(a, b) + \left(\frac{1}{s+1} \right)^{\frac{1}{q}} A(a^{s-1}, b^{s-1}) \right].$$

Proof. The assertion follow from Theorem 2.3, for $m = 1$ applied to the s -convex function $f(x) = x^s$.

Acknowledgements. The author thanks the editor and the referees for their comments.

REFERENCES

- [1] M. ALOMARI, M. DARUS, S. S. DRAGOMIR AND P. CERONE, *Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense*, Appl. Math. Lett., **23**, (2010), 1071–1076.
- [2] M. K. BAKULA, M. E. ÖZDEMİR AND J. PEČARIĆ, *Hadamard type inequalities for m -convex and (α, m) -convex functions*, J. Inequal. Pure Appl. Math., **9**, (2008), Article 96.
- [3] W. W. BRECKNER, *Stetigkeitsaussagen für eine klasse verallgemeinerter konvexer funktionen in topologischen linearen Räumen*, Publ. Inst. Math. (Beograd), **23**, (1978), 13–20.
- [4] V. MIHESAN, *A generalization of the convexity*, Seminar on Functional Equations, Approx. and Convex., Cluj-Napoca, Romania, 1993.
- [5] G. TOADER, *Some generalisations of the convexity*, Proc. Colloq. Approx. Optim., (1984), 329–338.
- [6] G. TOADER, *The hierarchy of convexity and some classic inequalities*, J. Math. Ineq., **3**, 3 (2009), 305–313.

(Received November 22, 2012)

Noha Eftekhari
Shahrekord University
Faculty of Mathematical sciences
P. O. Box 115, Shahrekord
88186-34141, Iran
e-mail: eftekharinoha@yahoo.com