

## NEUMAN-SÁNDOR MEAN, ASYMPTOTIC EXPANSIONS AND RELATED INEQUALITIES

NEVEN ELEZOVIĆ AND LENKA VUKŠIĆ

(Communicated by I. Perić)

*Abstract.* The subject of this paper is a systematic study of inequalities of the form

$$(1 - \mu)M_1 + \mu M_3 \leq M_2 \leq (1 - \nu)M_1 + \nu M_3$$

which cover Neuman-Sándor mean and some classical means. Furthermore, following inequalities with optimal parameters were proved:

$$\mu \frac{1}{H(s,t)} + (1 - \mu) \frac{1}{NS(s,t)} \leq \frac{1}{A(s,t)} \leq \nu \frac{1}{H(s,t)} + (1 - \nu) \frac{1}{NS(s,t)}$$

and

$$\mu \frac{1}{H(s,t)} + (1 - \mu) \frac{1}{N(s,t)} \leq \frac{1}{NS(s,t)} \leq \nu \frac{1}{H(s,t)} + (1 - \nu) \frac{1}{N(s,t)}.$$

### 1. Introduction

The Neuman-Sándor mean is a bivariate mean

$$NS(s,t) = \frac{s-t}{2 \operatorname{arcsinh}\left(\frac{s-t}{s+t}\right)}.$$

first defined by Neuman and Sándor [21]. Beside this mean, our analysis will include following well known means: harmonic mean (H), geometric mean (G), logarithmic mean (L), arithmetic mean (A), centroidal mean (C), root mean square (Q), and contra-harmonic mean (N).

$$\begin{aligned} H(s,t) &= \frac{2st}{s+t}, & G(s,t) &= \sqrt{st}, \\ L(s,t) &= \frac{t-s}{\log t - \log s}, & A(s,t) &= \frac{s+t}{2}, \\ C(s,t) &= \frac{2}{3} \cdot \frac{s^2 + st + t^2}{s+t}, & Q(s,t) &= \sqrt{\frac{s^2 + t^2}{2}}, \\ N(s,t) &= \frac{s^2 + t^2}{s+t}. \end{aligned}$$

*Mathematics subject classification* (2010): 26E60; 41A60.

*Keywords and phrases:* Neuman-Sándor mean, asymptotic expansion, optimal convex combination.

This work has been fully supported by Croatian Science Foundation under the project 5435.

Neuman-Sándor mean has been subject of investigation in various papers recently. Inequalities

$$\mu Q + (1 - \mu)A < NS < \nu Q + (1 - \nu)A \tag{1.1}$$

and

$$\mu N + (1 - \mu)A < NS < \nu N + (1 - \nu)A. \tag{1.2}$$

were studied in [20].

In [28] authors proved that double inequalities

$$\mu_1 H(s, t) + (1 - \mu_1)Q(s, t) < NS(s, t) < \nu_1 H(s, t) + (1 - \nu_1)Q(s, t) \tag{1.3}$$

$$\mu_2 G(s, t) + (1 - \mu_2)Q(s, t) < NS(s, t) < \nu_2 G(s, t) + (1 - \nu_2)Q(s, t) \tag{1.4}$$

$$\mu_3 H(s, t) + (1 - \mu_3)N(s, t) < NS(s, t) < \nu_3 H(s, t) + (1 - \nu_3)N(s, t) \tag{1.5}$$

hold for all  $s, t > 0$  with  $s \neq t$  if and only if  $\mu_1 \geq 2/9$ ,  $\nu_1 \leq 1 - 1/[\sqrt{2} \log(1 + \sqrt{2})]$ ,  $\mu_2 \geq 1/3$ ,  $\nu_2 \leq 1 - 1/[\sqrt{2} \log(1 + \sqrt{2})]$ ,  $\mu_3 \geq 1 - 1/[2 \log(1 + \sqrt{2})]$  and  $\nu_3 \leq 5/12$ .

Qian, Chu [23] proved that the double inequality

$$\mu C(s, t) + (1 - \mu)A(s, t) < NS(s, t) < \nu(s, t)C + (1 - \nu)A(s, t) \tag{1.6}$$

holds for all  $s, t > 0$  with  $s \neq t$  if and only if  $\mu \leq (3 - 3 \log(1 + \sqrt{2}))/\log(1 + \sqrt{2})$  and  $\nu \geq \frac{1}{2}$ . Other similar results can be found in [16, 22, 24, 25, 26, 27].

The aim of this paper is to give a systematic study of double inequalities of the type

$$(1 - \mu)M_1 + \mu M_3 \leq M_2 \leq (1 - \nu)M_1 + \nu M_3 \tag{1.7}$$

which apart from Neuman-Sándor mean also contains two classical means from the list given at the beginning of this section. The analysis of these inequalities will be made using technique of asymptotic expansions which was subject of research in [10]–[15]. After finding coefficients in asymptotic expansion of Neuman-Sándor mean we will be able to establish asymptotic inequalities between means of our interest. That technique will provide us to easily cover large number of relations between means.

## 2. Asymptotic expansion of Neuman-Sándor mean

Asymptotic expansion of any of these means has the following form

$$M(x + s, x + t) \sim x + c_1(s, t) + \frac{c_2(s, t)}{x} + \frac{c_3(s, t)}{x^2} + \dots \tag{2.1}$$

where  $c_n(s, t)$  is a homogeneous polynomial of order  $n$ .

The coefficients  $c_n$  will have simpler form if they are presented in terms of variables  $\alpha$  and  $\beta$  where

$$\alpha = \frac{t + s}{2}, \quad \beta = \frac{t - s}{2}.$$

Let's assume for a moment that  $\alpha = 0$  and that mean  $M$  has asymptotic expansion. Because of the homogeneity we have

$$M(x + s, x + t) = M(x - \beta, x + \beta) = \beta M\left(\frac{x}{\beta} - 1, \frac{x}{\beta} + 1\right) \sim \beta \left(\frac{x}{\beta} + c_1 + c_2 \frac{\beta}{x} + c_3 \frac{\beta^2}{x^2} + \dots\right)$$

i. e. coefficient by  $x^{-n}$  equals  $\beta^{n+1}$  multiplied by constant. Now using Theorem 7.1 from [14] we obtain that for general  $\alpha$ ,  $c_n$  is a homogeneous polynomial of degree  $n$  in variables  $\alpha$  and  $\beta$  and thus in variables  $s$  and  $t$ .

We shall need the first few coefficients in asymptotic expansion of the above mentioned means. These expansions were derived in [14]:

$$\begin{aligned}
 H(x+s, x+t) &\sim x + \alpha - \beta^2 x^{-1} + \alpha \beta^2 x^{-2} - \alpha^2 \beta^2 x^{-3} + \dots \\
 G(x+s, x+t) &\sim x + \alpha - \frac{1}{2} \beta^2 x^{-1} + \frac{1}{2} \alpha \beta^2 x^{-2} - \frac{1}{8} \beta^2 (4\alpha^2 + \beta^2) x^{-3} + \dots \\
 L(x+s, x+t) &\sim x + \alpha - \frac{1}{3} \beta^2 x^{-1} + \frac{1}{3} \alpha \beta^2 x^{-2} - \frac{1}{45} \beta^2 (15\alpha^2 + 4\beta^2) x^{-3} + \dots \\
 A(x+s, x+t) &\sim x + \alpha \\
 C(x+s, x+t) &\sim x + \alpha + \frac{1}{3} \beta^2 x^{-1} - \frac{1}{3} \alpha \beta^2 x^{-2} + \frac{1}{3} \alpha^2 \beta^2 x^{-3} + \dots \\
 Q(x+s, x+t) &\sim x + \alpha + \frac{1}{2} \beta^2 x^{-1} - \frac{1}{2} \alpha \beta^2 x^{-2} + \frac{1}{8} \beta^2 (4\alpha^2 - \beta^2) x^{-3} + \dots \\
 N(x+s, x+t) &\sim x + \alpha + \beta^2 x^{-1} - \alpha \beta^2 x^{-2} + \alpha^2 \beta^2 x^{-3} + \dots
 \end{aligned} \tag{2.2}$$

The following lemma will be used for obtaining asymptotic expansion of Neuman-Sándor mean. It is a special case of Lemma 1.1 from [15].

LEMMA 2.1. *Let function  $f(x)$  have following asymptotic expansion ( $a_0 \neq 0$ ):*

$$f(x) \sim \sum_{n=0}^{\infty} a_n x^{-n}, \quad x \rightarrow \infty.$$

Then

$$\frac{1}{f(x)} \sim \sum_{n=0}^{\infty} b_n x^{-n},$$

where coefficients  $b_n$  are defined by

$$\begin{aligned}
 b_0 &= \frac{1}{a_0}, \\
 b_n &= -\frac{1}{a_0} \sum_{k=1}^n a_k b_{n-k}.
 \end{aligned}$$

Now for  $\alpha = 0$  we have

$$NS(x+s, x+t) = NS(x-\beta, x+\beta) = \frac{\beta}{\operatorname{arcsinh}(\beta/x)} = x \sum_{n=0}^{\infty} b_n x^{-n}.$$

Coefficients  $(b_k)$  can be calculated by inverting Maclaurin series of  $\operatorname{arcsinh}$  function, using Lemma 2.1 with

$$a_k = (-1)^k \binom{k-\frac{1}{2}}{k} \frac{\beta^{2k+1}}{2k+1}.$$

Then the coefficients in the case  $\alpha \neq 0$  can be obtained using Theorem 7.1 from the paper [14]. It holds

$$\begin{aligned} c_0 &= 1, \\ c_1 &= \alpha, \\ c_{n+2} &= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} b_{k+2} \alpha^{n-k}. \end{aligned}$$

The first few coefficients of the Neuman-Sándor mean are:

$$\begin{aligned} c_0 &= 1, & c_4 &= \frac{1}{360} \beta^2 (60\alpha^2 - 17\beta^2), \\ c_1 &= \alpha, & c_5 &= -\frac{1}{120} \alpha \beta^2 (20\alpha^2 - 17\beta^2), \\ c_2 &= \frac{\beta^2}{6}, & c_6 &= \frac{1}{15120} \beta^2 (2520\alpha^4 - 4284\alpha^2 \beta^2 + 367\beta^4), \\ c_3 &= -\frac{\alpha \beta^2}{6}, & & \vdots \end{aligned}$$

### 3. Comparison of means. Asymptotic inequalities

It is natural to define a relation between means based on asymptotic expansion, see [11] for details.

DEFINITION 3.1. Let  $M_1$  and  $M_2$  be any two means, and

$$M_1(x + s, x + t) - M_2(x + s, x + t) = c_k(t, s)x^{-k+1} + \mathcal{O}(x^{-k}). \tag{3.1}$$

If  $c_k(s, t) > 0$  for all  $s$  and  $t$ , then we say that mean  $M_1$  is *asymptotically* greater than mean  $M_2$ , and write

$$M_1 \succ M_2.$$

Equivalently, we can say mean  $M_2$  is asymptotically smaller than mean  $M_1$  and write

$$M_2 \prec M_1.$$

We can easily see that asymptotic inequality is a necessary relation between two comparable means.

THEOREM 3.2. *If  $M_1 \geq M_2$ , then  $M_1 \succ M_2$ .*

*Proof.* For  $x$  large enough, the sign of the difference  $M_1(x + s, x + t) - M_2(x + s, x + t)$  is the same as the sign of the first term in its asymptotic expansion.  $\square$

Consider the asymptotic expansion of linear combination of three means:

$$M_2(x - \beta, x + \beta) - (1 - \mu)M_1(x - \beta, x + \beta) - \mu M_3(x - \beta, x + \beta) \sim$$

$$c_m(\mu)x^{-m+1} + c_{m+1}(\mu)x^{-m} + c_{m+2}(\mu)x^{-m-1} \dots \tag{3.2}$$

where  $c_j$  equals the corresponding combination of  $j$ -th coefficients of means  $M_i$ , that is

$$c_j(\mu) = M_2^{(j)} - (1 - \mu)M_1^{(j)} - \mu M_3^{(j)}, \quad j \in \mathbb{N}_0, \tag{3.3}$$

and  $c_m$  denotes the first coefficient that is not zero function of  $\mu$ . Since we deal here with comparable means, we shall assume  $M_1 \leq M_2 \leq M_3$ . From the Theorem 3.2 and formula (3.3) follows that  $c_m(\mu)$  is decreasing. Now suppose that mean  $M_2$  is greater than convex combination of means  $M_1$  and  $M_3$ :

$$M_2(s, t) \geq (1 - \mu)M_1(s, t) + \mu M_3(s, t) \tag{3.4}$$

for all values of arguments  $(s, t)$ , and let  $\mu$  be such that  $c_m(\mu) = 0$ . Taking smaller  $\mu$  will decrease  $c_m(\mu)$  to some negative value and the following asymptotic inequality will hold

$$M_2(s, t) < (1 - \mu)M_1(s, t) + \mu M_3(s, t) \tag{3.5}$$

which along with the Theorem 3.2 contradicts the inequality (3.4). Hence,  $\mu$  such that  $c_m(\mu) = 0$  is optimal. Analogous conclusion is made for the reverse inequality in (3.4). In the following table optimal values of parameter  $\mu$  are given.

$H$	$G$	$L$	$A$	$NS$	$C$	$Q$	$N$	$\times \beta^4/x^3$	$\geq 0$
4/7	-1			3/7				11/105	+
3/7		-1		4/7				13/210	+
-1/7			1	-6/7				17/420	[26]
1/8				-1	7/8			17/360	[25]
-2/9				1		-7/9		1/20	[28]
5/12				-1			7/12	17/360	[28]
	-3/4	1		-1/4				1/60	-
	-1/4		1	-3/4				1/15	+
	1/5			-1	4/5			1/45	-
	-1/3			1		-2/3		7/90	[28]
	-5/9			1			-4/9	1/45	[24]
		-1/3	1	-2/3				11/180	+
		1/4		-1	3/4			1/40	-
		-2/5		1		-3/5		19/300	+
		-5/8		1			-3/8	1/120	+
			1/2	-1	1/2			17/360	[23]
			2/3	-1		1/3		1/180	[20]
			5/6	-1			1/6	17/360	[20]
				-1/2	1	-1/2		31/360	+
				-4/5	1		-1/5	17/450	+
				3/5		-1	2/5	29/300	[22]

For example, the fourth row reads as

$$\frac{1}{8}H + \frac{7}{8}C - NS \sim \frac{17}{360}\beta^4x^{-3} + \dots$$

wherefrom it follows

$$\frac{1}{8}H + \frac{7}{8}C \succ NS. \tag{3.6}$$

There are combinations of means in the table for which the real inequality does not hold such as geometric and logarithmic mean and other marked with the  $-$  sign. The reference is given for those inequalities which are proved in literature. Other inequalities, marked with  $+$  are proved using computer algebra system (CAS).

By this procedure we get one side of the double inequality (1.7).

Optimal parameter for the other side of that inequality can be derived from the values of means at  $(0, 1)$ . Because of homogeneity and symmetry of means it suffices to observe relations between means on the line segment  $\{(s, 1 - s) : s \in [0, 1/2]\}$ . Now the main inequality (1.7) is equivalent to

$$\mu \leq \frac{M_2(s, 1 - s) - M_1(s, 1 - s)}{M_3(s, 1 - s) - M_1(s, 1 - s)} \leq \nu \tag{3.7}$$

and the problem reduces to finding infimum and supremum of the function in the middle. For the most of the combinations of classical means mentioned in the introduction, this function appear to be monotone on the  $[0, \frac{1}{2}]$  and takes the minimum and maximum value at the edges of that interval. Therefore, we can impose the condition

$$(1 - \nu)M_1(0, 1) + \nu M_3(0, 1) = M_2(0, 1)$$

which makes sense if the value of  $M_1$  differs from value of  $M_3$  at point  $(0, 1)$ . In the following table we give such parameters. Inequalities with those parameters were verified through CAS and  $+$  sign stands for those which appeared to be true. Some of them were already proved and the references are given in the introduction and in the previous table. Other still wait for analytic proof. To the best of our knowledge the following inequalities, with the best possible constants, have not been proved yet:

$$H \leq G \leq \frac{4}{7}H + \frac{3}{7}NS \tag{3.8}$$

$$H \leq L \leq \frac{3}{2}H + \frac{4}{7}NS \tag{3.9}$$

$$\frac{1}{4}G + \frac{3}{4}NS \leq A \leq (1 - \sigma)G + \sigma NS \tag{3.10}$$

$$\frac{1}{3}L + \frac{2}{3}NS \leq A \leq (1 - \sigma)L + \sigma NS \tag{3.11}$$

$$\frac{2}{5}L + \frac{3}{5}Q \leq NS \leq \frac{\sqrt{2}\sigma - 1}{\sqrt{2}\sigma}L + \frac{1}{\sqrt{2}\sigma}Q \tag{3.12}$$

$$\frac{5}{8}L + \frac{3}{8}N \leq NS \leq \frac{2\sigma - 1}{2\sigma}L + \frac{1}{2\sigma}N \tag{3.13}$$

$$\frac{1}{2}NS + \frac{1}{2}Q \leq C \leq \frac{(3\sqrt{2} - 4)\sigma}{3\sqrt{2}\sigma - 3}NS + \frac{3 - 4\sigma}{3 - 3\sqrt{2}\sigma}Q \tag{3.14}$$

$$\frac{4}{5}NS + \frac{1}{5}N \leq C \leq \frac{2\sigma}{6\sigma - 3}NS + \frac{3 - 4\sigma}{3 - 6\sigma}N \tag{3.15}$$

where  $\sigma = \operatorname{arcsinh}(1)$ .

$H$	$G$	$L$	$A$	$NS$	$C$	$Q$	$N$	$\times \beta^2/x$	$\geq 0$
-1	1			0				$\frac{1}{2}$	+
-1		1		0				$\frac{2}{3}$	+
$1 - \sigma$			-1	$\sigma$				$\frac{7\sigma-6}{6}$	+
$\frac{3-4\sigma}{4\sigma}$				1	$-\frac{3}{4\sigma}$			$\frac{7\sigma-6}{6\sigma}$	+
$\frac{\sqrt{2\sigma}-1}{\sqrt{2\sigma}}$				-1		$\frac{1}{\sqrt{2\sigma}}$		$\frac{9\sqrt{2}-14\sigma}{12\sigma}$	+
$\frac{1-2\sigma}{2\sigma}$				1			$-\frac{1}{2\sigma}$	$\frac{7\sigma-6}{6\sigma}$	+
	-1	1		0				$\frac{1}{6}$	+
	$1 - \sigma$		-1	$\sigma$				$\frac{4\sigma-3}{6}$	+
$\frac{4\sigma-3}{4\sigma}$				-1	$\frac{3}{4\sigma}$			$\frac{15-16\sigma}{24\sigma}$	+
$\frac{\sqrt{2\sigma}-1}{\sqrt{2\sigma}}$				-1		$\frac{1}{\sqrt{2\sigma}}$		$\frac{3\sqrt{2}-4\sigma}{6\sigma}$	+
$\frac{2\sigma-1}{2\sigma}$				-1			$\frac{1}{2\sigma}$	$\frac{9-8\sigma}{12\sigma}$	+
		$1 - \sigma$	-1	$\sigma$				$\frac{3\sigma-2}{6}$	+
		$\frac{4\sigma-3}{4\sigma}$		-1	$\frac{3}{4\sigma}$			$\frac{1-\sigma}{2\sigma}$	+
		$\frac{\sqrt{2\sigma}-1}{\sqrt{2\sigma}}$		-1		$\frac{1}{\sqrt{2\sigma}}$		$\frac{5\sqrt{2}-6\sigma}{12\sigma}$	+
		$\frac{2\sigma-1}{2\sigma}$		-1			$\frac{1}{2\sigma}$	$\frac{4-3\sigma}{6\sigma}$	+
			$\frac{3-4\sigma}{\sigma}$	1	$\frac{3\sigma-3}{\sigma}$			$\frac{7\sigma-6}{6\sigma}$	+
			$\frac{1-\sqrt{2}\sigma}{(\sqrt{2}-1)\sigma}$	1		$\frac{(\sigma-1)}{(\sqrt{2}-1)\sigma}$		$\frac{(2+\sqrt{2})\sigma-3}{6(\sqrt{2}-1)\sigma}$	+
			$\frac{1-2\sigma}{\sigma}$	1			$\frac{\sigma-1}{\sigma}$	$\frac{7\sigma-6}{6\sigma}$	+
				$\frac{(3\sqrt{2}-4)\sigma}{3\sqrt{2}\sigma-3}$	-1	$\frac{3-4\sigma}{3-3\sqrt{2}\sigma}$		$\frac{(8-3\sqrt{2})\sigma-3}{18(\sqrt{2}\sigma-1)}$	+
				$\frac{2\sigma}{6\sigma-3}$	-1		$\frac{3-4\sigma}{3-6\sigma}$	$\frac{7\sigma-6}{18\sigma-9}$	+
				$\frac{(\sqrt{2}-2)\sigma}{2\sigma-1}$		1	$\frac{1-\sqrt{2}\sigma}{2\sigma-1}$	$\frac{3-(5\sqrt{2}-4)\sigma}{12\sigma-6}$	+

For the combination of harmonic and centroidal mean with Neuman-Sándor mean we read from the table:

$$\left(1 - \frac{3}{4\sigma}\right)H + \frac{3}{4\sigma}C < NS. \tag{3.16}$$

Combining (3.6) with (3.16) suggests that the inequality

$$\left(1 - \frac{3}{4\sigma}\right)H + \frac{3}{4\sigma}C \leq NS \leq \frac{1}{8}H + \frac{7}{8}C \tag{3.17}$$

should be true. Expression like this, with sharp inequalities, was proved in [25]. Now some known inequalities are immediate consequence. Inequality (1.7) is equivalent to

$$\mu \leq \frac{M_2(1,t) - M_1(1,t)}{M_3(1,t) - M_1(1,t)} \leq \nu. \tag{3.18}$$

Since the means are symmetric, it suffices to consider the case  $t \geq 1$ . Because of the arc-hyperbolic sine function in the definition of Neuman-Sándor mean it is convenient to make a substitution

$$t = \frac{1 + \sinh \varphi}{1 - \sinh \varphi}, \quad \varphi \in [0, \operatorname{arcsinh}(1)). \tag{3.19}$$

Let

$$M_{HNSC}(\varphi) = \frac{NS(1,t) - H(1,t)}{C(1,t) - H(1,t)} \tag{3.20}$$

with  $t$  defined above. Now we have

$$M_{HNSC}(\varphi) = \frac{\frac{\sinh \varphi}{\varphi(1-\sinh \varphi)} - (\sinh \varphi + 1)}{\frac{\sinh \varphi + 3}{3(1-\sinh \varphi)} - (\sinh \varphi + 1)} = \frac{3}{4} \left( 1 + \frac{1}{\varphi \sinh \varphi} - \frac{1}{\sinh^2 \varphi} \right). \tag{3.21}$$

Denote

$$M(\varphi) = 1 + \frac{1}{\varphi \sinh \varphi} - \frac{1}{\sinh^2 \varphi} = \frac{4}{3} M_{HNSC}(\varphi) \tag{3.22}$$

Then result from [25]:

$$\frac{3}{4\sigma} < M_{HNSC}(\varphi) < \frac{7}{8}, \quad \varphi \in (0, \operatorname{arcsinh}(1)) \tag{3.23}$$

implies

$$\frac{1}{\sigma} < M(\varphi) < \frac{7}{6}, \quad \varphi \in (0, \operatorname{arcsinh}(1)). \tag{3.24}$$

In the same manner we define functions

$$M_{HANS}(\varphi) = \frac{A(1,t) - H(1,t)}{NS(1,t) - H(1,t)} = \frac{1}{M(\varphi)}, \tag{3.25}$$

$$M_{HNSN}(\varphi) = \frac{NS(1,t) - H(1,t)}{N(1,t) - H(1,t)} = \frac{1}{2} M(\varphi), \tag{3.26}$$

$$M_{ANSC}(\varphi) = \frac{NS(1,t) - A(1,t)}{C(1,t) - A(1,t)} = 3(M(\varphi) - 1), \tag{3.27}$$

$$M_{ANSN}(\varphi) = \frac{NS(1,t) - A(1,t)}{N(1,t) - A(1,t)} = M(\varphi) - 1, \tag{3.28}$$

$$M_{NSCN}(\varphi) = \frac{C(1,t) - NS(1,t)}{N(1,t) - NS(1,t)} = \frac{4 - 3M(\varphi)}{6 - 3M(\varphi)}. \tag{3.29}$$



Inequality (1.5), stated as Theorem 1.3. in [28], follow easily since (3.26) and (3.24) yield

$$\frac{1}{2\sigma} < M_{HNSN}(\varphi) < \frac{7}{12}$$

and further

$$\frac{1}{2\sigma}N(1,t) + \left(1 - \frac{1}{2\sigma}\right)H(1,t) < NS(1,t) < \frac{7}{12}N(1,t) + \frac{5}{12}H(1,t).$$

For the arithmetic and centroidal mean we have

$$3\left(\frac{1}{\sigma} - 1\right) < M_{ANSC}(\varphi) < \frac{1}{2}$$

and

$$3\left(\frac{1}{\sigma} - 1\right)C(1,t) + \left(4 - \frac{3}{\sigma}\right)A(1,t) < NS(1,t) < \frac{1}{2}C(1,t) + \frac{1}{2}A(1,t).$$

Hence, the inequality (1.6) holds.

Similarly,

$$\frac{1}{\sigma} - 1 < M_{ANSN}(\varphi) < \frac{1}{6}$$

gives

$$\left(\frac{1}{\sigma} - 1\right)N(1,t) + \left(2 - \frac{1}{\sigma}\right)A(1,t) < NS(1,t) < \frac{1}{6}N(1,t) + \frac{5}{6}A(1,t)$$

which was proved in [20] as Theorem 3.2. Theorem 6 in [26] is a consequence of (3.25) and (3.24).

Finally, by combining (3.24) and (3.29) we obtain new double inequality for centroidal, Neuman-Sándor and contraharmonic mean.

**THEOREM 3.3. Inequality**

$$(1 - \mu)NS(s,t) + \mu N(s,t) < C(s,t) < (1 - \nu)NS(s,t) + \nu N(s,t)$$

holds for all  $s, t > 0$ , with  $s \neq t$ , if and only if  $\mu \leq \frac{1}{5}$  and  $\nu \geq \frac{4\sigma-3}{6\sigma-3}$ .

**4. Reciprocal of means**

Besides the classical combinations we can observe convex combinations of reciprocal of means:

$$\mu \frac{1}{M_1} + (1 - \mu) \frac{1}{M_3} \leq \frac{1}{M_2} \leq \nu \frac{1}{M_1} + (1 - \nu) \frac{1}{M_3} \tag{4.1}$$

which cover Neuman-Sándor mean. If the mean  $M$  has asymptotic expansion of the form (2.1) with  $\alpha = 0$ , then by Lemma 2.1 reciprocal of the mean  $M$  has the following asymptotic expansion

$$M(x - \beta, x + \beta) \sim x - c_2x^{-1} + (c_2^2 - c_4)x^{-3} + \dots$$

As a consequence, we can find optimal parameter for the one side of the inequality (1.7) in the first table. Same as before, observing linear combinations near point  $(0, 1)$  will should provide optimal parameter for the other side.

In this case some new inequalities can be proved.

**THEOREM 4.1. Inequality**

$$\mu \frac{1}{H(s,t)} + (1 - \mu) \frac{1}{NS(s,t)} < \frac{1}{A(s,t)} < \nu \frac{1}{H(s,t)} + (1 - \nu) \frac{1}{NS(s,t)} \tag{4.2}$$

holds for all  $s, t \in \langle 0, \infty \rangle$ ,  $s \neq t$ , if and only if  $\mu = 0$  and  $\frac{1}{7} \leq \nu \leq 1$ .

*Proof.* Similarly as before, define function  $R(\varphi)$

$$R(\varphi) = \frac{1/A(1,t) - 1/NS(1,t)}{1/H(1,t) - 1/NS(1,t)}, \quad \varphi \in \langle 0, \sigma \rangle$$

and  $t$  is defined by (3.19).  $R(\varphi)$  after arranging becomes

$$R(\varphi) = \frac{(\sinh \varphi - \varphi)(1 - \sinh^2 \varphi)}{-\varphi + \sinh \varphi + \varphi \sinh^2 \varphi} = 1 - \frac{1}{\frac{\varphi}{\sinh \varphi} M(\varphi)}$$

It is easily seen that  $\frac{\varphi}{\sinh \varphi}$  takes values between  $\sigma$  and 1 which together with (3.24) gives

$$0 < R(\varphi) < \frac{1}{7}.$$

Optimality of the parameters is assured by discussion before the theorem.  $\square$

**THEOREM 4.2. Inequality**

$$\mu \frac{1}{H(s,t)} + (1 - \mu) \frac{1}{N(s,t)} < \frac{1}{NS(s,t)} < \nu \frac{1}{H(s,t)} + (1 - \nu) \frac{1}{N(s,t)} \tag{4.3}$$

holds for all  $s, t \in \langle 0, \infty \rangle$ , with  $s \neq t$ , if and only if  $\mu = 0$  and  $\frac{5}{12} \leq \nu \leq 1$ .

*Proof.* Let

$$R(\varphi) = \frac{1/NS(1,t) + 1/N(1,t)}{1/H(1,t) - 1/N(1,t)}, \quad \varphi \in \langle 0, \sigma \rangle \tag{4.4}$$

where  $t$  is defined by (3.19). It is easily seen that

$$\lim_{\varphi \rightarrow 0} R(\varphi) = \frac{5}{12} \text{ and } \lim_{\varphi \rightarrow \sigma} R(\varphi) = 0. \tag{4.5}$$

It remains to prove  $R$  is decreasing.

$$\begin{aligned} R'(\varphi) &= \frac{1}{2 \sinh^4(\varphi)} (\sinh \varphi - \sinh^5 \varphi + \sinh(2\varphi) - \varphi \cosh \varphi (3 + \sinh^4 \varphi)) \\ &= \frac{1}{2 \sinh^4(\varphi)} g(\varphi) \end{aligned}$$

Taylor series for the function  $g$  equals

$$\begin{aligned} g(\varphi) &= \frac{3}{8} \sinh \varphi + \sinh(2\varphi) + \frac{5}{16} \sinh(3\varphi) - \frac{1}{16} \sinh(5\varphi) \\ &\quad - \varphi \left[ \frac{25}{8} \cosh \varphi - \frac{3}{16} \cosh(3\varphi) + \frac{1}{16} \cosh(5\varphi) \right] \\ &= \sum_{n=0}^{\infty} \left[ \frac{3}{8} + 2^{2n+1} + \frac{5}{16} 3^{2n+1} - \frac{1}{16} 5^{2n+1} \right. \\ &\quad \left. - (2n+1) \left( \frac{25}{8} - \frac{3}{16} 3^{2n} + \frac{1}{16} 5^{2n} \right) \right] \frac{\varphi^{2n+1}}{(2n+1)!} \end{aligned}$$

The first two coefficients of this series is equal to zero, and for  $n \geq 2$  we have

$$\begin{aligned} &6 + 32 \cdot 2^{2n} + 15 \cdot 3^{2n} - 5 \cdot 5^{2n} \\ &< 6 + 32 \cdot 2^{2n} + 15 \cdot 3^{2n} - 5 \cdot 2^{2n} - 15n \cdot 2^{2n} - \frac{20n}{3} \cdot 3^{2n} - 5 \cdot 3^{2n} < 0 \end{aligned}$$

and also

$$50 - 3 \cdot 3^{2n} + 5^{2n} > 0$$

which makes all coefficients negative. Hence,  $R$  is strictly decreasing and the statement of theorem is proved.  $\square$

*Acknowledgement.* The authors would thank anonymous referees for their useful remarks which improve presentation of this paper.

#### REFERENCES

- [1] J. ACZEL, Z. PÁLES, *The behaviour of means under equal increments of their variables*, Amer. Math. Monthly, **95**, (1988), 856–860.
- [2] P. S. BULLEN, *Averages still on the move*, Math. Mag., **63** (1990), 250–255.
- [3] P. S. BULLEN, *Handbook of Means and Their Inequalities*, Kluwer Academic Publishers, Dordrecht, 2003.
- [4] P. S. BULLEN, D. S. MITRINOVIĆ, P. M. VASIĆ, *Means and theirs inequalities*, D Reidel, Dordrecht, 1988.
- [5] T. BURIĆ, N. ELEZOVIĆ, *Bernoulli polynomials and asymptotic expansions of the quotient of gamma functions*, J. Comput. Appl. Math., **235**, 11 (2011), 3315–3331.
- [6] T. BURIĆ, N. ELEZOVIĆ, *New asymptotic expansions of the gamma function and improvements of Stirling's type formulas*, J. Comput. Anal. Appl., **13**, 4 (2011), 785–795.
- [7] T. BURIĆ, N. ELEZOVIĆ, *New asymptotic expansions of the quotient of gamma functions*, Integral Transforms Spec. Funct., **23** (2012), 355–368.

- [8] T. BURIĆ, N. ELEZOVIĆ, *Approximations of the Euler-Mascheroni constant and harmonic numbers*, Appl. Math. Comput., **222** (2013), 604–611.
- [9] T. BURIĆ, N. ELEZOVIĆ, *Asymptotic expansions of the binomial coefficients*, J. Appl. Math. Comput., **46** (2014), 135–145.
- [10] C.-P. CHEN, N. ELEZOVIĆ, L. VUKŠIĆ, *Asymptotic expansion of integral mean of polygamma functions* Math. Inequal. Appl. **18**, 1 (2015), 255–266.
- [11] N. ELEZOVIĆ, *Asymptotic inequalities and comparison of classical means*, J. Math. Inequal., **9**, 1 (2015), 177–196.
- [12] N. ELEZOVIĆ, *Asymptotic expansions of gamma and related functions, binomial coefficients, inequalities and means*, J. Math. Inequal **9**, 4 (2015), 1001–1054.
- [13] N. ELEZOVIĆ, L. VUKŠIĆ, *Asymptotic expansions of integral means and applications to the ratio of gamma functions*, Appl. Math. Comput. **235** (2014), 187–200.
- [14] N. ELEZOVIĆ, L. VUKŠIĆ, *Asymptotic expansions of bivariate classical means and related inequalities*, J. Math. Inequal., **8**, 4 (2014), 707–724.
- [15] N. ELEZOVIĆ, L. VUKŠIĆ, *Asymptotic expansions and comparison of bivariate parameter means*, Math. Inequal. Appl., **17**, 4 (2014), 1225–1244.
- [16] W.-M. GONG, X.-H. SHEN, Y.-M. CHU, *Bounds for the Neuman-Sándor mean in terms of logarithmic, quadratic or contraharmonic means*, Int. Math. Forum, **8**, 30 (2013), 1467–1475.
- [17] L. HOEHN, I. NIVEN, *Averages on the move*, Math. Mag., **58** (1985), 151–156.
- [18] W.-D. JIANG, *Some sharp inequalities involving reciprocals of the Seiffert and other means*, J. Math. Inequal., **6**, 4 (2012), 593–599.
- [19] D. S. MITRINOVIĆ, J. E. PEČARIĆ, A. M. FINK, *Classical and New Inequalities in Analysis*, D Reidel, Dordrecht, 1993.
- [20] E. NEUMAN, *A note on certain bivariate mean*, J. Math. Inequal., **6**, 4 (2012), 637–643.
- [21] E. NEUMAN, J. SÁNDOR, *On the Schwab-Borchardt mean*, Mathematica Pannonica, **14**, 2 (2003), 253–266.
- [22] FENG QI, WE-HUI LI, *A unified proof of several inequalities and some new inequalities involving Neuman-Sándor mean*, Miskolc Math. Notes, **15**, 2 (2014), 665–675
- [23] W.-M. QIAN, Y.-M. CHU, *On certain inequalities for Neuman-Sándor mean*, Abstr. Appl. Anal., **2013**, Article ID 790783, 6 pages.
- [24] H. SUN, X.-H. SHEN, T.-H. ZHAO, Y.-M. CHU, *Optimal bounds for the Neuman-Sándor means in terms of geometric and contraharmonic means*, Appl. Math. Sci., **7**, 88 (2013), 4363–4373.
- [25] W. XIA, Y. CHU, *Optimal inequalities between Neuman-Sándor, centroidal and harmonic means*, J. Math. Inequal., **7**, 4 (2013), 593–600.
- [26] F. ZHANG, Y.-M. CHU, W.-M. QIAN, *Bounds for the arithmetic mean in terms of the Neuman-Sándor and other bivariate means*, J. Appl. Math., **2013**, Article ID 582504, 7 pages.
- [27] T.-H. ZHAO, Y.-M. CHU, Y.-L. JIANG, Y.-M. LI, *Best possible bounds for Neuman-Sándor mean by the identric, quadratic and contraharmonic means*, Abstr. Appl. Anal., **2013**, Article ID 348326, 12 pages.
- [28] T.-H. ZHAO, Y.-M. CHU, B.-Y. LIU, *Optimal bounds for Neuman-Sándor mean in terms of the convex combinations of harmonic, geometric, quadratic, and contraharmonic means*, Abstr. Appl. Anal., **2012**, Article ID 302635, 9 pages.

(Received July 10, 2014)

*Neven Elezović*  
 Faculty of Electrical Engineering and Computing  
 University of Zagreb  
 Unska 3, 10000 Zagreb, Croatia  
 e-mail: neven.elez@fer.hr  
  
*Lenka Vukšić*  
 Faculty of Electrical Engineering and Computing  
 University of Zagreb  
 Unska 3, 10000 Zagreb, Croatia  
 e-mail: lenka.vuksic@fer.hr