

GENERALIZED STEFFENSEN'S INEQUALITY

ASFAND FAHAD, JOSIP PEČARIĆ AND MARJAN PRALJAK

(Communicated by N. Elezović)

Abstract. We give a generalization of Steffensen's inequality by extending the results of Pečarić [4] and Rabier [5]. We make use of the n -order Taylor expansion of a composition of functions and Faà di Bruno's formula for higher order derivatives of the composition.

1. Introduction

Steffensen [6] proved the following inequality: if $f, h : [\alpha, \beta] \rightarrow \mathbb{R}$, $0 \leq h \leq 1$ and f is decreasing, then

$$\int_{\alpha}^{\beta} f(t)h(t) dt \leq \int_{\alpha}^{\alpha+\gamma} f(t) dt, \quad \text{where } \gamma = \int_{\alpha}^{\beta} h(t) dt. \quad (1)$$

A few hundred papers are devoted to studying generalizations of Steffensen's inequality (1). One recent is given by Rabier [5].

THEOREM 1.1. *Let $\phi : [0, \infty) \rightarrow \mathbb{R}$ be convex and continuous with $\phi(0) = 0$. If $b > 0$ and $h \in L^{\infty}(0, b)$, $h \geq 0$ and $\|h\|_{\infty} \leq 1$, then $h\phi^{(1)} \in L^1(0, b)$ and*

$$\phi\left(\int_0^b h(t) dt\right) \leq \int_0^b h(t)\phi^{(1)}(t) dt \quad (2)$$

In fact, Rabier's result is closely related to another generalization of Steffensen's inequality given by Pečarić [4].

THEOREM 1.2. *Let $g : [a, b] \rightarrow \mathbb{R}$ be a nondecreasing and differentiable function and $f : I \rightarrow \mathbb{R}$ be a nondecreasing function (I is an interval in \mathbb{R} such that $a, b, g(a), g(b) \in I$).*

(a) *If $g(x) \leq x$, then*

$$\int_a^b f(t)g^{(1)}(t) dt \geq \int_{g(a)}^{g(b)} f(t) dt. \quad (3)$$

Mathematics subject classification (2010): 26D10, 26D15.

Keywords and phrases: Steffensen's inequality, Faà di Bruno's formula.

This research work is funded by Higher Education Commission Pakistan. The research of the second author was supported by the Croatian Ministry of Science, Education and Sports under the Research Grants 117- 1170889-0888. The research of the third author was supported by the Croatian Ministry of Science, Education and Sports under the Research Grant 202339.

(b) If $g(x) \geq x$, then the reverse of the above inequality holds.

REMARK 1.3. The assumptions of Theorem 1.2 can be weakened and differentiability of g can be replaced with absolute continuity. Indeed, for a nondecreasing function f , the function $F(x) = \int_a^x f(t) dt$ is well defined and satisfies $F'(x) = f(x)$ at all points except at most countably many points. For absolutely continuous nondecreasing function g the substitution $z = g(t)$ in the integral is justified (see [2, Corollary 20.5]), so

$$F(g(b)) - F(g(a)) = \int_{g(a)}^{g(b)} f(z) dz = \int_a^b f(g(t))g'(t) dt \leq \int_a^b f(t)g'(t) dt, \quad (4)$$

where the last inequality holds when $g(t) \leq t$.

Steffensen's inequality (1) follows from Theorem 1.2 by making substitutions $g(x) \mapsto \int_a^x h(t + \alpha - a) dt + a$ and $f(x) \mapsto -f(x + \alpha - a)$ and taking $b = \beta - \alpha + a$.

REMARK 1.4. Theorem 1.1 follows from Theorem 1.2 with weaker assumptions. Indeed, the convex function ϕ from Theorem 1.1 has a nondecreasing right-sided derivative $\phi_+^{(1)}$ such that $\phi(x) = \int_0^x \phi_+^{(1)}(t) dt$. Furthermore, for a function $h : [0, b] \rightarrow [0, 1]$, the function $g(x) = \int_0^x h(t) dt$ is absolutely continuous and satisfies $g(x) \leq x$ and $g^{(1)} = h$ almost everywhere. Therefore, by taking $a = 0$, $f = \phi_+^{(1)}$ and $g(x) = \int_0^x h(t) dt$ in Theorem 1.2 (under the weaker assumptions) we get Theorem 1.1.

On the other hand, an absolutely continuous nondecreasing function $g : [0, b] \rightarrow \mathbb{R}$ can satisfy $g(x) \leq x$ without satisfying $0 \leq g^{(1)}(x) \leq 1$.

The goal of this paper is to generalize the inequality from Theorem 1.2 by replacing the equality

$$F(g(x)) = F(g(a)) + \int_{g(a)}^{g(x)} f(t) dt$$

with the n -th order Taylor expansion of the composition $F \circ g$ with the remainder given in the integral form. In the process we will use Faà di Bruno's formula for higher order derivatives of the composition $F \circ g$. The formula states that

$$\frac{d^m}{dx^m} F(g(x)) = \sum_{k=1}^m F^{(k)}(g(x)) B_{m,k}(g^{(1)}(x), \dots, g^{(m-k+1)}(x)) \quad (5)$$

where $B_{m,k}(x_1, x_2, \dots, x_{m-k+1})$ are the Bell polynomials

$$B_{m,k}(x_1, x_2, \dots, x_{m-k+1}) = \sum \frac{m!}{j_1! j_2! \dots j_{m-k+1}!} \left(\frac{x_1}{1!}\right)^{j_1} \left(\frac{x_2}{2!}\right)^{j_2} \dots \left(\frac{x_{m-k+1}}{(m-k+1)!}\right)^{j_{m-k+1}}$$

where the sum is taken over all sequences $j_1, j_2, \dots, j_{m-k+1}$ of non-negative integers such that

$$j_1 + j_2 + \dots + j_k = k \quad \text{and} \quad j_1 + 2j_2 + 3j_3 + \dots = m.$$

For a historical overview of Faà di Bruno's formula and its various forms see [3].

In this paper we will give a generalization of Steffensen's inequality by making use of the n -order Taylor expansion of the composition $F \circ g$. Moreover, we will give the inequalities for the special case when g is of the form $g(x) = \int_0^x h(t) dt$ and use it to obtain a Hardy-type inequality.

2. Main results

The following theorem states our main result.

THEOREM 2.1. *Let $n \in \mathbb{N}$. Let $g : [a, b] \rightarrow \mathbb{R}$ and $F : I \rightarrow \mathbb{R}$ (where I is an interval in \mathbb{R} such that $a, b, g(a), g(b) \in I$) be two n times differentiable functions such that $g, g^{(1)}, \dots, g^{(n-1)}, F^{(1)}, F^{(2)}, \dots, F^{(n)}$ are nondecreasing functions.*

(a) *If $g(x) \leq x$, then*

$$F(g(b)) \leq F(g(a)) + \sum_{k=1}^{n-1} F^{(k)}(g(a)) \sum_{i=k}^{n-1} B_{i,k}(g^{(1)}(a), \dots, g^{(i-k+1)}(a)) \frac{(b-a)^i}{i!} + \int_a^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(t) B_{n,k}(g^{(1)}(t), \dots, g^{(n-k+1)}(t)) dt.$$

(b) *If $g(x) \geq x$, then the reverse of the above inequality holds.*

Proof. The $(n - 1)$ -th Taylor expansion of a function H with the remainder in the integral form is given by

$$H(x) = \sum_{k=0}^{n-1} H^{(k)}(a) \frac{(x-a)^k}{k!} + \int_a^x H^{(n)}(t) \frac{(x-t)^{n-1}}{(n-1)!} dt$$

Applying this formula for the composition $H = F \circ g$ and using Faà di Bruno's formula (5) gives

$$\begin{aligned} (F \circ g)(b) &= (F \circ g)(a) + \sum_{i=1}^{n-1} \sum_{k=1}^i F^{(k)}(g(a)) B_{i,k}(g^{(1)}(a), \dots, g^{(i-k+1)}(a)) \frac{(b-a)^i}{i!} \\ &\quad + \int_a^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(g(t)) B_{n,k}(g^{(1)}(t), \dots, g^{(n-k+1)}(t)) dt \\ &= (F \circ g)(a) + \sum_{k=1}^{n-1} F^{(k)}(g(a)) \sum_{i=k}^{n-1} B_{i,k}(g^{(1)}(a), \dots, g^{(i-k+1)}(a)) \frac{(b-a)^i}{i!} \\ &\quad + \int_a^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(g(t)) B_{n,k}(g^{(1)}(t), \dots, g^{(n-k+1)}(t)) dt. \end{aligned}$$

By the assumptions of the theorem, $g^{(i)} \geq 0$ for $i = 1, \dots, n$, so the Bell polynomials evaluated at the derivatives of g in the above expression are nonnegative. Therefore, for $g(x) \leq x$ the inequality

$$\int_a^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(g(t)) B_{n,k}(g^{(1)}(t), \dots, g^{(n-k+1)}(t)) dt \leq \int_a^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(t) B_{n,k}(g^{(1)}(t), \dots, g^{(n-k+1)}(t)) dt$$

holds, while for $g(x) \geq x$ the reversed inequality holds. \square

A special case of the previous theorem is given by the following corollary.

COROLLARY 2.2. *Let $n \in \mathbb{N}$, let $F : [0, b] \rightarrow \mathbb{R}$ be n times differentiable function such that $F^{(1)}, F^{(2)}, \dots, F^{(n)}$ are nondecreasing functions and let $h : [0, b] \rightarrow [0, +\infty)$ be $n - 1$ times differentiable function such that $h, h^{(1)}, \dots, h^{(n-1)}$ are nonnegative.*

(a) *If $\int_0^x h(t) dt \leq x$ for every $x \in [0, b]$, then*

$$F\left(\int_0^b h(t) dt\right) \leq F(0) + \sum_{k=1}^{n-1} F^{(k)}(0) \sum_{i=k}^{n-1} B_{i,k}(h(0), h^{(1)}(0), \dots, h^{(i-k)}(0)) \frac{b^i}{i!} + \int_0^b \frac{(b-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(t) B_{n,k}(h(t), h^{(1)}(t), \dots, h^{(n-k)}(t)) dt.$$

(b) *If $x \leq \int_0^x h(t) dt$ for every $x \in [0, b]$, then the reverse of the above inequality holds.*

Proof. Follows from Theorem 2.1 by taking $a = 0$ and $g(x) = \int_0^x h(t) dt$. \square

By inserting the Bell polynomials we get the explicit forms of the inequalities from Theorem 2.1 and Corollary 2.2. Since $B_{1,1}(x_1) = x_1$, the inequalities (from part (a)) for $n = 1$ are given by (4) and

$$F\left(\int_0^b h(t) dt\right) \leq F(0) + \int_0^b F^{(1)}(t) h(t) dt.$$

The subsequent Bell polynomials are equal to:

$$n = 2 : B_{2,1}(x_1, x_2) = x_2, \quad B_{2,2}(x_1) = x_1^2,$$

$$n = 3 : B_{3,1}(x_1, x_2, x_3) = x_3, \quad B_{3,2}(x_1, x_2) = 3x_1x_2, \quad B_{3,3}(x_1) = x_1^3,$$

$$n = 4 : B_{4,1}(x_1, x_2, x_3, x_4) = x_4, \quad B_{4,2}(x_1, x_2, x_3) = 3x_1^2x_3 + 3x_2^2, \\ B_{4,3}(x_1, x_2) = 6x_1^2x_2, \quad B_{4,4}(x_1) = x_1^4,$$

$$n = 5 : B_{5,1}(x_1, x_2, x_3, x_4, x_5) = x_5, \quad B_{5,2}(x_1, x_2, x_3, x_4) = 5x_1x_4 + 10x_2x_3, \\ B_{5,3}(x_1, x_2, x_3) = 10x_1^2x_3 + 15x_1x_2^2, \\ B_{5,4}(x_1, x_2) = 10x_1^3x_2, \quad B_{5,5}(x_1) = x_1^5.$$

Therefore, for $n = 2$ the inequality from Theorem 2.1 (a) is of the form

$$F(g(b)) \leq F(g(a)) + (b - a)F^{(1)}(g(a))g^{(1)}(a) + \int_a^b (b - t) \left[F^{(1)}(t)g^{(2)}(t) + F^{(2)}(t)g^{(1)}(t)^2 \right] dt$$

and the inequality from Corollary 2.2(a) is of the form

$$F \left(\int_0^b h(t) dt \right) \leq F(0) + bF^{(1)}(0)h(0) + \int_0^b (b - t) \left[F^{(1)}(t)h^{(1)}(t) + F^{(2)}(t)h(t)^2 \right] dt.$$

Next, we will use the inequality from Corollary 2.2 to obtain the following result.

THEOREM 2.3. *Let $n \in \mathbb{N}$, h and F be as in Corollary 2.2, $k : [0, b] \rightarrow [0, +\infty)$ and denote $K_i(t) = \int_t^b \frac{(x-t)^{i-1}}{(i-1)!} k(x) dx$ for $i \in \mathbb{N}$.*

(a) *If $\int_0^x h(t) dt \leq x$ for every $x \in [0, b]$, then*

$$\begin{aligned} & \int_0^b k(x) F \left(\int_0^x h(t) dt \right) dx \\ & \leq F(0)K_1(0) + \sum_{k=1}^{n-1} F^{(k)}(0) \sum_{i=k}^{n-1} B_{i,k}(h(0), h^{(1)}(0), \dots, h^{(i-k)}(0)) K_{i+1}(0) \\ & \quad + \int_0^b K_n(t) \sum_{k=1}^n F^{(k)}(t) B_{n,k}(h(t), h^{(1)}(t), \dots, h^{(n-k)}(t)) dt. \end{aligned}$$

(b) *If $x \leq \int_0^x h(t) dt$ for every $x \in [0, b]$, then the reverse of the above inequality holds.*

Proof. Applying Corollary 2.2 with $b = x$ and multiplying by $k(x) \geq 0$ we get

$$\begin{aligned} & k(x) F \left(\int_0^x h(t) dt \right) \\ & \leq F(0)k(x) + \sum_{k=1}^{n-1} F^{(k)}(0) \sum_{i=k}^{n-1} B_{i,k}(h(0), h^{(1)}(0), \dots, h^{(i-k)}(0)) \frac{x^i}{i!} k(x) \\ & \quad + k(x) \int_0^x \frac{(x-t)^{n-1}}{(n-1)!} \sum_{k=1}^n F^{(k)}(t) B_{n,k}(h(t), h^{(1)}(t), \dots, h^{(n-k)}(t)) dt. \end{aligned}$$

Integrating the above inequality with respect to x from 0 to b and applying Fubini's theorem on the right hand side we obtain the stated inequality. \square

For $p > 1$, $l > 1$ and a nonnegative function h such that $x^{1-p/l} h \in L^p(0, b)$, inequality

$$\int_0^b x^{-l} \left(\int_0^x h(t) dt \right)^p dx \leq \left(\frac{p}{l-1} \right)^p \int_0^b x^{p-l} h(x)^p dx \tag{6}$$

holds. The classical Hardy’s inequality is inequality (6) with $b = \infty$ and in that case the constant in (6) is sharp, while for finite $b > 0$ the constant is not sharp. In fact, for finite b the following inequality due to Čižmešija and Pečarić [1] holds

$$\int_0^b x^{-l} \left(\int_0^x h(t) dt \right)^p dx \leq \left(\frac{p}{l-1} \right)^p \int_0^b \left[1 - \left(\frac{x}{b} \right)^{\frac{l-1}{p}} \right] x^{p-l} h(x)^p dx \tag{7}$$

and the constant in (7) is sharp. For $l = p$ inequality (6) can be written as

$$\|Ah\|_p \leq \frac{p}{p-1} \|h\|_p,$$

where A is the operator $Ah(x) = \frac{1}{x} \int_0^x h(t) dt$. In the following example we will use results obtained in this paper to derive some inequalities similar to (6) and (7).

EXAMPLE 2.4. Applying Theorem 2.3(a) with $F(t) = t^p$, $p > n$, we obtain the inequality

$$\int_0^b k(x) \left(\int_0^x h(t) dt \right)^p dx \leq \int_0^b K_n(t) \sum_{k=1}^n (p)_k t^{p-k} B_{n,k}(h(t), h^{(1)}(t), \dots, h^{(n-k)}(t)) dt, \tag{8}$$

where $(p)_k = p(p-1)\dots(p-k+1)$ is the Pochhammer symbol.

Furthermore, for $p > 1$

$$\begin{aligned} \int_0^b k(x) \left(\int_0^x h(t) dt \right)^p dx &\leq p \int_0^b t^{p-1} K(t) h(t) dt \\ &\leq p \left[\int_0^b h(t)^p dt \right]^{\frac{1}{p}} \left[\int_0^b t^p K(t)^{\frac{p}{p-1}} dt \right]^{\frac{p-1}{p}}, \end{aligned}$$

where the first inequality follows from (8) with $n = 1$ and the second inequality follows by applying Hölder’s inequality. In particular, for $k(x) = x^{-p}$ we have

$$\begin{aligned} \int_0^b \left(\frac{1}{x} \int_0^x h(t) dt \right)^p dx &\leq \frac{p}{p-1} \int_0^b \left(1 - \left(\frac{t}{b} \right)^{p-1} \right) h(t) dt \\ &\leq \frac{p}{p-1} \left[\int_0^b h(t)^p dt \right]^{\frac{1}{p}} \left[\int_0^b \left(1 - \left(\frac{t}{b} \right)^{p-1} \right)^{\frac{p}{p-1}} dt \right]^{\frac{p-1}{p}}. \end{aligned} \tag{9}$$

The last inequality can be written as

$$\|Ah\|_p^p \leq C \|h\|_p^p,$$

where the constant C does not depend on h . By applying the substitution $y = (t/b)^{p-1}$ in the last integral on the right hand side of (9), one can calculate

$$C = \frac{pb}{(p-1)^2} B \left(\frac{1}{p-1}, \frac{2p-1}{p-1} \right),$$

where B is the beta function. In particular, for $p = 2$ we have

$$\int_0^b \left(\frac{1}{x} \int_0^x h(t) dt \right)^2 dx \leq \frac{2}{3} b \|h\|_2.$$

REFERENCES

- [1] A. ČIŽMEŠIJA AND J. PEČARIĆ, *Some new generalizations of inequalities of Hardy and Levin-Cochran-Lee*, Bull. Aust. Math. Soc. **63**, 1 (2001), 105–113.
- [2] E. HEWITT AND K. STROMBERG, *Real and abstract analysis*, 3rd edition, Springer, New York, 1975.
- [3] W. P. JOHNSON, *The Curious History of Faà di Bruno's Formula*, Amer. Math. Monthly **109** (2002), 217–234.
- [4] J. PEČARIĆ, *Connections among some inequalities of Gauss, Steffensen and Ostrowski*, Southeast Asian Bull. Math. **13**, 2 (1989), 89–91.
- [5] P. RABIER, *Steffensen's inequality and $L^1 - L^\infty$ estimates of weighted integrals*, Proc. Amer. Math. Soc. **140**, 2 (2012), 665–675.
- [6] J. F. STEFFENSEN, *On certain inequalities between mean values, and their application to actuarial problems*, Skand. Aktuarietidskr. **1** (1918), 82–97.

(Received May 22, 2013)

Asfand Fahad
Abdus Salam School of Mathematical Sciences, GC University
Lahore, Pakistan
e-mail: asfandfahad1@yahoo.com

Josip Pečarić
University Of Zagreb, Faculty Of Textile Technology
Zagreb, Croatia
and

Abdus Salam School of Mathematical Sciences, GC University
Lahore, Pakistan
e-mail: pecaric@mahazu.hazu.hr

Marjan Praljak
University Of Zagreb, Faculty of Food Technology and Biotechnology
Zagreb, Croatia
and

Abdus Salam School of Mathematical Sciences, GC University
Lahore, Pakistan
e-mail: mpraljak@pbf.hr