

EXTENSION OF DETERMINANTAL INEQUALITIES OF POSITIVE DEFINITE MATRICES

XIAOHUI FU, YANG LIU AND SHUNQIN LIU

(Communicated by J. Pečarić)

Abstract. In this short note, we extend some known determinantal inequalities of positive definite matrices to a larger class of matrices, namely, matrices whose numerical range is contained in a sector.

1. Introduction

Let $\mathbb{M}_n(\mathbb{C})$ denote the set of $n \times n$ complex matrices. For $A \in \mathbb{M}_n(\mathbb{C})$, the conjugate transpose of A is denoted by A^* , and recall the Cartesian decomposition ([8, p. 6]) $A = \Re A + i\Im A$, where $\Re A = \frac{1}{2}(A + A^*)$ and $\Im A = \frac{1}{2i}(A - A^*)$. For two Hermitian matrices $A, B \in \mathbb{M}_n(\mathbb{C})$, $A \prec B$ ($A \preceq B$) means that $B - A$ is positive definite (semidefinite). In particular, a positive definite (positive semidefinite) matrix A can be expressed as $A \succ 0$ ($A \succeq 0$). We also consider $A \in \mathbb{M}_n(\mathbb{C})$ to be partitioned as

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}, \quad (1.1)$$

where diagonal blocks are square matrices. $\text{diag}(D_1, \dots, D_k)$ denotes the block diagonal matrix whose diagonal blocks are D_1, \dots, D_k . In (1.1), if A_{11} is nonsingular, the Schur complement of A_{11} in A is defined by $A/A_{11} = A_{22} - A_{21}A_{11}^{-1}A_{12}$.

The numerical range of $A \in \mathbb{M}_n(\mathbb{C})$ is defined by

$$W(A) = \{x^*Ax \mid x \in \mathbb{C}^n, x^*x = 1\}.$$

For $\alpha \in [0, \frac{\pi}{2})$, let S_α be the sector in the complex plane given by

$$S_\alpha = \{z \in \mathbb{C} \mid \Re z > 0, |\Im z| \preceq (\Re z) \tan \alpha\} = \{re^{i\theta} \mid r > 0, |\theta| \leq \alpha\}.$$

Clearly, if A is positive definite, $W(A) \subset S_0$.

Mathematics subject classification (2010): 47A63, 15A45.

Keywords and phrases: Determinantal inequality, positive definite matrices, numerical range, sector, determinants of block matrices.

For fundamentals of numerical range, the readers can refer to [9, 10]. As $0 \notin S_\alpha$, if $W(A) \subset S_\alpha$, A is necessarily nonsingular.

Based on a recent result of Lin [3], Choi [1] proved the following determinantal inequality for positive definite matrices.

THEOREM 1.1. [1] *Let A_i , $i = 1, \dots, m$, be positive definite matrices whose diagonal blocks are n_j -square matrices $A_i^{(j)}$ for $j = 1, \dots, k$. Then*

$$\det\left(\sum_{i=1}^m A_i^{-1}\right) \geq \det\left(\sum_{i=1}^m (A_i^{(1)})^{-1}\right) \cdots \det\left(\sum_{i=1}^m (A_i^{(k)})^{-1}\right). \quad (1.2)$$

In [7], Haynsworth gave the result for the Schur complement of 2×2 block matrices.

THEOREM 1.2. [7] *Suppose $A, B \in \mathbb{M}_n(\mathbb{C})$ are Hermitian matrices, partitioned as in (1.1), $A = (A_{ij})$, $B = (B_{ij})$, $i, j = 1, 2$, where A_{11} and B_{11} are square of order m . If $A \succeq 0$, $B \succeq 0$, $A_{11} \succ 0$, $B_{11} \succ 0$, then*

$$\det((A+B)/(A_{11}+B_{11})) \geq \det A / \det A_{11} + \det B / \det B_{11}. \quad (1.3)$$

This result (1.3) has been extended in [4]. In this paper, we extend the results (1.2) and (1.3) to the case of matrices whose numerical ranges are contained in a sector.

2. Main results

The following lemmas are useful for proving the above theorems.

LEMMA 2.1. [2, p. 68] *Let $A \in \mathbb{M}_n(\mathbb{C})$. If $\Re A$ is positive definite, then*

$$\det(\Re A) \leq |\det(A)|. \quad (2.1)$$

LEMMA 2.2. [5, 6] *Let $A \in \mathbb{M}_n(\mathbb{C})$ with $W(A) \subset S_\alpha$. Then A can be decomposed as $A = XZX^*$ for some invertible matrix $X \in \mathbb{M}_n(\mathbb{C})$ and $Z = \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n})$ with $|\theta_j| \leq \alpha$ for all j .*

LEMMA 2.3. [4, Lemma 2.6] *Let $A \in \mathbb{M}_n(\mathbb{C})$ with $W(A) \subset S_\alpha$. Then*

$$\sec^n(\alpha) \det(\Re A) \geq |\det(A)|. \quad (2.2)$$

Proof. By Lemma 2.2, we have $A = XZX^*$ for some invertible matrix $X \in \mathbb{M}_n(\mathbb{C})$ and $Z = \text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n})$ with $|\theta_j| \leq \alpha$ for all j . Thus, $|\det(A)| = |\det(XZX^*)|$. After dividing by $|\det X|^2$, the inequality holds: $|\det A| / |\det X|^2 = |\det Z| = 1$. Furthermore, $\sec^n \alpha \det(\Re Z) \geq 1$. Therefore, $\sec^n \alpha \det(\Re Z) \geq |\det A| / |\det X|^2$. \square

LEMMA 2.4. [4, Lemma 2.4] Let $A \in \mathbb{M}_n(\mathbb{C})$ with $\Re A$ positive definite. Then

$$(\Re A)^{-1} \succeq \Re(A^{-1}). \quad (2.3)$$

Now we present some generalizations of Theorem 1.1 and Theorem 1.2. First of all, Theorem 1.1 is extended to the class of matrices whose numerical ranges are contained in a sector as follows:

THEOREM 2.5. Let A_i , $i = 1, \dots, m$, be a sequence of $n \times n$ matrices whose diagonal blocks are n_j -square matrices $A_i^{(j)}$ for $j = 1, \dots, k$. Assume that $W(A_i) \subset S_\alpha$, $i = 1, \dots, m$, $\alpha \in [0, \frac{\pi}{2})$. Then

$$|\det(\sum_{i=1}^m A_i^{-1})| \geq \cos^{3n}(\alpha) |\det(\sum_{i=1}^m (A_i^{(1)})^{-1})| \cdots |\det(\sum_{i=1}^m (A_i^{(k)})^{-1})|. \quad (2.4)$$

Proof. Compute

$$\begin{aligned} |\det(\sum_{i=1}^m A_i^{-1})| &\geq \det(\Re(\sum_{i=1}^m A_i^{-1})) = \det(\Re(A_1^{-1} + \cdots + A_m^{-1})) && \text{(by (2.1))} \\ &= \det \Re((X_1^{-1})^* Z_1^{-1} X_1^{-1} + \cdots + (X_m^{-1})^* Z_m^{-1} X_m^{-1}) && \text{(by Lemma 2.2)} \\ &= \det((X_1^{-1})^* \Re(Z_1^{-1}) X_1^{-1} + \cdots + (X_m^{-1})^* \Re(Z_m^{-1}) X_m^{-1}) \\ &\geq \det(\cos^2(\alpha) (X_1^{-1})^* (\Re Z_1)^{-1} X_1^{-1} + \cdots + \cos^2(\alpha) (X_m^{-1})^* (\Re Z_m)^{-1} X_m^{-1}) \\ &\quad \text{(by } \Re(Z^{-1}) \succeq \cos^2(\alpha) (\Re Z)^{-1} \text{)} \\ &= \cos^{2n}(\alpha) \det((X_1^{-1})^* (\Re Z_1)^{-1} X_1^{-1} + \cdots + (X_m^{-1})^* (\Re Z_m)^{-1} X_m^{-1}) \\ &= \cos^{2n}(\alpha) \det((X_1 \Re Z_1 X_1^*)^{-1} + \cdots + (X_m \Re Z_m X_m^*)^{-1}) \\ &\geq \cos^{2n}(\alpha) \det(\sum_{i=1}^m \Re A_i^{(1)})^{-1} \cdots \det(\sum_{i=1}^m \Re A_i^{(k)})^{-1} && \text{(by (1.2))} \\ &\geq \cos^{2n}(\alpha) \det(\sum_{i=1}^m \Re((A_i^{(1)})^{-1})) \cdots \det(\sum_{i=1}^m \Re((A_i^{(k)})^{-1})) && \text{(by (2.3))} \\ &\geq \cos^{3n}(\alpha) |\det(\sum_{i=1}^m (A_i^{(1)})^{-1})| \cdots |\det(\sum_{i=1}^m (A_i^{(k)})^{-1})|, && \text{(by (2.2))} \end{aligned}$$

where X_i ($i = 1, \dots, m$) and Z_i ($i = 1, \dots, m$) correspond to the invertible matrices and diagonal matrices in Lemma 2.2, respectively. \square

REMARK 2. When $\alpha = 0$, our Theorem 2.5 reduces to Theorem 1.1.

Next, Theorem 1.2 is extended to the class of matrices whose numerical ranges are contained in a sector as follows.

THEOREM 2.6. Let $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix} \in \mathbb{M}_n(\mathbb{C})$ with $A_{11}, B_{11} \in \mathbb{M}_m(\mathbb{C})$ and $W(A), W(B) \subset S_\alpha, \alpha \in [0, \frac{\pi}{2})$, then

$$\frac{|\det(A+B)|}{|\det(A_{11}+B_{11})|} \geq \cos^{3(n-m)} \alpha \left(\frac{|\det A|}{|\det A_{11}|} + \frac{|\det B|}{|\det B_{11}|} \right). \tag{2.5}$$

Proof. Compute

$$\begin{aligned} \frac{|\det(A+B)|}{|\det(A_{11}+B_{11})|} &\geq \frac{\det \Re(A+B)}{|\det(A_{11}+B_{11})|} && \text{(by (2.1))} \\ &\geq \cos^{2(n-m)} \alpha \det(\Re(A/A_{11}) + \Re(B/B_{11})) && \text{(by [4, Theorem 3.1])} \\ &\geq \cos^{2(n-m)} \alpha (\det(\Re(A/A_{11})) + \det(\Re(B/B_{11}))) \\ &\geq \cos^{3(n-m)} \alpha (|\det(A/A_{11})| + |\det(B/B_{11})|). && \text{(by (2.2)) } \square \end{aligned}$$

REMARK 3. When $\alpha = 0$, our Theorem 2.6 reduces to Haynsworth’s result (Theorem 1.2). In Theorem 2.6, if $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \in \mathbb{M}_{k+1}(\mathbb{C})$ with $A_{11} \in \mathbb{M}_k(\mathbb{C})$, then

$$\sec^3 \alpha \left| \frac{\det(A_{k+1} + B_{k+1})}{\det(A_k + B_k)} \right| \geq \left| \frac{\det A_{k+1}}{\det A_k} \right| + \left| \frac{\det B_{k+1}}{\det B_k} \right|,$$

which is the result in [4, (4.1)]. Thus, our result is a generalization of (4.1) in [4].

Acknowledgement. This research was supported by the key project of the applied mathematics of Hainan Normal University, the natural science foundation of Hainan Province (No. 20161005), Chongqing Graduate Student Research Innovation Project (No. CYS14020) and Doctoral scientific research foundation of Hainan Normal University.

REFERENCES

- [1] D. CHOI, *Determinantal inequalities of positive definite matrices*, Math. Inequal. Appl., Preprint.
- [2] S. G. WANG, M. X. WU, Z. Z. JIA, *Matrix Inequalities*, 2nd ed., Science Press, Beijing, 2004 (in Chinese).
- [3] M. LIN, *Determinantal inequalities for block triangular matrices*, Math. Inequal. Appl. **18** (2015) 1079–1086.
- [4] M. LIN, *Extension of a result of Haynsworth and Hartfiel*, Arch. Math. **104** (2015) 93–100.
- [5] S. W. DRURY, *Fischer determinantal inequalities and Higham’s Conjecture*, Linear Algebra Appl. **439** (2013) 3129–3133.
- [6] F. ZHANG, *A matrix decomposition and its applications*, Linear Multilinear Algebra **63** (2015), no. 10, 2038–2042.
- [7] E. V. HAYNSWORTH, *Applications of an inequality for the Schur complement*, Proc. Amer. Math. Soc. **24** (1970) 512–516.
- [8] R. BHATIA, *Matrix Analysis*, Grad. Texts in Math., vol. **169**, Springer-Verlag, New York, 1997.

- [9] K. E. GUSTAFSON, D. K. M. RAO, *Numerical Range: The Field of Values of Linear Operators and Matrices*, Springer, New York, 1997.
- [10] R. A. HORN, C. R. JOHNSON, *Topics in Matrix Analysis*, Cambridge University Press, 1997.

(Received April 21, 2016)

Xiaohui Fu
School of Mathematics and Statistics
Hainan Normal University
Haikou, 571158, P. R. China
and
Hainan Center for Mathematical Research
Hainan Normal University
Haikou, City Hainan Province
e-mail: fxh6662@sina.com

Yang Liu
College of Mathematics and Statistics
Chongqing University
Chongqing, 401331, P. R. China

Shunqin Liu
College of Information Science and Technology
Xiamen University Tan Kah Kee College
Xiamen, 363123, P. R. China