

## SOME NEW NONLINEAR WEAKLY SINGULAR INTEGRAL INEQUALITIES AND THEIR APPLICATIONS

RUN XU

(Communicated by Q.-H. Ma)

*Abstract.* The purpose of the present note is to establish some new weakly singular integral inequalities of Wendroff type, which generalized some known weakly singular inequalities for functions in two variables. The inequalities given here can be used in the analysis of various problems in the theory of certain classes of integral equations and evolution equations.

### 1. Introduction

It is well known that singular integral inequalities played very important roles in the qualitative theory of differential and integral equations. In the study of differential and integral equations, one often deal with certain integral inequalities. The Gronwall-Bellman type integral inequality and its various linear and nonlinear generalizations are crucial in the discussion of existence, uniqueness, continuation, boundedness, and stability properties of solutions. During the past few years, many papers [1–6, 8–16] have appeared in the literature which deal with integral and sum inequalities involving functions of one and more than one independent variables. Usually, the integrals concerning this type inequalities have regular or continuous kernels, but some problems of theory and practicality require us to solve integral inequalities with singular kernels. For example, Henry [7] used this type integral inequalities to prove a global existence and an exponential decay result for a parabolic Cauchy problem. Recently, Ma et al. [10] studied the inequality

$$u^p(t) \leq a(t) + b(t) \int_0^t (t^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} f(s) u^q(s) ds, \quad t \in \mathbb{R}_+.$$

In 2016, Xu and Meng [11] presented Gronwall-Bellman type integral inequalities with nonlinear weakly singular integral kernel of the form

$$u^p(t) \leq a(t) + b(t) \int_0^t (t-s)^{\beta-1} c(s) u^m(s) ds + d(t) \int_0^t (t^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} f(s) u^q(s) ds.$$

---

*Mathematics subject classification* (2010): 42B20, 26D07, 26D15.

*Keywords and phrases:* Integral inequality, weakly singular kernel, nonlinear.

This research is supported by National Science Foundation of China (11671227).

Cheung et al.[11] investigated the inequality in two variables

$$u^p(x,y) \leq a(x,y)+b(x,y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} f(s,t) u^q(s,t) dt ds, \\ (x,y) \in D.$$

In this paper, motivated mainly by the work of [9–12], and applying Medved’s method of desingularization of weakly singular inequalities, we establish some new singular version of the Wendroff inequality for functions in two variables. An example is included to illustrate the usefulness of our results.

### 2. Preliminary knowledge

In what follows,  $R$  denotes the set of real numbers,  $R_+ = [0, +\infty)$ ,  $C^i(M, S)$  denotes the class of all  $i$ -times continuously differentiable functions defined on set  $M$  with range in the set  $S$  ( $i = 1, 2, \dots$ ),  $C^0(M, S) = C(M, S)$ . For convenience, we cite some useful lemmas and definitions in the discussion of our proof as follows.

LEMMA 2.1. (See [8]) *Let  $a \geq 0$ ,  $p \geq q \geq 0$  and  $p \neq 0$ , then*

$$a^{\frac{q}{p}} \leq \frac{q}{p} K^{\frac{q-p}{p}} a + \frac{p-q}{p} K^{\frac{q}{p}},$$

for any  $K > 0$ .

DEFINITION 2.2. (See [9]) Let  $[x, y, z]$  be an ordered parameter group of non-negative real numbers. The group is called belong to the first class distribution and denoted by  $[x, y, z] \in I$  if conditions  $x \in (0, 1]$ ,  $y \in (\frac{1}{2}, 1)$  and  $z \geq \frac{3}{2} - y$  are satisfied. The group is called belong to the second class distribution and denoted by  $[x, y, z] \in II$ , if  $x \in (0, 1]$ ,  $y \in (0, \frac{1}{2}]$  and  $z > \frac{1-2y^2}{1-y^2}$  are satisfied.

LEMMA 2.3. (See [12, page 296]) *Let  $\alpha, \beta, \gamma$  and  $p$  be positive constants, then*

$$\int_0^t (t^\alpha - s^\beta)^{p(\beta-1)} s^{p(r-1)} ds = \frac{t^\theta}{\alpha} B \left[ \frac{p(\gamma-1)+1}{\alpha}, p(\beta-1)+1 \right], \quad t \in R_+,$$

where  $B[\xi, \eta] = \int_0^1 s^{\xi-1} (1-s)^{\eta-1} ds$  ( $\Re \xi > 0, \Re \eta > 0$ ) is the well-known  $B$ -function and  $\theta = p[\alpha(\beta-1) + \gamma - 1] + 1$ .

LEMMA 2.4. (See [9]) *Suppose that the positive constants  $\alpha, \beta, \gamma, p_1$  and  $p_2$  satisfy conditions:*

- (a) if  $[\alpha, \beta, \gamma] \in I$ ,  $p_1 = \frac{1}{\beta}$ ;
- (b) if  $[\alpha, \beta, \gamma] \in II$ ,  $p_2 = \frac{1+4\beta}{1+3\beta}$ ,

then  $B[\frac{p_i(\gamma-1)+1}{\alpha}, p_i(\beta-1)+1] \in (0, +\infty)$  and  $\theta_i = p_i[\alpha(\beta-1) + \gamma - 1] + 1 \geq 0$ ,  $i = 1, 2$ .

LEMMA 2.5. (See [6, page 329]) *Let  $u(x, y)$ ,  $p(x, y)$ ,  $q(x, y)$  and  $k(x, y)$  be non-negative continuous functions defined for  $x, y \in R_+$ . If*

$$u(x, y) \leq p(x, y) + q(x, y) \int_0^x \int_0^y k(s, t) u(s, t) dt ds, \tag{2.1}$$

then

$$u(x, y) \leq p(x, y) + q(x, y) \left( \int_0^x \int_0^y k(s, t) p(s, t) dt ds \right) \exp \left( \int_0^x \int_0^y k(s, t) q(s, t) dt ds \right). \tag{2.2}$$

### 3. Main result

THEOREM 3.1. *Let  $u(x, y)$ ,  $a(x, y)$ ,  $b(x, y)$ ,  $f(x, y)$ ,  $h(x, y)$  be nonnegative continuous functions for  $(x, y) \in D$ ,  $D = [0, X] \times [0, Y]$  ( $0 < X \leq \infty$ ,  $0 < Y \leq \infty$ ),  $a(x, y)$  and  $b(x, y)$  are nondecreasing in  $D$  and  $u(x, y)$  satisfies the following form of integral inequality*

$$u^p(x, y) \leq a(x, y) + b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \times \left[ f(s, t) u^q(s, t) + h(s, t) u^r(\sigma(s), \sigma(t)) \right] dt ds, \tag{3.1}$$

with the initial condition

$$u(x, y) = \phi(x, y), \quad (x, y) \in D' = [\mu, 0] \times [\mu, 0],$$

$$\phi(\sigma(x), \sigma(y)) \leq (a(x, y))^{\frac{1}{p}} \text{ for } (x, y) \in D \text{ with } \sigma(x) \leq 0, \sigma(y) \leq 0, \tag{3.2}$$

where  $p \neq 0$ ,  $p \geq q > 0$ ,  $p \geq r > 0$ ,  $p, q, r$  be constants,  $\sigma(x), \sigma(y) \in C(R_+, R)$ ,  $\sigma(t) \leq t, t \in R_+$ .  $-\infty < \mu = \inf\{\sigma(t), t \in R_+\} \leq 0$  and  $\phi(x, y) \in C([\mu, 0] \times [\mu, 0], R_+)$ . Then

(i) If  $[\alpha, \beta, \gamma] \in I$ ,

$$u(x, y) \leq \left\{ a(x, y) + \left[ P_1(x, y) + Q_1(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1}{1-\beta}} P_1(s, t) dt ds \right) \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1}{1-\beta}} Q_1(s, t) dt ds \right) \right]^{1-\beta} \right\}^{\frac{1}{p}}, \quad (x, y) \in D, \tag{3.3}$$

where

$$\begin{aligned}
 M_1 &= \frac{1}{\alpha} B \left[ \frac{\beta + \gamma - 1}{\alpha \beta}, \frac{2\beta - 1}{\beta} \right], \\
 M &= \max \left\{ \frac{q}{p} K^{\frac{q-p}{p}}, \frac{r}{p} K^{\frac{r-p}{p}} \right\}, \\
 A_1(x, y) &= \frac{q}{p} K^{\frac{q-p}{p}} a(x, y) + \frac{p-q}{p} K^{\frac{q}{p}}, \quad A_2(x, y) = \frac{r}{p} K^{\frac{r-p}{p}} a(x, y) + \frac{p-r}{p} K^{\frac{r}{p}}, \\
 \mathcal{A}_1(x, y) &= \int_0^x \int_0^y (f(s, t) A_1(s, t) + h(s, t) A_2(s, t))^{\frac{1}{1-\beta}} dt ds, \\
 P_1(x, y) &= 2^{\frac{\beta}{1-\beta}} M_1^{\frac{2\beta}{1-\beta}}(xy)^{\frac{(\alpha+1)(\beta-1)+\gamma}{1-\beta}} b^{\frac{1}{1-\beta}}(x, y) \mathcal{A}_1(x, y), \\
 Q_1(x, y) &= 2^{\frac{\beta}{1-\beta}} M_1^{\frac{2\beta}{1-\beta}}(xy)^{\frac{(\alpha+1)(\beta-1)+\gamma}{1-\beta}} b^{\frac{1}{1-\beta}}(x, y) M^{\frac{1}{1-\beta}}.
 \end{aligned}$$

(ii) If  $[\alpha, \beta, \gamma] \in \Pi$ ,

$$\begin{aligned}
 u(x, y) &\leq \left\{ a(x, y) + \left[ P_2(x, y) + Q_2(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1+4\beta}{\beta}} P_2(s, t) dt ds \right) \right. \right. \\
 &\quad \left. \left. \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1+4\beta}{\beta}} Q_2(s, t) dt ds \right) \right]^{\frac{\beta}{1+4\beta}} \right\}^{\frac{1}{\beta}}, \quad (x, y) \in D,
 \end{aligned} \tag{3.4}$$

where

$$\begin{aligned}
 M_2 &= \frac{1}{\alpha} B \left[ \frac{\gamma(1+4\beta) - \beta}{\alpha(1+3\beta)}, \frac{4\beta^2}{1+3\beta} \right], \\
 \mathcal{A}_2(x, y) &= \int_0^x \int_0^y (f(s, t) A_1(s, t) + h(s, t) A_2(s, t))^{\frac{1+4\beta}{\beta}} dt ds, \\
 P_2(x, y) &= 2^{\frac{1+3\beta}{\beta}} M_2^{\frac{2(1+3\beta)}{\beta}}(xy)^{\frac{(1+4\beta)[\alpha(\beta-1)+\gamma-\beta]}{\beta}} b^{\frac{1+4\beta}{\beta}}(x, y) \mathcal{A}_2(x, y), \\
 Q_2(x, y) &= 2^{\frac{1+3\beta}{\beta}} M_2^{\frac{2(1+3\beta)}{\beta}}(xy)^{\frac{(1+4\beta)[\alpha(\beta-1)+\gamma-\beta]}{\beta}} b^{\frac{1+4\beta}{\beta}}(x, y) M^{\frac{1+4\beta}{\beta}}.
 \end{aligned}$$

*Proof.* For any fixing positive number  $X, Y$ , we define a function  $\omega(x, y)$  by

$$\begin{aligned}
 \omega^p(x, y) &= a(X, Y) + b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\
 &\quad \times \left[ f(s, t) u^q(s, t) + h(s, t) u^r(\sigma(s), \sigma(y)) \right], \quad (x, y) \in D.
 \end{aligned} \tag{3.5}$$

Then we can easily see that  $\omega(x, y)$  is a nonnegative and nondecreasing function and

$$u(x, y) \leq \omega(x, y), \quad (x, y) \in D, \tag{3.6}$$

therefore, for  $x \in [0, X), y \in [0, Y)$  with  $\sigma(x) \geq 0, \sigma(y) \geq 0$ , we have

$$u(\sigma(x), \sigma(y)) \leq \omega(\sigma(x), \sigma(y)) \leq \omega(x, y), \quad (x, y) \in D, \tag{3.7}$$

on the other hand, for  $x \in [0, X)$ ,  $y \in [0, Y)$  with  $\sigma(x) \leq 0$ ,  $\sigma(y) \leq 0$ , using the initial condition (3.2) and the nondecreasing function  $a(x, y)$ , we have

$$u(\sigma(x), \sigma(y)) = \phi(\sigma(x), \sigma(y)) \leq (a(x, y))^{\frac{1}{p}} \leq (a(X, Y))^{\frac{1}{p}} \leq \omega(x, y), \quad (x, y) \in D, \tag{3.8}$$

and it follows from (3.7) and (3.8) that

$$u(\sigma(x), \sigma(y)) \leq \omega(x, y), \quad (x, y) \in D, \tag{3.9}$$

from (3.5) and (3.9), there is

$$\begin{aligned} \omega^p(x, y) &\leq a(X, Y) + b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s, t) \omega^q(s, t) + h(s, t) \omega^r(s, t) \right] dt ds. \end{aligned} \tag{3.10}$$

If taking  $x = X$ ,  $y = Y$  in (3.10), we have

$$\begin{aligned} \omega^p(X, Y) &\leq a(X, Y) + b(X, Y) \int_0^X \int_0^Y (X^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (Y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s, t) \omega^q(s, t) + h(s, t) \omega^r(s, t) \right] dt ds, \end{aligned} \tag{3.11}$$

for  $0 < X \leq \infty$ ,  $0 < Y \leq \infty$  be arbitrary constant, it proves that

$$\begin{aligned} \omega^p(x, y) &\leq a(x, y) + b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s, t) \omega^q(s, t) + h(s, t) \omega^r(s, t) \right] dt ds, \quad (x, y) \in D, \end{aligned} \tag{3.12}$$

set

$$\begin{aligned} v(x, y) &= b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s, t) \omega^q(s, t) + h(s, t) \omega^r(s, t) \right] dt ds, \end{aligned} \tag{3.13}$$

then

$$\omega^p(x, y) \leq a(x, y) + v(x, y), \tag{3.14}$$

or

$$\omega(x, y) \leq (a(x, y) + v(x, y))^{\frac{1}{p}}, \tag{3.15}$$

by Lemma 2.1 and inequality (3.15), for any  $K > 0$ , we have

$$\omega^q(s, t) \leq (a(s, t) + v(s, t))^{\frac{q}{p}} \leq \frac{q}{p} K^{\frac{q-p}{p}} (a(s, t) + v(s, t)) + \frac{p-q}{p} K^{\frac{q}{p}}, \tag{3.16}$$

$$\omega^r(s, t) \leq (a(s, t) + v(s, t))^{\frac{r}{p}} \leq \frac{r}{p} K^{\frac{r-p}{p}} (a(s, t) + v(s, t)) + \frac{p-r}{p} K^{\frac{r}{p}}, \tag{3.17}$$

it follows from (3.13), (3.16) and (3.17) that

$$\begin{aligned} v(x, y) &\leq b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s, t) \left( \frac{q}{p} K^{\frac{q-p}{p}} (a(s, t) + v(s, t)) + \frac{p-q}{p} K^{\frac{q}{p}} \right) \right. \\ &\quad \left. + h(s, t) \left( \frac{r}{p} K^{\frac{r-p}{p}} (a(s, t) + v(s, t)) + \frac{p-r}{p} K^{\frac{r}{p}} \right) \right] dt ds \\ &= b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} (f(s, t) A_1(s, t) + h(s, t) A_2(s, t)) \\ &\quad + Mb(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} (f(s, t) + h(s, t)) v(s, t) dt ds, \end{aligned} \tag{3.18}$$

where

$$\begin{aligned} A_1(x, y) &= \frac{q}{p} K^{\frac{q-p}{p}} a(x, y) + \frac{p-q}{p} K^{\frac{q}{p}}, \quad A_2(x, y) = \frac{r}{p} K^{\frac{r-p}{p}} a(x, y) + \frac{p-r}{p} K^{\frac{r}{p}}, \\ M &= \max \left\{ \frac{q}{p} K^{\frac{q-p}{p}}, \frac{r}{p} K^{\frac{r-p}{p}} \right\}. \end{aligned}$$

If  $[\alpha, \beta, \gamma] \in I$ , let  $p_1 = \frac{1}{\beta}$ ,  $q_1 = \frac{1}{1-\beta}$ ; if  $[\alpha, \beta, \gamma] \in II$ , let  $p_2 = \frac{1+4\beta}{1+3\beta}$ ,  $q_2 = \frac{1+4\beta}{\beta}$ , then  $\frac{1}{p_i} + \frac{1}{q_i} = 1$  for  $i = 1, 2$ . By applying Hölder's inequality with indices  $p_i, q_i$  to (3.18), we get

$$\begin{aligned} v(x, y) &\leq b(x, y) \left[ \int_0^x \int_0^y (x^\alpha - s^\alpha)^{p_i(\beta-1)} s^{p_i(\gamma-1)} (y^\alpha - t^\alpha)^{p_i(\beta-1)} t^{p_i(\gamma-1)} dt ds \right]^{\frac{1}{p_i}} \\ &\quad \times \left[ \int_0^x \int_0^y (f(s, t) A_1(s, t) + h(s, t) A_2(s, t))^{q_i} dt ds \right]^{\frac{1}{q_i}} \\ &\quad + Mb(x, y) \left[ \int_0^x \int_0^y (x^\alpha - s^\alpha)^{p_i(\beta-1)} s^{p_i(\gamma-1)} (y^\alpha - t^\alpha)^{p_i(\beta-1)} t^{p_i(\gamma-1)} dt ds \right]^{\frac{1}{p_i}} \\ &\quad \times \left[ \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} v^{q_i}(s, t) dt ds \right]^{\frac{1}{q_i}}, \end{aligned} \tag{3.19}$$

by Lemma 2.3 and Lemma 2.4, the last inequality can be rewritten as

$$\begin{aligned} v(x, y) &\leq b(x, y) (M_i^2(xy)^{\theta_i})^{\frac{1}{p_i}} \mathcal{A}_i^{\frac{1}{q_i}}(x, y) + Mb(x, y) (M_i^2(xy)^{\theta_i})^{\frac{1}{p_i}} \\ &\quad \times \left[ \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} v^{q_i}(s, t) dt ds \right]^{\frac{1}{q_i}}, \quad (x, y) \in D, \end{aligned} \tag{3.20}$$

where

$$M_i = \frac{1}{\alpha} B \left[ \frac{p_i(\gamma - 1) + 1}{\alpha}, p_i(\beta - 1) + 1 \right],$$

$$\mathcal{A}_i(x, y) = \int_0^x \int_0^y (f(s, t)A_1(s, t) + h(s, t)A_2(s, t))^{q_i} dt ds,$$

and  $\theta_i$  is given as in Lemma 2.4 for  $i = 1, 2$ .

We also need the following well-know consequence of Jensen inequality

$$(A_1 + A_2 + \dots + A_n)^r \leq n^{r-1} (A_1^r + A_2^r + \dots + A_n^r) \quad (JI)$$

for  $A_i \geq 0$  ( $i = 1, 2, \dots, n$ ) and  $r \geq 1$ . Applying inequality (JI) to (3.20), we get

$$v^{q_i}(x, y) \leq 2^{q_i-1} (M_i^2(xy)^{\theta_i})^{\frac{q_i}{p_i}} b^{q_i}(x, y) \mathcal{A}_i(x, y) + 2^{q_i-1} M^{q_i} b^{q_i}(x, y) (M_i^2(xy)^{\theta_i})^{\frac{q_i}{p_i}}$$

$$\times \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} v^{q_i}(s, t) dt ds, \tag{3.21}$$

by Lemma 2.5 and the last inequality, we have

$$v^{q_i}(x, y) \leq P_i(x, y) + Q_i(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} P_i(s, t) dt ds \right)$$

$$\times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} Q_i(s, t) dt ds \right), \tag{3.22}$$

where

$$P_i(x, y) = 2^{q_i-1} (M_i^2(xy)^{\theta_i})^{\frac{q_i}{p_i}} b^{q_i}(x, y) \mathcal{A}_i(x, y),$$

$$Q_i(x, y) = 2^{q_i-1} (M_i^2(xy)^{\theta_i})^{\frac{q_i}{p_i}} b^{q_i}(x, y) M^{q_i},$$

finally, substituting (3.22) into (3.15) leads to

$$\omega(x, y) \leq \left\{ a(x, y) + \left[ P_i(x, y) + Q_i(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} P_i(s, t) dt ds \right) \right. \right.$$

$$\left. \left. \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} Q_i(s, t) dt ds \right) \right]^{\frac{1}{q_i}} \right\}^{\frac{1}{p}}, \tag{3.23}$$

from (3.6), we have

$$u(x, y) \leq \left\{ a(x, y) + \left[ P_i(x, y) + Q_i(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} P_i(s, t) dt ds \right) \right. \right.$$

$$\left. \left. \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{q_i} Q_i(s, t) dt ds \right) \right]^{\frac{1}{q_i}} \right\}^{\frac{1}{p}}. \tag{3.24}$$

Considering two situations for  $i = 1, 2$  and using parameters  $\alpha, \beta$  and  $\gamma$  to denote  $p_i, q_i$  and  $\theta_i$  in (3.24), we can get the desired estimations.  $\square$

COROLLARY 3.2. Let functions  $u(x, y)$ ,  $a(x, y)$ ,  $b(x, y)$ ,  $f(x, y)$  and  $h(s, t)$  be defined as in Theorem 3.1, and  $q, r$  be a constant with  $0 < q \leq 1$ ,  $0 < r \leq 1$ , suppose that

$$u(x, y) \leq a(x, y) + b(x, y) \int_0^x \int_0^y (x-s)^{\beta-1} s^{\gamma-1} (y-t)^{\beta-1} t^{\gamma-1} \times \left[ f(s, t)u^q(s, t) + h(s, t)u^r(\sigma(s), \sigma(t)) \right] dt ds, \quad (x, y) \in D, \quad (3.25)$$

with the initial condition of Theorem 3.1, then one has the following

(i) If  $\beta \in (\frac{1}{2}, 1)$ ,

$$u(x, y) \leq a(x, y) + \left[ \bar{P}_1(x, y) + \bar{Q}_1(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1}{1-\beta}} \bar{P}_1(s, t) dt ds \right) \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1}{1-\beta}} \bar{Q}_1(s, t) dt ds \right) \right]^{1-\beta}, \quad (x, y) \in D, \quad (3.26)$$

where

$$\begin{aligned} \bar{M}_1 &= B \left[ \frac{\beta + \gamma - 1}{\beta}, \frac{2\beta - 1}{\beta} \right], \\ \bar{M} &= \max \left\{ qK^{q-1}, rK^{r-1} \right\}, \\ \bar{A}_1(x, y) &= qK^{q-1}a(x, y) + (1 - q)K^q, \quad \bar{A}_2(x, y) = rK^{r-1}a(x, y) + (1 - r)K^r, \\ \bar{\mathcal{A}}_1(x, y) &= \int_0^x \int_0^y (f(s, t)\bar{A}_1(s, t) + h(s, t)\bar{A}_2(s, t))^{\frac{1}{1-\beta}} dt ds, \\ \bar{P}_1(x, y) &= 2^{\frac{\beta}{1-\beta}} \bar{M}_1^{\frac{2\beta}{1-\beta}} (xy)^{\frac{2\beta+\gamma-2}{1-\beta}} b^{\frac{1}{1-\beta}} (x, y) \bar{\mathcal{A}}_1(x, y), \\ \bar{Q}_1(x, y) &= 2^{\frac{\beta}{1-\beta}} \bar{M}_1^{\frac{2\beta}{1-\beta}} (xy)^{\frac{2\beta+\gamma-2}{1-\beta}} b^{\frac{1}{1-\beta}} (x, y) \bar{M}^{\frac{1}{1-\beta}}. \end{aligned}$$

(ii) If  $\beta \in (0, \frac{1}{2}]$ ,

$$u(x, y) \leq a(x, y) + \left[ \bar{P}_2(x, y) + \bar{Q}_2(x, y) \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1+4\beta}{\beta}} \bar{P}_2(s, t) dt ds \right) \times \exp \left( \int_0^x \int_0^y (f(s, t) + h(s, t))^{\frac{1+4\beta}{\beta}} \bar{Q}_2(s, t) dt ds \right) \right]^{\frac{\beta}{1+4\beta}}, \quad (x, y) \in D, \quad (3.27)$$



where

$$\begin{aligned} \overline{M}_2 &= B \left[ \frac{\gamma(1+4\beta) - \beta}{1+3\beta}, \frac{4\beta^2}{1+3\beta} \right], \\ \overline{\mathcal{A}}_2(x,y) &= \int_0^x \int_0^y (f(s,t)\overline{A}_1(s,t) + h(s,t)\overline{A}_2(s,t))^{\frac{1+4\beta}{\beta}} dt ds, \\ \overline{P}_2(x,y) &= 2^{\frac{1+3\beta}{\beta}} \overline{M}_2^{\frac{2(1+3\beta)}{\beta}} (xy)^{\frac{(4\beta+1)(\gamma-1)+4\beta^2}{\beta}} b^{\frac{1+4\beta}{\beta}}(x,y) \overline{\mathcal{A}}_2(x,y), \\ \overline{Q}_2(x,y) &= 2^{\frac{1+3\beta}{\beta}} \overline{M}_2^{\frac{2(1+3\beta)}{\beta}} (xy)^{\frac{(4\beta+1)(\gamma-1)+4\beta^2}{\beta}} b^{\frac{1+4\beta}{\beta}}(x,y) \overline{M}^{\frac{1+4\beta}{\beta}}. \end{aligned}$$

*Proof.* Inequalities (3.26) and (3.27) follow by letting  $p = \alpha = 1$  and  $0 < q \leq 1, 0 < r \leq 1$  in Theorem 3.1 and by simple computation. Details are omitted here.  $\square$

**COROLLARY 3.3.** *Let functions  $u(x,y), a(x,y), b(x,y), f(s,t)$  and  $h(x,y)$  be defined as in Theorem 3.1, suppose that*

$$\begin{aligned} u^2(x,y) &\leq a(x,y) + b(x,y) \int_0^x \int_0^y (x-s)^{\beta-1} s^{\gamma-1} (y-t)^{\beta-1} t^{\gamma-1} \\ &\quad \times \left[ f(s,t)u(s,t) + h(s,t)u(\sigma(s), \sigma(t)) \right] dt ds, \quad (x,y) \in D, \end{aligned} \tag{3.28}$$

with the initial condition of Theorem 3.1, then for any  $K > 0$ , one has the following

(i) If  $\beta \in (\frac{1}{2}, 1)$ ,

$$\begin{aligned} u^2(x,y) &\leq a(x,y) + \left[ \tilde{P}_1(x,y) + \tilde{Q}_1(x,y) \left( \int_0^x \int_0^y (f(s,t) + h(s,t))^{\frac{1}{1-\beta}} \tilde{P}_1(s,t) dt ds \right) \right. \\ &\quad \left. \times \exp \left( \int_0^x \int_0^y (f(s,t) + h(s,t))^{\frac{1}{1-\beta}} \tilde{Q}_1(s,t) dt ds \right) \right]^{1-\beta}, \quad (x,y) \in D, \end{aligned} \tag{3.29}$$

where

$$\begin{aligned} \tilde{M}_1 &= B \left[ \frac{\beta + \gamma - 1}{\beta}, \frac{2\beta - 1}{\beta} \right], \\ \tilde{A}_1(x,y) &= \tilde{A}_2(x,y) = \frac{1}{2} K^{-\frac{1}{2}} a(x,y) + \frac{1}{2} K^{\frac{1}{2}}, \\ \tilde{\mathcal{A}}_1(x,y) &= \int_0^x \int_0^y \left[ (f(s,t) + h(s,t))\tilde{A}_1(s,t) \right]^{\frac{1}{1-\beta}} dt ds, \\ \tilde{P}_1(x,y) &= 2^{\frac{1}{1-\beta}} \tilde{M}_1^{\frac{2\beta}{1-\beta}} (xy)^{\frac{2\beta+\gamma-2}{1-\beta}} b^{\frac{1}{1-\beta}}(x,y) \tilde{\mathcal{A}}_1(x,y), \\ \tilde{Q}_1(x,y) &= 2^{\frac{\beta}{1-\beta}} \tilde{M}_1^{\frac{2\beta}{1-\beta}} (xy)^{\frac{2\beta+\gamma-2}{1-\beta}} b^{\frac{1}{1-\beta}}(x,y) \tilde{M}^{\frac{1}{1-\beta}}. \end{aligned}$$

(ii) If  $\beta \in (0, \frac{1}{2}]$ ,

$$\begin{aligned}
 u^2(x,y) \leq & a(x,y) + \left[ \tilde{P}_2(x,y) + \tilde{Q}_2(x,y) \left( \int_0^x \int_0^y (f(s,t) + h(s,t)) \frac{1+4\beta}{\beta} \tilde{P}_2(s,t) dt ds \right) \right. \\
 & \left. \times \exp \left( \int_0^x \int_0^y (f(s,t) + h(s,t)) \frac{1+4\beta}{\beta} \tilde{Q}_2(s,t) dt ds \right) \right]^{\frac{\beta}{1+4\beta}}, \quad (x,y) \in D,
 \end{aligned}
 \tag{3.30}$$

where

$$\begin{aligned}
 \tilde{M}_2 &= B \left[ \frac{\gamma(1+4\beta) - \beta}{1+3\beta}, \frac{4\beta^2}{1+3\beta} \right], \\
 \tilde{\mathcal{A}}_2(x,y) &= \int_0^x \int_0^y \left[ (f(s,t) + h(s,t)) \tilde{A}_2(s,t) \right]^{\frac{1+4\beta}{\beta}} dt ds, \\
 \tilde{P}_2(x,y) &= 2^{\frac{1+4\beta}{\beta}} \tilde{M}_2^{\frac{2(1+3\beta)}{\beta}} (xy)^{\frac{(4\beta+1)(\gamma-1)+4\beta^2}{\beta}} b^{\frac{1+4\beta}{\beta}} (x,y) \tilde{\mathcal{A}}_2(x,y), \\
 \tilde{Q}_1(x,y) &= 2^{\frac{1+3\beta}{\beta}} \tilde{M}_2^{\frac{2(1+3\beta)}{\beta}} (xy)^{\frac{(4\beta+1)(\gamma-1)+4\beta^2}{\beta}} b^{\frac{1+4\beta}{\beta}} (x,y) \tilde{M}^{\frac{1+4\beta}{\beta}}.
 \end{aligned}$$

*Proof.* Inequality (3.29) and (3.30) follow by letting  $p = 2$ ,  $q = r = \alpha = 1$  in Theorem 3.1 and by simple computation. Details are omitted.  $\square$

### 4. Applications

In this section, we will indicate the usefulness of our main result in the study of the boundedness of solutions for certain partial integral equations with weakly singular kernel.

Consider the partial integral equation:

$$\begin{aligned}
 z^p(x,y) = & l(x,y) + g(x,y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\
 & \times \left[ F_1(s,t, z(s,t)) + F_2(s,t) z^r(\sigma(s), \sigma(t)) \right] dt ds,
 \end{aligned}
 \tag{4.1}$$

for  $(x,y) \in D = [0,X) \times [0,Y)$ , where  $l(x,y)$ ,  $g(x,y)$ ,  $z(x,y)$ ,  $F_2(s,t) \in C(D, \mathbb{R})$ ,  $F_1(s,t, z(s,t)) \in (D \times \mathbb{R}, \mathbb{R})$ . Suppose that

$$\begin{aligned}
 |l(x,y)| &\leq a(x,y), \\
 |g(x,y)| &\leq b(x,y), \\
 |F_1(s,t, z)| &\leq f(s,t) |z^q|, \\
 |F_2(s,t)| &\leq h(s,t),
 \end{aligned}
 \tag{4.2}$$

if  $z(x, y)$ ,  $(x, y) \in D$  is any solution of (4.1), we have

$$|z^p(x, y)| \leq a(x, y) + b(x, y) \int_0^x \int_0^y (x^\alpha - s^\alpha)^{\beta-1} s^{\gamma-1} (y^\alpha - t^\alpha)^{\beta-1} t^{\gamma-1} \\ \times \left[ f(s, t) |z^q(s, t)| + h(s, t) |z^r(\sigma(s), \sigma(t))| \right] dt ds. \quad (4.3)$$

with the initial condition

$$|z(x, y)| = \phi(x, y), \quad (x, y) \in D' = [\mu, 0] \times [\mu, 0], \\ \phi(\sigma(x), \sigma(y)) \leq (a(x, y))^{\frac{1}{p}} \text{ for } (x, y) \in D \text{ with } \sigma(x) \leq 0, \sigma(y) \leq 0, \quad (4.4)$$

where  $p \neq 0$ ,  $p \geq q > 0$ ,  $p \geq r > 0$ ,  $p, q, r$  be constants, the functions  $a(x, y)$ ,  $b(x, y)$ ,  $f(x, y)$ ,  $h(s, t)$  be nonnegative continuous functions for  $(x, y) \in D$  and  $\sigma(x)$ ,  $\sigma(y) \in C(R_+, R)$ ,  $\sigma(t) \leq t$ ,  $t \in R_+$ .  $-\infty < \mu = \inf\{\sigma(t), t \in R_+\} \leq 0$  and  $\phi(x, y) \in C([\mu, 0] \times [\mu, 0], R_+)$ .

It is easy for us to find that the inequality (4.3) is similar to (3.1), and applying Theorem 3.1, we have the same results as (3.3) and (3.4).

#### REFERENCES

- [1] E. F. BECKENBACH AND R. BELLMAN, *Inequalities*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, Berlin, Germany, 1961.
- [2] V. LAKSHMIKANTHAM AND S. LEELA, *Differential and Integral Inequalities: Theory and Applications*, vol. I: Ordinary Differential Equations, Mathematics in Science and Engineering, Academic Press, New York, NY, USA, 1969.
- [3] V. LAKSHMIKANTHAM AND S. LEELA, *Differential and Integral Inequalities: Theory and Applications*, vol. II: Functional, Partial, Abstract, and Complex Differential Equations, Mathematics in Science and Engineering, Academic Press, New York, NY, USA, 1969.
- [4] D. BANOV AND P. SIMEONOV, *Integral Inequalities and Applications*, vol. 57 of Mathematics and Its Applications, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1992.
- [5] R. P. AGARWAL, *Difference Equations and Inequalities*, Marcel Dekker, New York, NY, USA, 1993.
- [6] B. G. PACHPATTE, *Inequalities for Differential and Integral Equations*, vol. 197 of Mathematics in Science and Engineering, Academic Press, San Diego, Calif, USA, 1998.
- [7] D. HENRY, *Geometric Theory of Semilinear Parabolic Equations*, vol. 840 of Lecture Notes in Mathematics, Springer, Berlin, Germany, 1981.
- [8] F. C. JIANG, F. W. MENG, *Explicit bounds on some new nonlinear integral inequalities with delay*, J. Comput. Appl. Math. **205** (2007) 479–486.
- [9] Q. H. MA AND E. H. YANG, *Estimates on solutions of some weakly singular Volterra integral inequalities*, Acta Mathematicae Applicatae Sinica, vol. **25**, no. 3, pp. 505–515, 2002.
- [10] Q. H. MA AND J. PEČARIĆ, *Some new explicit bounds for weakly singular integral inequalities with applications to fractional differential and integral equations*, Journal of Mathematical Analysis and Applications, vol. 341, no. 2, pp. 894–905, 2008.
- [11] RUN XU, FANWEI MENG, *Some new weakly singular integral inequalities and their applications to fractional differential equations*, Journal of Inequalities and Applications **2016** (1), 1–16.
- [12] W. S. CHEUNG, *On some new integral differential inequalities of the Gronwall and Wendroff type*, Journal of Mathematical Analysis and Applications, vol. 178, no. 2, pp. 438–449, 1993.
- [13] A. P. PRUDNIKOV, YU. A. BRYCHKOV, AND O. I. MARICHEV, *Integrals and Series*, Elementary Functions, vol. 1, “Nauka”, Moscow, Russia, 1981.
- [14] RUN XU, FANWEI MENG, CUIHUA SONG, *On Some Integral Inequalities on Time Scales and Their Applications*, Journal of Inequalities and Applications, Volume 2010, Article ID 464976, 1–13.

- [15] RUN XU, YING ZHANG, *Generalized Gronwall fractional summation inequalities and their applications*, Journal of Inequalities and Applications, (2015) 2015:242.
- [16] ZHAOWEN ZHENG, XIN GAO AND JING SHAO, *Some new generalized retarded inequalities for discontinuous functions and their applications*, Journal of Inequalities and Applications, **2016** (1), 1–14, doi 10.1186/s13660-015-0943-6.

(Received February 23, 2016)

Run Xu  
Department of Mathematics  
Qufu Normal University  
Qufu, 273165, Shandong, People's Republic of China  
e-mail: xurun2005@163.com