

ON TWO PROBLEMS FOR GAUSS COMPOUND MEAN

LADISLAV MATEJÍČKA

Dedicated to my father

(Communicated by T. Burić)

Abstract. In this paper, we give answers to two open problems concerning Gauss Compound mean posed in the paper: [ZH. H. YANG, *A New Proof of Inequalities for Gauss Compound Mean*, Int. Journal of Math. Analysis, Vol. 4, (2010), no. 21, 1013–1018.] One of the problems is equivalent to the Vamanamurthy problem.

1. Introduction

Within the past years, means have been the subject of intensive research. Many interesting results and inequalities for means can be found in the literature [1]–[9]. The well known Gauss compound mean of positive numbers a, b with $a \neq b$ is the limit

$$AGM(a, b) = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$$

where

$$a_1 = a, \quad b_1 = b, \quad a_{n+1} = \frac{a_n + b_n}{2} = A(a_n, b_n), \quad b_{n+1} = \sqrt{a_n b_n} = G(a_n, b_n). \quad (1)$$

The power mean of positive a and b is defined by

$$M_t(a, b) = M^{1/t}(a^t, b^t) \quad \text{if } t \neq 0 \quad \text{and} \quad M_0(a, b) = \sqrt{ab}. \quad (2)$$

The t -order logarithmic mean of $a, b > 0$ is defined by

$$L_t(a, b) = \begin{cases} \left(\frac{b^t - a^t}{t(\ln b - \ln a)} \right)^{\frac{1}{t}}, & t \neq 0 \\ \sqrt{ab}, & t = 0, \end{cases} \quad (3)$$

where $L_1(a, b) = L(a, b) = \frac{b-a}{\ln b - \ln a}$ is the logarithmic mean.

Mathematics subject classification (2010): 26D07, 26E60.

Keywords and phrases: Gauss compound mean, logarithmic mean, inequality, open problems.

We note, that AGM is connected with elliptic integrals by Gauss’s fantastic result:

$$AGM(1, r') = \frac{\pi}{2k(r)}$$

where $k(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 x)^{-1/2} dx$, $r' = \sqrt{1 - r^2}$, $0 \leq r \leq 1$ (see [2]).

In the paper [8], the author presented new sophisticated proofs for famous inequalities

$$L(a, b) < AGM(a, b) < L^{2/3}(a^{3/2}, b^{3/2})$$

where the first inequality was proved by B. C. Carlson and M. Vuorinen [4], while the second one is due to J. and P. Borwein [2].

Denote by

$$D_n(t) = b_{n+1}^{-t} \left(\frac{b'_n - a'_n}{\ln b_n - \ln a_n} - \frac{b'_{n+1} - a'_{n+1}}{\ln b_{n+1} - \ln a_{n+1}} \right), \tag{4}$$

where a_n, b_n is defined by (1). It was showed in [8] that $t \mapsto D_n(t)$ is continuous on \mathbb{R} and increasing on $(1, \infty)$ with $D_n(1) < 0$, $D_n(3/2) > 0$. So $D_n(t)$ has an unique zero $t_0^{(n)}(a, b)$ on $(1, 3/2)$.

Denote by $t_{0i}(a, b) = \inf_{n \in \mathbb{N}} \{t_0^{(n)}(a, b)\}$ and $t_{0s}(a, b) = \sup_{n \in \mathbb{N}} \{t_0^{(n)}(a, b)\}$. It is evident that $1 \leq t_{0i}(a, b) < 3/2$, $1 < t_{0s}(a, b) \leq 3/2$.

At the end in the same paper [8], the author posed two open problems and remarks as follows:

PROBLEM 1. Is $t_{0i}(a, b)$ equal to 1?

REMARK 1. If the answer is “no”, then there must exist $t_1 \in (1, t_{0i}(a, b))$ such that $D_n(t_1) < 0$ for all n , whence

$$L(a_n^{t_1}, b_n^{t_1}) < L(a_{n+1}^{t_1}, b_{n+1}^{t_1}) \text{ for all } n,$$

which implies

$$L_{t_1}(a, b) < AGM(a, b), \tag{5}$$

where $L_t(a, b)$ is defined by (3).

Furthermore, are there some concrete constants $t_1 \in (1, t_{0i}(a, b))$ such that (5) holds?

PROBLEM 2. Is $t_{0s}(a, b)$ equal to 3/2?

REMARK 2. If the answer is “no”, then there must exist $t_2 \in (t_{0s}(a, b), 3/2)$ such that $D_n(t_2) > 0$ for all n , whence

$$L(a_n^{t_2}, b_n^{t_2}) > L(a_{n+1}^{t_2}, b_{n+1}^{t_2}) \text{ for all } n,$$

which implies

$$AGM(a, b) < L_{t_2}(a, b), \tag{6}$$

where $L_t(a, b)$ is defined by (3).

Furthermore, are there some concrete constants $t_2 \in (t_{0s}(a, b), 3/2)$ such that (6) holds?

REMARK 3. If replacing a^t, b^t by a, b in $L_t(a, b)$ and a, b by $a^{1/t}, b^{1/t}$ in $AGM(a, b)$, then the inequality (5) is equivalent to

$$L(a, b) < AGM_{1/t_1}(a, b) \text{ for } 1/t_1 \in (1/t_{0i}(a, b), 1). \tag{7}$$

This shows that the open problem presented by Vamanamurthy is equivalent to the problem 1.

In this note, we will prove that

- for $\forall a, b > 0$ with $a \neq b$ we have $t_{0i}(a, b) > 1$, and $\inf_{a, b > 0, a \neq b} \{t_{0i}(a, b)\} = 1$;
- for $\forall a, b > 0$ with $a \neq b$ we have $t_{0s}(a, b) = 3/2$.

These solve two open problems as above.

2. Two lemmas

In order to solve two open problems posed in [8], we need two important lemmas. Without loss of generality, we assume that $b < a$. Denote by $s_n = s_n(a, b) = b_n/a_n$, then $0 < s_n = s_n(a, b) < 1$. We have

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{1 + b_n/a_n}{2} = \frac{1 + s_n}{2}, \\ \frac{b_{n+1}}{a_{n+1}} &= \frac{\sqrt{a_n b_n}}{(a_n + b_n)/2} = 2 \frac{\sqrt{b_n/a_n}}{1 + b_n/a_n} = 2 \frac{\sqrt{s_n}}{1 + s_n}, \\ \frac{a_n}{b_{n+1}} &= \frac{a_n}{\sqrt{a_n b_n}} = \frac{1}{\sqrt{b_n/a_n}} = \frac{1}{\sqrt{s_n}}. \end{aligned}$$

Then $D_n(t)$ and $D'_n(t)$ can be expressed as

$$\begin{aligned} D_n(t) &= b_{n+1}^{-t} \left(\frac{b_n^t - a_n^t}{\ln b_n - \ln a_n} - \frac{b_{n+1}^t - a_{n+1}^t}{\ln b_{n+1} - \ln a_{n+1}} \right) \\ &= \left(\frac{a_n}{b_{n+1}} \right)^t \left(\frac{(b_n/a_n)^t - 1}{\ln(b_n/a_n)} \right) - \frac{1 - (a_{n+1}/b_{n+1})^t}{\ln(b_{n+1}/a_{n+1})} \\ &= \frac{s_n^{t/2} - s_n^{-t/2}}{\ln s_n} - \frac{1 - \left(\frac{1+s_n}{2\sqrt{s_n}}\right)^t}{\ln \frac{2\sqrt{s_n}}{1+s_n}}, \end{aligned} \tag{8}$$

$$D'_n(t) = \frac{s_n^{-t/2} + s_n^{t/2}}{2} - \left(\frac{s_n^{-1/2} + s_n^{1/2}}{2} \right)^t.$$

We now show that $t \mapsto D_n(t)$, $n \in \mathbb{N}$ is a convex function on $[1, \infty)$.

LEMMA 1. *Let $D_n(t)$ $n \in \mathbb{N}$ be defined in (2). Then $D''_n(t) > 0$ for $n \in \mathbb{N}$, $t \in [1, \infty)$.*

Proof. Differentiation gives

$$\begin{aligned} s_n^{t/2} D''_n(t) &= \frac{s_n^t - 1}{4} \ln s_n - \left(\frac{1 + s_n}{2} \right)^t \ln \left(\frac{1 + s_n}{2\sqrt{s_n}} \right) := g(t), \\ g'(t) &= \left(\frac{1 + s_n}{2} \right)^t \ln \left(\frac{2\sqrt{s_n}}{1 + s_n} \right) \ln \left(\frac{1 + s_n}{2} \right) + \frac{s_n^t \ln^2 s_n}{4} \geq 0, \end{aligned}$$

where the inequality holds due to $0 < s_n < 1$.

Clearly, the proof will be done if we prove that

$$g(1) = h(s_n) = \left(\frac{1 + s_n}{2} \right) \ln \left(\frac{2\sqrt{s_n}}{1 + s_n} \right) + \frac{s_n - 1}{4} \ln s_n \geq 0.$$

Differentiation again yields that for $0 < s_n < 1$,

$$h'(s_n) = \frac{1}{2} \ln \left(\frac{2s_n}{1 + s_n} \right) \leq 0,$$

which indicates that $h(s_n) > h(1) = 0$ and the proof of our lemma is complete. \square

LEMMA 2. *For $n \in \mathbb{N}$ and $a, b > 0$ with $a \neq b$, let*

$$r_n = 1 - \frac{1}{2} \frac{D_n(1)}{D_n(3/2) - D_n(1)},$$

where $D_n(t)$ is given by (2). Then we have

$$\lim_{n \rightarrow \infty} r_n = \frac{3}{2}.$$

Proof. To complete the proof, we write r_n as

$$r_n = 1 + \frac{1}{2} \frac{1}{D_n(3/2)/(-D_n(1)) + 1}.$$

By (8) we have

$$\begin{aligned} \frac{D_n(3/2)}{-D_n(1)} &= \left[\frac{s_n^{-3/4} - s_n^{3/4}}{\ln s_n} - \frac{\left(\frac{1+s_n}{2\sqrt{s_n}}\right)^{3/2} - 1}{\ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right)} \right] \bigg/ \left[\frac{\frac{1+s_n}{2\sqrt{s_n}} - 1}{\ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right)} - \frac{\frac{1}{\sqrt{s_n}} - \sqrt{s_n}}{\ln s_n} \right] \\ &= \frac{1}{s_n^{1/4}} \frac{\left(1 - s_n^{3/2}\right) \frac{\ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right)}{\ln s_n} - \left(\left(\frac{1+s_n}{2}\right)^{3/2} - s_n^{3/4}\right)}{\frac{1+s_n}{2} - \sqrt{s_n} - \frac{\ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right)}{\ln s_n} (1 - s_n)} \\ &= \frac{1}{s_n^{1/4}} \frac{\left(1 - s_n^{3/2}\right) \ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right) - \left(\left(\frac{1+s_n}{2}\right)^{3/2} - s_n^{3/4}\right) \ln s_n}{\left(\frac{1+s_n}{2} - \sqrt{s_n}\right) \ln s_n - \ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right) (1 - s_n)} \\ &:= \frac{1}{s_n^{1/4}} \frac{f_1(s_n)}{f_2(s_n)}. \end{aligned}$$

From

$$s_{n+1} = \frac{b_{n+1}}{a_{n+1}} = \frac{2\sqrt{s_n}}{1 + s_n},$$

we see that $1 > s_{n+1}(a, b) > s_n(a, b) > 0$, which is due to $\sqrt{x}(1+x) \leq 2$ for $0 < x < 1$. This implies that sequence $\{s_n(a, b)\}_{n=1}^\infty$ is convergent, and it is easily deduced that $\lim_{n \rightarrow \infty} s_n(a, b) = 1$.

To prove $\lim_{n \rightarrow \infty} r_n = \lim_{s_n \rightarrow 1} r_n = \frac{3}{2}$, it suffices to show that $\lim_{s_n \rightarrow 1} [D_n(3/2)/(-D_n(1))] = 0$. To this end, we use power series expansion to get

$$\begin{aligned} f_1(s_n) &= \left(1 - s_n^{3/2}\right) \ln\left(\frac{2\sqrt{s_n}}{1 + s_n}\right) - \left(\left(\frac{1 + s_n}{2}\right)^{3/2} - s_n^{3/4}\right) \ln s_n \\ &= \frac{21}{5 \times 2^{15}}(s_n - 1)^7 + O((s_n - 1)^8), \\ f_2(s_n) &= \left(\frac{1 + s_n}{2} - \sqrt{s_n}\right) \ln s_n - \ln\left(\frac{2\sqrt{s_n}}{1 + s_n}\right) (1 - s_n) \\ &= \frac{1}{384}(s_n - 1)^5 + O((s_n - 1)^6). \end{aligned}$$

Then we obtain

$$\begin{aligned} \lim_{s_n \rightarrow 1} \frac{D_n(3/2)}{-D_n(1)} &= \lim_{s_n \rightarrow 1} \frac{1}{s_n^{1/4}} \frac{f_1(s_n)}{f_2(s_n)} \\ &= \lim_{s_n \rightarrow 1} \frac{\frac{21}{5 \times 2^{15}}(s_n - 1)^7 + O((s_n - 1)^8)}{\frac{1}{384}(s_n - 1)^5 + O((s_n - 1)^6)} = 0. \end{aligned}$$

This completes the proof. \square

3. Main result

Now we are in a position to state and prove our main result.

THEOREM 1. For $n \in \mathbb{N}$, let $t_0^{(n)} = t_0^{(n)}(a, b)$ denote the unique zero of $D_n(t)$ defined by (2) on $(1, 3/2)$. Then we have

$$t_{0s} = \sup_{n \in \mathbb{N}} \{t_0^{(n)}(a, b)\} = \frac{3}{2}, \quad t_{0i} = \inf_{n \in \mathbb{N}} \{t_0^{(n)}(a, b)\} > 1$$

and

$$\inf_{a>0, b>0, a \neq b} \{t_{0i}(a, b)\} = 1.$$

Proof. Lemma 1 tells us that $t \mapsto D_n(t)$ is convex on $[1, \infty)$, which implies that for $t_0^{(n)} \in [1, 3/2]$ we have

$$\frac{D_n(3/2) - D_n(1)}{3/2 - 1} > \frac{D_n(t_0^{(n)}) - D_n(1)}{t_0^{(n)} - 1} = \frac{-D_n(1)}{t_0^{(n)} - 1}.$$

This gives

$$r_n = 1 - \frac{1}{2} \frac{D_n(1)}{D_n(3/2) - D_n(1)} < t_0^{(n)} < \frac{3}{2}.$$

It follows from Lemma 2 that $\lim_{n \rightarrow \infty} t_0^{(n)}(a, b) = 3/2$, and therefore, $t_{0s} = \sup_{n \in \mathbb{N}} \{t_0^{(n)}(a, b)\} = \frac{3}{2}$.

Denote by

$$\begin{aligned} F(s_n, t) &= \left(\frac{2\sqrt{s_n}}{1+s_n}\right)^t \ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right) \times D_n(t) \\ &= \left(\ln\left(\frac{2\sqrt{s_n}}{1+s_n}\right)\right) \left(\frac{2}{1+s_n}\right)^t \left(\frac{s_n^t - 1}{\ln s_n}\right) - \left(\frac{2\sqrt{s_n}}{1+s_n}\right)^t + 1. \end{aligned}$$

Put $t_0 = \inf_{a>0, b>0, a \neq b} t_{0i}(a, b)$. Let $\{s_k\}_{k=1}^\infty$ be a sequence such that $0 < s_{k+1} < s_k < 1$, $\lim_{k \rightarrow \infty} s_k = 0$. Let $\{t(s_k)\}_{k=1}^\infty$ be a sequence of solutions $F(s_k, t(s_k)) = 0$ in $(1, 3/2)$. Then $\{t(s_k)\}_{k=1}^\infty$ is a bounded sequence. It implies that the sequence has a convergent subsequence $\{t(s_{k_l})\}_{l=1}^\infty$. Denote by $t_0^* = \lim_{l \rightarrow \infty} t(s_{k_l})$. It is clear that

$$D_n(t(a, b)) = 0 \iff F(s_n(a, b), t(a, b)) = 0.$$

So from $\lim_{l \rightarrow \infty} F(s_{k_l}, t(s_{k_l})) = 0$ we obtain $2^{t_0^*-1} = 1$. It implies $t_0 = 1$. \square

CONCLUSION 1. For all $a, b > 0$ with $a \neq b$ there is $t_1 \in (1, t_{0i}(a, b))$ such that

$$L_{t_1}(a, b) < AGM(a, b),$$

but $\sup_{a>0, b>0, a\neq b} \{t_1(a, b)\} = 1$.

For all $a, b > 0$ with $a \neq b$ there is no $t_2 \in (1, 3/2)$ such that

$$L_{t_2}(a, b) > AGM(a, b).$$

So $L_1(a, b)$ and $L_{3/2}(a, b)$ are optimal lower and upper bounds for $AGM(a, b)$, respectively.

Acknowledgements. The work was supported by VEGA grant No. 1/0649/17 and by KEGA grant No. 007TnUAD-4/2017. The author thanks to the faculty FPT TnUAD in Púchov, Slovakia for its kind support and he is deeply grateful to the unknown reviewers for their valuable remarks and suggestions.

Competing interests. The author declares that he has no competing interests.

REFERENCES

- [1] J. M. BORWEIN AND P. B. BORWEIN, *Pi and the AGM*, John Wiley and Sons, New York (1987).
- [2] J. M. BORWEIN AND P. B. BORWEIN, *Inequalities for compound mean iterations with logarithmic asymptotes*, J. Math. Anal. Appl. **177** (1993), no. 2, 572–582.
- [3] T. BURIČ, N. ELEZOVIĆ, *Asymptotic expansion of the arithmetic-geometric mean and related inequalities*, Journal of Mathematical Inequalities **9** (2015), 1181–1190.
- [4] B. C. CARLSON AND M. VUORINEN, *An inequality of the AGM and the logarithmic mean*, SIAM Rev. **33** (1991), 655, Problem 91–117.
- [5] E. NEUMAN AND J. SÁNDOR, *On certain means of two arguments and their extensions*, Intern. J. Math. Math. Sci. **2003**, no. 16, 981–993.
- [6] M. K. VAMANAMURTHY AND M. VUORINEN, *Inequalities for means*, J. Math. Anal. Appl. **183** (1994), no. 1, 155–166.
- [7] ZH. H. YANG, *Some identities for means and applications*, RGMIA Res. Rep. Coll. **8** (2005), no. 3 Art. 17; available online at <http://rgmia.vu.edu.au/v8n3.html>.
- [8] ZH. H. YANG, *A New Proof of Inequalities for Gauss Compound Mean*, Int. Journal of Math. Analysis, Vol. **4**, (2010), no. 21, 1013–1018.
- [9] ZH. H. YANG, Y. Q. SONG AND Y. M. CHU, *Sharp bounds for the arithmetic-geometric mean*, J. Inequal. Appl. (2014): **192**, available online at <http://www.journalofinequalitiesandapplications.com/content/2014/1/192>.

(Received October 11, 2016)

Ladislav Matejíčka
 Faculty of Industrial Technologies in Púchov
 Trenčín University of Alexander Dubček in Trenčín
 I. Krasku 491/30, 02001 Púchov, Slovakia
 e-mail: ladislav.matejicka@tnuni.sk