

## ON REVERSES OF THE GOLDEN–THOMPSON TYPE INEQUALITIES

MOHAMMAD BAGHER GHAEMI, VENUS KALEIBARY AND SHIGERU FURUICHI

(Communicated by Y. Seo)

*Abstract.* In this paper we present some reverses of the Golden-Thompson type inequalities: Let  $H$  and  $K$  be Hermitian matrices such that  $e^s e^H \preceq_{ols} e^K \preceq_{ols} e^t e^H$  for some scalars  $s \leq t$ , and  $\alpha \in [0, 1]$ . Then for all  $p > 0$  and  $k = 1, 2, \dots, n$

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq (\max\{S(e^{sp}), S(e^{tp})\})^{\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}},$$

where  $A \#_{\alpha} B = A^{\frac{1}{2}}(A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-\frac{1}{2}})^{\alpha} A^{\frac{1}{2}}$  is  $\alpha$ -geometric mean,  $S(t)$  is the so called Specht ratio and  $\preceq_{ols}$  is the so called Olson order. The same inequalities are also provided with other constants. The obtained inequalities improve some known results.

### 1. Introduction

In what follows, capital letters  $A, B, H$  and  $K$  stand for  $n \times n$  matrices or bounded linear operators on an  $n$ -dimensional complex Hilbert space  $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ . For a pair  $A, B$  of Hermitian matrices, we say  $A \leq B$  if  $B - A \geq 0$ . Let  $A$  and  $B$  be two positive definite matrices. For each  $\alpha \in [0, 1]$ , the weighted geometric mean  $A \#_{\alpha} B$  of  $A$  and  $B$  in the sense of Kubo-Ando [9] is defined by

$$A \#_{\alpha} B = A^{\frac{1}{2}}(A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-\frac{1}{2}})^{\alpha} A^{\frac{1}{2}}.$$

Also for positive semidefinite matrices  $A$  and  $B$ , the weak log-majorization  $A \prec_{wlog} B$  means that

$$\prod_{j=1}^k \lambda_j(A) \leq \prod_{j=1}^k \lambda_j(B), \quad k = 1, 2, \dots, n,$$

where  $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$  are the eigenvalues of  $A$  listed in decreasing order. If equality holds when  $k = n$ , we have the log-majorization  $A \prec_{log} B$ . It is known that the weak log-majorization  $A \prec_{wlog} B$  implies  $\|A\|_u \leq \|B\|_u$  for any unitarily invariant norm  $\|\cdot\|_u$ , i.e.  $\|UAV\|_u = \|A\|_u$  for all  $A$  and all unitaries  $U, V$ . See [2] for theory of majorization.

*Mathematics subject classification* (2010): 15A42, 15A60, 47A63.

*Keywords and phrases:* Ando-Hiai inequality, Golden-Thompson inequality, eigenvalue inequality, geometric mean, Olson order, Specht ratio, generalized Kantorovich constant, unitarily invariant norm.

The author (S. F.) was partially supported by JSPS KAKENHI Grant Number 16K05257.

In [14], Specht obtained an inequality for the arithmetic and geometric means of positive numbers: Let  $x_1 \geq \dots \geq x_n > 0$  and set  $t = x_1/x_n$ . Then

$$\frac{x_1 + \dots + x_n}{n} \leq S(t)(x_1 \dots x_n)^{\frac{1}{n}},$$

where

$$S(t) = \frac{(t-1)t^{1/(t-1)}}{e \log t} \quad (t \neq 1) \quad \text{and} \quad S(1) = 1 \tag{1}$$

is called the Specht ratio at  $t$ . Note that  $\lim_{p \rightarrow 0} S(t^p)^{\frac{1}{p}} = 1$ ,  $S(t^{-1}) = S(t) > 1$  for  $t \neq 1$ ,  $t > 0$  [5]. Specht’s inequality is a ratio type reverse inequality of the classical arithmetic-geometric mean inequality. Using this nice ratio we can state our main result in Section 2.

The Golden-Thompson trace inequality, which is of importance in statistical mechanics and in the theory of random matrices, states that  $Tr e^{H+K} \leq Tr e^H e^K$  for arbitrary Hermitian matrices  $H$  and  $K$ . This inequality has been complemented in several ways [1, 8]. Ando and Hiai in [1] proved that for every unitarily invariant norm  $\|\cdot\|_u$  and  $p > 0$

$$\|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u \leq \|e^{(1-\alpha)H + \alpha K}\|_u. \tag{2}$$

Seo in [12] found some upper bounds on  $\|e^{(1-\alpha)H + \alpha K}\|_u$  in terms of scalar multiples of  $\|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u$ , which show reverse of the Golden-Thompson type inequality (2). In this paper we establish another reverses of this inequality, which improve and refine Seo’s results. In fact the general sandwich condition  $sA \leq B \leq tA$  for positive definite matrices, is the key for our statements. Also, the so called Olson order  $\preceq_{ols}$  is used. For positive operators,  $A \preceq_{ols} B$  if and only if  $A^r \leq B^r$  for every  $r \geq 1$  [11]. Our results are parallel to eigenvalue inequalities obtained in [3] and [7].

### 2. Reverse inequalities via Specht ratio

To study the Golden-Thompson inequality, Ando-Hiai in [1] developed the following log-majorizations:

$$A^r \#_{\alpha} B^r \prec_{\log} (A \#_{\alpha} B)^r, \quad r \geq 1,$$

or equivalently

$$(A^p \#_{\alpha} B^p)^{\frac{1}{p}} \prec_{\log} (A^q \#_{\alpha} B^q)^{\frac{1}{q}}, \quad 0 < q \leq p.$$

There are some literatures [13] on the converse of these inequalities in terms of unitarily invariant norm  $\|\cdot\|_u$ . By the following lemmas, we obtain a new reverse of these inequalities in terms of eigenvalue inequalities.

LEMMA 1. Let  $A$  and  $B$  be positive definite matrices such that  $sA \leq B \leq tA$  for some scalars  $0 < s \leq t$ , and  $\alpha \in [0, 1]$ . Then

$$A^r \sharp_{\alpha} B^r \leq (\max\{S(s), S(t)\})^r (A \sharp_{\alpha} B)^r, \quad 0 < r \leq 1, \quad (3)$$

where  $S(t)$  is the Specht ratio defined as (1).

*Proof.* Let  $f$  be an operator monotone function on  $[0, \infty)$ . Then according to the proof of Theorem 1 in [6], we have

$$f(A) \sharp_{\alpha} f(B) \leq f(M(A \sharp_{\alpha} B)),$$

where  $M = \max\{S(s), S(t)\}$ . Putting  $f(t) = t^r$  for  $0 < r \leq 1$ , we reach inequality (3).  $\square$

LEMMA 2. Let  $A$  and  $B$  be positive definite matrices such that  $sA \preceq_{ols} B \preceq_{ols} tA$  for some scalars  $0 < s \leq t$ , and  $\alpha \in [0, 1]$ . Then

$$\lambda_k(A \sharp_{\alpha} B)^r \leq \max\{S(s^r), S(t^r)\} \lambda_k(A^r \sharp_{\alpha} B^r), \quad r \geq 1, \quad (4)$$

and hence,

$$\lambda_k(A^q \sharp_{\alpha} B^q)^{\frac{1}{q}} \leq (\max\{S(s^p), S(t^p)\})^{\frac{1}{p}} \lambda_k(A^p \sharp_{\alpha} B^p)^{\frac{1}{p}}, \quad 0 < q \leq p, \quad (5)$$

where  $S(t)$  is the Specht ratio defined as (1) and  $k = 1, 2, \dots, n$ .

*Proof.* First note that the condition  $sA \preceq_{ols} B \preceq_{ols} tA$  is equivalent to the condition  $s^v A^v \leq B^v \leq t^v A^v$  for every  $v \geq 1$ . In particular, we have  $sA \leq B \leq tA$  for  $v = 1$ . Also, for  $r \geq 1$  we have  $0 < \frac{1}{r} \leq 1$  and by (3)

$$A^{\frac{1}{r}} \sharp_{\alpha} B^{\frac{1}{r}} \leq (\max\{S(s), S(t)\})^{\frac{1}{r}} (A \sharp_{\alpha} B)^{\frac{1}{r}}. \quad (6)$$

On the other hand, from the condition  $s^v A^v \leq B^v \leq t^v A^v$  for every  $v \geq 1$  and letting  $v = r$ , we have

$$s^r A^r \leq B^r \leq t^r A^r.$$

Now if we let  $X = A^r$ ,  $Y = B^r$ ,  $w = s^r$  and  $z = t^r$ , then

$$wX \leq Y \leq zX. \quad (7)$$

Using (6) under the condition (7), we have

$$X^{\frac{1}{r}} \sharp_{\alpha} Y^{\frac{1}{r}} \leq (\max\{S(w), S(z)\})^{\frac{1}{r}} (X \sharp_{\alpha} Y)^{\frac{1}{r}},$$

and this is the same as

$$A \sharp_{\alpha} B \leq (\max\{S(s^r), S(t^r)\})^{\frac{1}{r}} (A^r \sharp_{\alpha} B^r)^{\frac{1}{r}}.$$

Hence

$$\lambda_k(A \#_{\alpha} B) \leq (\max\{S(s^r), S(t^r)\})^{\frac{1}{r}} \lambda_k(A^r \#_{\alpha} B^r)^{\frac{1}{r}}.$$

By taking  $r$ -th power on both sides and using the spectral mapping theorem, we get the desired inequality (4). Note that from the minimax characterization of eigenvalues of a Hermitian matrix [2] it follows immediately that  $A \leq B$  implies  $\lambda_k(A) \leq \lambda_k(B)$  for each  $k$ . Similarly since  $p/q \geq 1$ , from inequality (4)

$$\lambda_k(A \#_{\alpha} B)^{\frac{p}{q}} \leq \max\{S(s^{\frac{p}{q}}), S(t^{\frac{p}{q}})\} \lambda_k(A^{\frac{p}{q}} \#_{\alpha} B^{\frac{p}{q}}). \tag{8}$$

Replacing  $A$  and  $B$  by  $A^q$  and  $B^q$  in (8), and using the sandwich condition  $s^q A^q \leq B^q \leq t^q A^q$ , we have

$$\lambda_k(A^q \#_{\alpha} B^q)^{\frac{p}{q}} \leq \max\{S(s^p), S(t^p)\} \lambda_k(A^p \#_{\alpha} B^p).$$

This completes the proof.  $\square$

Note that eigenvalue inequalities immediately imply log-majorization and unitarily invariant norm inequalities.

**COROLLARY 1.** *Let  $A$  and  $B$  be positive definite matrices such that  $mI \leq A, B \leq MI$  for some scalars  $0 < m \leq M$  with  $h = M/m$ , and let  $\alpha \in [0, 1]$ . Then*

$$A^r \#_{\alpha} B^r \leq S(h)^r (A \#_{\alpha} B)^r, \quad 0 < r \leq 1, \tag{9}$$

and hence

$$\lambda_k(A \#_{\alpha} B)^r \leq S(h^r) \lambda_k(A^r \#_{\alpha} B^r), \quad r \geq 1, \tag{10}$$

$$\lambda_k(A^q \#_{\alpha} B^q)^{\frac{1}{q}} \leq S(h^p)^{\frac{1}{p}} \lambda_k(A^p \#_{\alpha} B^p)^{\frac{1}{p}}, \quad 0 < q \leq p, \tag{11}$$

where  $S(t)$  is the Specht ratio defined as (1) and  $k = 1, 2, \dots, n$ .

*Proof.* Since  $mI \leq A, B \leq MI$  implies  $\frac{m}{M}A \leq B \leq \frac{M}{m}A$ , the inequality (9) is obtained by letting  $s = m/M$ ,  $t = M/m$  in Lemma 1. Also from  $mI \leq A, B \leq MI$ , we have  $m^{\nu}I \leq A^{\nu}, B^{\nu} \leq M^{\nu}I$  for every  $\nu \geq 1$ , and so

$$\left(\frac{m}{M}\right)^{\nu} A^{\nu} \leq B^{\nu} \leq \left(\frac{M}{m}\right)^{\nu} A^{\nu}. \tag{12}$$

Using Lemma 2 under the condition (12), we reach inequalities (10) and (11). Note that  $S(h) = S(\frac{1}{h})$  for every  $h > 0$ .  $\square$

**REMARK 1.** We remark that the matrix inequality (9) is more stronger than corresponding norm inequality obtained by Seo in [12, Corollary 3.2]. Also, inequality (11) is presented in [12, Lemma 3.1].

In the sequel we show a reverse of the Golden-Thompson type inequality (2), which is our main result.

**THEOREM 1.** *Let  $H$  and  $K$  be Hermitian matrices such that  $e^s e^H \preceq_{ols} e^K \preceq_{ols} e^t e^H$  for some scalars  $s \leq t$ , and let  $\alpha \in [0, 1]$ . Then for all  $p > 0$ ,*

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq (\max\{S(e^{sp}), S(e^{tp})\})^{\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}},$$

where  $S(t)$  is the Specht ratio defined as (1) and  $k = 1, 2, \dots, n$ .

*Proof.* Replacing  $A$  and  $B$  by  $e^H$  and  $e^K$  in the inequality (5) of Lemma 2, we can write

$$\lambda_k(e^{qH} \#_{\alpha} e^{qK})^{\frac{1}{q}} \leq (\max\{S(e^{sp}), S(e^{tp})\})^{\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}, \quad 0 < q \leq p.$$

By [8, Lemma 3.3], we have

$$e^{(1-\alpha)H+\alpha K} = \lim_{q \rightarrow 0} (e^{qH} \#_{\alpha} e^{qK})^{\frac{1}{q}},$$

and hence it follows that for each  $p > 0$ ,

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq (\max\{S(e^{sp}), S(e^{tp})\})^{\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}. \quad \square$$

**COROLLARY 2.** *Let  $H$  and  $K$  be Hermitian matrices such that  $e^s e^H \preceq_{ols} e^K \preceq_{ols} e^t e^H$  for some scalars  $s \leq t$ , and let  $\alpha \in [0, 1]$ . Then for every unitarily invariant norm  $\|\cdot\|_u$  and all  $p > 0$ ,*

$$\|e^{(1-\alpha)H+\alpha K}\|_u \leq (\max\{S(e^{sp}), S(e^{tp})\})^{\frac{1}{p}} \|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u, \quad (13)$$

and the right-hand side of (13) converges to the left-hand side as  $p \downarrow 0$ . In particular,

$$\|e^{H+K}\|_u \leq \max\{S(e^{2s}), S(e^{2t})\} \|(e^{2H} \#_{\alpha} e^{2K})\|_u.$$

**COROLLARY 3.** [12, Theorem 3.3–Theorem 3.4] *Let  $H$  and  $K$  be Hermitian matrices such that  $mI \leq H, K \leq MI$  for some scalars  $m \leq M$ , and let  $\alpha \in [0, 1]$ . Then for all  $p > 0$ ,*

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq S(e^{(M-m)p})^{\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}, \quad k = 1, 2, \dots, n.$$

So, for every unitarily invariant norm  $\|\cdot\|_u$

$$\|e^{(1-\alpha)H+\alpha K}\|_u \leq S(e^{(M-m)p})^{\frac{1}{p}} \|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u,$$

and the right-hand side of these inequalities converges to the left-hand side as  $p \downarrow 0$ .

*Proof.* From  $mI \leq H, K \leq MI$ , we have  $e^{vm} \leq e^{vH}, e^{vK} \leq e^{vM}$  for every  $v \geq 1$  and so we can derive  $e^{m-M} e^H \preceq_{ols} e^K \preceq_{ols} e^{M-m} e^H$ . Now the assertion is obtained by applying Theorem 1 and the fact that for every  $t > 0$ ,  $S(t) = S(\frac{1}{t})$ .  $\square$

### 3. Reverse inequalities via Kantorovich constant

A well-known matrix version of the Kantorovich inequality [10] asserts that if  $A$  and  $U$  are two matrices such that  $0 < mI \leq A \leq MI$  and  $UU^* = I$ , then

$$UA^{-1}U^* \leq \frac{(m+M)^2}{4mM}(UAU^*)^{-1}. \tag{14}$$

Let  $w > 0$ . The generalized Kantorovich constant  $K(w, \alpha)$  is defined by

$$K(w, \alpha) := \frac{(w^\alpha - w)}{(\alpha - 1)(w - 1)} \left( \frac{\alpha - 1}{\alpha} \frac{w^\alpha - 1}{w^\alpha - w} \right)^\alpha, \tag{15}$$

for any real number  $\alpha \in \mathbb{R}$  [5]. In fact,  $K(\frac{M}{m}, -1) = K(\frac{M}{m}, 2)$  is the constant occurring in (14).

Now as a result of the following statement, we have another reverse Golden-Thompson type inequality which refines corresponding inequality in [12].

**PROPOSITION 1.** [7, Theorem 3] Let  $H$  and  $K$  be Hermitian matrices such that  $e^s e^H \preceq_{ols} e^K \preceq_{ols} e^t e^H$  for some scalars  $s \leq t$ , and let  $\alpha \in [0, 1]$ . Then

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq K(e^{p(t-s)}, \alpha)^{-\frac{1}{p}} \lambda_k(e^{pH} \#_\alpha e^{pK})^{\frac{1}{p}}, \quad p > 0, \tag{16}$$

where  $K(w, \alpha)$  is the generalized Kantorovich constant defined as (15).

**THEOREM 2.** Let  $H$  and  $K$  be Hermitian matrices such that  $mI \leq K, H \leq MI$  for some scalars  $m \leq M$  and let  $\alpha \in [0, 1]$ . Then for every  $p > 0$ ,

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} \lambda_k(e^{pH} \#_\alpha e^{pK})^{\frac{1}{p}}, \quad k = 1, 2, \dots, n,$$

and the right-hand side of this inequality converges to the left-hand side as  $p \downarrow 0$ . In particular,

$$\lambda_k(e^{H+K}) \leq \frac{e^{2M} + e^{2m}}{2e^M e^m} \lambda_k(e^{2H} \# e^{2K}), \quad k = 1, 2, \dots, n.$$

*Proof.* Since  $mI \leq K, H \leq MI$  implies  $e^{m-M} e^H \preceq_{ols} e^K \preceq_{ols} e^{M-m} e^H$ , desired inequalities are obtained by letting  $s = m - M$  and  $t = M - m$  in Proposition 1. For the convergence, we know that  $\frac{2w^{\frac{1}{4}}}{w^{\frac{1}{2}} + 1} \leq K(w, \alpha) \leq 1$ , for every  $\alpha \in [0, 1]$ . So, for every  $p > 0$

$$1 \leq K(w^p, \alpha)^{-\frac{1}{p}} \leq \left( \frac{2w^{\frac{p}{4}}}{w^{\frac{p}{2}} + 1} \right)^{-\frac{1}{p}}.$$

A simple calculation shows that

$$\lim_{p \rightarrow 0} -\frac{1}{p} \log \left( \frac{2w^{\frac{p}{4}}}{w^{\frac{p}{2}} + 1} \right) = \lim_{p \rightarrow 0} \frac{(w^{\frac{p}{2}} - 1) \log(w)}{4(w^{\frac{p}{2}} + 1)} = 0,$$

and hence  $\lim_{p \rightarrow 0} \left( \frac{2w^{\frac{p}{4}}}{w^{\frac{p}{2}} + 1} \right)^{-\frac{1}{p}} = 1$ . Now by using the sandwich condition and letting  $w = e^{2(M-m)}$ , we have  $\lim_{p \rightarrow 0} K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} = 1$ .  $\square$

REMARK 2. Under the assumptions of Theorem 2, Seo in [12, Theorem 4.2] proved that

$$\|e^{(1-\alpha)H + \alpha K}\|_u \leq K(e^{(M-m)}, p)^{-\frac{\alpha}{p}} K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} \|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u, \quad 0 < p \leq 1,$$

and

$$\|e^{(1-\alpha)H + \alpha K}\|_u \leq K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} \|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u, \quad p \geq 1.$$

But the first inequality of Theorem 2 shows that the sharper constant for all  $p > 0$  is  $K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}}$ . Since for  $0 < p \leq 1$ ,  $K(e^{(M-m)}, p)^{-\frac{\alpha}{p}} \geq 1$  and hence

$$K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} \leq K(e^{(M-m)}, p)^{-\frac{\alpha}{p}} K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}}.$$

#### 4. Some related results

It has been shown [6] that if  $f : [0, \infty) \rightarrow [0, \infty)$  is operator monotone function and  $0 < ml \leq A \leq B \leq MI \leq I$  with  $h = \frac{M}{m}$ , then for all  $\alpha \in [0, 1]$

$$f(A) \#_{\alpha} f(B) \leq \exp \left( \alpha(1 - \alpha) \left( 1 - \frac{1}{h} \right)^2 \right) f(A \#_{\alpha} B). \tag{17}$$

This new ratio has been introduced by Furuichi and Minculete in [4], which is different from Specht ratio and Kantorovich constant. By applying (17) for  $f(t) = t^r$ ,  $0 < r \leq 1$  we have the following results similar to Lemma 1 and Lemma 2.

LEMMA 3. Let  $A$  and  $B$  be positive definite matrices such that  $0 < ml \leq A \leq B \leq MI \leq I$  with  $h = M/m$ , and let  $\alpha \in [0, 1]$ . Then

$$A^r \#_{\alpha} B^r \leq \exp \left( r\alpha(1 - \alpha) \left( 1 - \frac{1}{h} \right)^2 \right) (A \#_{\alpha} B)^r, \quad 0 < r \leq 1.$$

LEMMA 4. Let  $A$  and  $B$  be positive definite matrices such that  $0 < ml \preceq_{ots} A \preceq_{ots} B \preceq_{ots} MI \preceq_{ots} I$  with  $h = M/m$ , and let  $\alpha \in [0, 1]$ . Then for all  $k = 1, 2, \dots, n$ ,

$$\lambda_k(A \#_{\alpha} B)^r \leq \exp \left( \alpha(1 - \alpha) \left( 1 - \frac{1}{h^r} \right)^2 \right) \lambda_k(A^r \#_{\alpha} B^r), \quad r \geq 1,$$

$$\lambda_k(A^q \#_{\alpha} B^q)^{\frac{1}{q}} \leq \exp \left( \frac{1}{p} \alpha(1 - \alpha) \left( 1 - \frac{1}{h^p} \right)^2 \right) \lambda_k(A^p \#_{\alpha} B^p)^{\frac{1}{p}}, \quad 0 < q \leq p. \tag{18}$$

**THEOREM 3.** *Let  $H$  and  $K$  be Hermitian matrices such that  $e^m I \preceq_{ols} e^H \preceq_{ols} e^K \preceq_{ols} e^M I \preceq_{ols} I$  for some scalars  $m \leq M$ , and let  $\alpha \in [0, 1]$ . Then for all  $p > 0$  and  $k = 1, 2, \dots, n$*

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq \exp\left(\frac{1}{p}\alpha(1-\alpha)\left(1 - \frac{1}{e^{p(M-m)}}\right)^2\right) \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}},$$

and so, for every unitarily invariant norm  $\|\cdot\|_u$

$$\|e^{(1-\alpha)H+\alpha K}\|_u \leq \exp\left(\frac{1}{p}\alpha(1-\alpha)\left(1 - \frac{1}{e^{p(M-m)}}\right)^2\right) \|(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}\|_u.$$

*Proof.* The proof is similar to that of Theorem 1, by replacing  $A$  and  $B$  with  $e^H$  and  $e^K$ , and  $h = e^{M-m}$  in the inequality (18).  $\square$

**REMARK 3.** Under the different conditions, the different coefficients are not comparable. But it is known that if we have a certain statement under the sandwich condition  $0 < sA \leq B \leq tA$ , then the same statement is also true under the condition  $0 < ml \leq A, B \leq MI$  and  $0 < ml \leq A \leq B \leq MI \leq I$ . Hence, we can compare the following special cases:

(1) Comparison of the constants in Theorem 3 and in Theorem 2:

Let  $e^m I \preceq_{ols} e^H \preceq_{ols} e^K \preceq_{ols} e^M I \preceq_{ols} I$ . Operator monotony of  $\log(t)$  leads to  $ml \leq H \leq K \leq MI \leq I$ , and so  $ml \leq H, K \leq MI$ . Now by applying Theorem 2 we have

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq K(e^{2p(M-m)}, \alpha)^{-\frac{1}{p}} \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}, \quad p > 0.$$

Also, by Theorem 3

$$\lambda_k(e^{(1-\alpha)H+\alpha K}) \leq \exp\left(\frac{1}{p}\alpha(1-\alpha)\left(1 - \frac{1}{e^{p(M-m)}}\right)^2\right) \lambda_k(e^{pH} \#_{\alpha} e^{pK})^{\frac{1}{p}}, \quad p > 0.$$

Letting  $h = e^{M-m} \geq 1$ , the following numerical examples show that there is no ordering between these inequalities.

(i) Take  $\alpha = \frac{1}{2}$ ,  $p = \frac{1}{2}$  and  $h = 2$ , then we have

$$K(h^{2p}, \alpha)^{-\frac{1}{p}} - \exp\left(\frac{1}{p}\alpha(1-\alpha)\left(1 - \frac{1}{h^p}\right)^2\right) \simeq -0.0134963.$$

(ii) Take  $\alpha = \frac{1}{2}$ ,  $p = \frac{1}{2}$  and  $h = 8$ , then we have

$$K(h^{2p}, \alpha)^{-\frac{1}{p}} - \exp\left(\frac{1}{p}\alpha(1-\alpha)\left(1 - \frac{1}{h^p}\right)^2\right) \simeq 0.0631159.$$

(2) Comparison of the constants in Lemma 3 and in Lemma 1:

Let  $0 < ml \leq A \leq B \leq MI \leq I$ . Then the following sandwich condition is obtained

$$m \leq \frac{m}{M} \leq 1 \leq A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \leq \frac{M}{m} \leq \frac{1}{m}.$$



Now by letting  $s = 1$  and  $t = \frac{M}{m} = h$  in Lemma 1, we get

$$A^r \sharp_{\alpha} B^r \leq S(h)^r (A \sharp_{\alpha} B)^r, \quad 0 < r \leq 1. \quad (19)$$

Also, by Lemma 3

$$A^r \sharp_{\alpha} B^r \leq \exp\left(r\alpha(1-\alpha)\left(1-\frac{1}{h}\right)^2\right) (A \sharp_{\alpha} B)^r, \quad 0 < r \leq 1. \quad (20)$$

It is shown in [4, Remark 2.4] that there is no ordering between coefficients of (19) and (20). Therefore, we may conclude evaluation of Lemma 3 and Lemma 1 are different.

#### REFERENCES

- [1] T. ANDO AND F. HIAI, *Log-majorization and complementary Golden-Thompson type inequalities*, Linear Algebra Appl. **197/198** (1994), 113–131.
- [2] R. BHATIA, *Matrix Analysis*, Grad. Texts in Math., vol. **169**, Springer-Verlag, 1997.
- [3] J.-C. BOURIN AND Y. SEO, *Reverse inequality to Golden-Thompson type inequalities: comparison of  $e^{A+B}$  and  $e^{Ae^B}$* , Linear Algebra Appl. **426** (2007), 312–316.
- [4] S. FURUICHI AND N. MINCULETE, *Alternative reverse inequalities for Young's inequality*, J. Math. Inequal. **5** (2011), 595–600.
- [5] T. FURUTA, J. MIĆIĆ, J. E. PEČARIĆ AND Y. SEO, *Mond-Pečarić method in operator inequalities*, Monographs in Inequalities 1, Element, Zagreb, 2005.
- [6] M. B. GHAEMI AND V. KALEIBARY, *Some inequalities involving operator monotone functions and operator means*, Math. Inequal. Appl. **19** (2016), 757–764.
- [7] M. B. GHAEMI AND V. KALEIBARY, *Eigenvalue inequalities related to the Ando-Hiai inequality*, Math. Inequal. Appl. **20** (2017), 217–223.
- [8] F. HIAI AND D. PETZ, *The Golden-Thompson trace inequality is complemented*, Linear Algebra Appl. **181** (1993), 153–185.
- [9] F. KUBO AND T. ANDO, *Means of positive linear operators*, Math. Ann. **246** (1980), 205–224.
- [10] A. W. MARSHALL AND I. OLKIN, *Matrix versions of the Cauchy and Kantorovich inequalities*, Aequationes Math. **40** (1990), 89–93.
- [11] M. P. OLSON, *The selfadjoint operators of a von Neumann algebra from a conditionally complete lattice*, Proc. Amer. Math. Soc. **28** (1971), 537–544.
- [12] Y. SEO, *Reverses of the Golden-Thompson type inequalities due to Ando-Hiai-Petz*, Banach J. Math. Anal. **2** (2008), 140–149.
- [13] Y. SEO, *On a reverse of Ando-Hiai inequality*, Banach J. Math. Anal. **4** (2010), 87–91.
- [14] W. SPECHT, *Zur Theorie der elementaren Mittel*, Math. Z. **74** (1960), 91–98.

(Received July 28, 2016)

Mohammad Bagher Ghaemi  
School of Mathematics  
Iran University of Science and Technology  
Narmak, Tehran 16846-13114, Iran  
e-mail: mghaemi@iust.ac.ir

Venus Kaleibary  
School of Mathematics  
Iran University of Science and Technology  
Narmak, Tehran 16846-13114, Iran  
e-mail: v.kaleibary@gmail.com

Shigeru Furuichi  
Department of Information Science  
College of Humanities and Sciences, Nihon University  
3-25-40, Sakurajyousui, Setagaya-ku, Tokyo 156-8550, Japan  
e-mail: furuichi@chs.nihon-u.ac.jp