

## IMPROVEMENTS OF GENERALIZED HÖLDER'S INEQUALITIES AND THEIR APPLICATIONS

JING-FENG TIAN, MING-HU HA\* AND CHAO WANG

(Communicated by J. Pečarić)

*Abstract.* In this paper, we present some new improvements of generalized Hölder's inequalities, and then we obtain a new refinement of Minkowski inequality. Moreover, the obtained results are used to improve Chung's inequality and Beckenbach-type inequality proposed by Wang.

### 1. Introduction

The classical Hölder inequality states that if  $a_k \geq 0$ ,  $b_k \geq 0$  ( $k = 1, 2, \dots, n$ ),  $p \geq q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , then

$$\sum_{k=1}^n a_k b_k \leq \left( \sum_{k=1}^n a_k^p \right)^{\frac{1}{p}} \left( \sum_{k=1}^n b_k^q \right)^{\frac{1}{q}}. \quad (1)$$

The sign of inequality (1) is reversed for  $p < 0$ . (For  $p < 0$ , we assume that  $a_k, b_k > 0$ .) Hölder's inequality is of cardinal importance in mathematical analysis and in the field of applied mathematics. Due to the importance of Hölder's inequality (1), it has received considerable attention by many authors, and has motivated a large number of research papers giving it various generalizations, improvements and applications (e.g. [1, 2, 3, 7, 9, 10, 11, 12, 13, 14, 15, 18, 19, 20, 21, 22, 23]). Among various generalizations of (1), Vasić and Pečarić in [16] presented the following interesting theorem.

**THEOREM A.** Let  $A_{ij} > 0$  ( $i = 1, 2, \dots, n$ ,  $j = 1, 2, \dots, m$ ).

(a) If  $\beta_j \geq 0$ , and if  $\sum_{j=1}^m \beta_j \geq 1$ , then

$$\sum_{i=1}^n \prod_{j=1}^m A_{ij}^{\beta_j} \leq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij} \right)^{\beta_j}. \quad (2)$$

(b) If  $\beta_j \leq 0$  ( $j = 1, 2, \dots, m$ ), then

$$\sum_{i=1}^n \prod_{j=1}^m A_{ij}^{\beta_j} \geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij} \right)^{\beta_j}. \quad (3)$$

*Mathematics subject classification* (2010): 26D15, 26D10.

*Keywords and phrases:* Hölder's inequality, Minkowski inequality, Beckenbach-type inequality, Chung's inequality.

\* The corresponding author.

(c) If  $\beta_1 > 0$ ,  $\beta_j \leq 0$  ( $j = 2, 3, \dots, m$ ), and if  $\sum_{j=1}^m \beta_j \leq 1$ , then

$$\sum_{i=1}^n \prod_{j=1}^m A_{ij}^{\beta_j} \geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij} \right)^{\beta_j}. \tag{4}$$

The above inequalities are called as generalized Hölder’s inequalities.

The main objective of this paper is to establish some new improvements of inequalities (1)–(4), and their applications. The rest of this paper is organized as follows. In Section 2, we present a series of sharpened versions of inequalities (1)–(4). And then we obtain a new refinement of Minkowski inequality. In Section 3, we apply the results to improve Chung’s inequality and Beckenbach-type inequality proposed by Wang.

### 2. Improvements of generalized Hölder’s inequalities

We begin this section with four lemmas, which will be used in the sequel.

LEMMA 2.1. Let  $0 < X_{ij} < 1$  ( $i = 1, 2, \dots, n$ ,  $j = 1, 2, \dots, m$ ),  $m \geq 2$ , let  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m > 0$ ,  $\sum_{j=1}^m \frac{1}{\lambda_j} \geq 1$  such that  $1 - \sum_{i=1}^n X_{ij}^{\lambda_j} > 0$ , and let  $X_{i(m+1)} = X_{i1}$ ,  $\lambda_{m+1} = \lambda_1$ . Then

$$\prod_{j=1}^m \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} + \sum_{i=1}^n \prod_{j=1}^m X_{ij} \leq \prod_{j=1}^m \left[ 1 - \left( \sum_{i=1}^n X_{ij}^{\lambda_j} - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^2 \right]^{\frac{1}{2\lambda_j}}. \tag{5}$$

*Proof.* From the hypotheses of Lemma 2.1, it is easy to see that

$$\frac{1}{\lambda_m} \geq \frac{1}{\lambda_{m-1}} \geq \dots \geq \frac{1}{\lambda_2} \geq \frac{1}{\lambda_1} > 0,$$

and

$$\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j} \geq 0 \quad (j = 1, 2, \dots, m - 1).$$

Noting that  $(\frac{1}{2\lambda_2} - \frac{1}{2\lambda_1}) + \frac{1}{2\lambda_1} + \frac{1}{2\lambda_1} + (\frac{1}{2\lambda_3} - \frac{1}{2\lambda_2}) + \frac{1}{2\lambda_2} + \frac{1}{2\lambda_2} + \dots + (\frac{1}{2\lambda_m} - \frac{1}{2\lambda_{m-1}}) + \frac{1}{2\lambda_{m-1}} + \frac{1}{2\lambda_{m-1}} + (\frac{1}{2\lambda_m} - \frac{1}{2\lambda_1}) + \frac{1}{2\lambda_1} + \frac{1}{2\lambda_1} = \frac{1}{\lambda_1} + \frac{1}{\lambda_2} + \dots + \frac{1}{\lambda_m} \geq 1$ , from the inequality (2), we get

$$\begin{aligned} & \prod_{j=1}^m \left[ 1 - \left( \sum_{i=1}^n X_{ij}^{\lambda_j} - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &= \prod_{j=1}^m \left\{ \left[ \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right) + \sum_{i=1}^n X_{ij}^{\lambda_j} \right]^{\frac{1}{2\lambda_j}} \left[ \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right) + \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right]^{\frac{1}{2\lambda_j}} \right. \\ & \quad \left. \times \left[ \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right) + \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right]^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right\} \end{aligned}$$

$$\begin{aligned}
 &\geq \prod_{j=1}^m \left\{ \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_j}} \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{2\lambda_j}} \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right\} \\
 &\quad + \prod_{j=1}^m \left[ \left( X_{1j}^{\lambda_j} \right)^{\frac{1}{2\lambda_j}} \left( X_{1(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_j}} \left( X_{1(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \\
 &\quad + \prod_{j=1}^m \left[ \left( X_{2j}^{\lambda_j} \right)^{\frac{1}{2\lambda_j}} \left( X_{2(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_j}} \left( X_{2(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \\
 &\quad \dots \\
 &\quad + \prod_{j=1}^m \left[ \left( X_{nj}^{\lambda_j} \right)^{\frac{1}{2\lambda_j}} \left( X_{n(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_j}} \left( X_{n(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \\
 &= \prod_{j=1}^m \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} + \sum_{i=1}^n \prod_{j=1}^m X_{ij}, \tag{6}
 \end{aligned}$$

and then (5) follows. The proof of Lemma 2.1 is completed.  $\square$

LEMMA 2.2. *Let  $X_{ij} > 1$  ( $i = 1, 2, \dots, n, j = 1, 2, \dots, m$ ),  $m \geq 2$ , let  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_m < 0$  such that  $1 - \sum_{i=1}^n X_{ij}^{\lambda_j} > 0$ , and let  $X_{i(m+1)} = X_{i1}$ ,  $\lambda_{m+1} = \lambda_1$ . Then*

$$\prod_{j=1}^m \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} + \sum_{i=1}^n \prod_{j=1}^m X_{ij} \geq \prod_{j=1}^m \left[ 1 - \left( \sum_{i=1}^n X_{ij}^{\lambda_j} - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^2 \right]^{\frac{1}{2\lambda_j}}. \tag{7}$$

*Proof.* By the same method as in Lemma 2.1, and using inequality (3), we can deduce the desired inequality (7).  $\square$

LEMMA 2.3. *Let  $0 < X_{im} < 1$ ,  $X_{ij} > 1$  ( $i = 1, 2, \dots, n, j = 1, 2, \dots, m-1$ ),  $m \geq 2$ , let  $\lambda_m > 0$ ,  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{m-1} < 0$ ,  $\sum_{j=1}^m \frac{1}{\lambda_j} \leq 1$  such that  $1 - \sum_{i=1}^n X_{ij}^{\lambda_j} > 0$ , and let  $X_{i(m+1)} = X_{i1}$ ,  $\lambda_{m+1} = \lambda_1$ . Then*

$$\prod_{j=1}^m \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} + \sum_{i=1}^n \prod_{j=1}^m X_{ij} \geq \prod_{j=1}^{m-1} \left[ 1 - \left( \sum_{i=1}^n X_{ij}^{\lambda_j} - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^2 \right]^{\frac{1}{2\lambda_j}}. \tag{8}$$

*Proof.* In view of the assumptions in Lemma 2.3, we have

$$\frac{1}{\lambda_m} > 0, \frac{1}{\lambda_{m-1}} \leq \frac{1}{\lambda_{m-2}} \leq \dots \leq \frac{1}{\lambda_2} \leq \frac{1}{\lambda_1} < 0$$

and

$$\frac{1}{2\lambda_j} - \frac{1}{2\lambda_{j-1}} \leq 0 \quad (j = 2, 3, \dots, m-1), \quad \frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}} > 0.$$

Consequently, according to  $\frac{1}{2\lambda_1} + \frac{1}{2\lambda_1} + \left( \frac{1}{2\lambda_2} - \frac{1}{2\lambda_1} \right) + \frac{1}{2\lambda_2} + \frac{1}{2\lambda_2} + \left( \frac{1}{2\lambda_3} - \frac{1}{2\lambda_2} \right) + \dots + \frac{1}{2\lambda_{m-2}} + \frac{1}{2\lambda_{m-2}} + \left( \frac{1}{2\lambda_{m-1}} - \frac{1}{2\lambda_{m-2}} \right) + \frac{1}{2\lambda_{m-1}} + \frac{1}{2\lambda_{m-1}} + \left( \frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}} \right) + \frac{1}{2\lambda_1} = \frac{1}{\lambda_1} +$

$\frac{1}{\lambda_2} + \dots + \frac{1}{\lambda_m} \leq 1$ , by using inequality (4), we obtain

$$\begin{aligned}
 & \prod_{j=1}^{m-1} \left[ 1 - \left( \sum_{i=1}^n X_{ij}^{\lambda_j} - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\
 &= \left\{ \prod_{j=1}^{m-1} \left[ \sum_{i=1}^n X_{ij}^{\lambda_j} + \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right) \right]^{\frac{1}{2\lambda_j}} \right\} \\
 & \quad \times \left\{ \prod_{j=1}^{m-1} \left[ \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} + \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right) \right]^{\frac{1}{2\lambda_j}} \right\} \\
 & \quad \times \left\{ \prod_{j=1}^{m-2} \left[ \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} + \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right) \right]^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right\} \\
 & \quad \times \left[ \sum_{i=1}^n X_{im}^{\lambda_m} + \left( 1 - \sum_{i=1}^n X_{im}^{\lambda_m} \right) \right]^{\frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}}} \left[ \sum_{i=1}^n X_{i1}^{\lambda_1} + \left( 1 - \sum_{i=1}^n X_{i1}^{\lambda_1} \right) \right]^{\frac{1}{2\lambda_1}} \\
 & \leq \left\{ \prod_{j=1}^{m-2} \left[ (X_{1j}^{\lambda_j})^{\frac{1}{2\lambda_j}} (X_{1(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_j}} (X_{1(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \right\} \\
 & \quad \times (X_{1(m-1)}^{\lambda_{m-1}})^{\frac{1}{2\lambda_{m-1}}} (X_{1m}^{\lambda_m})^{\frac{1}{2\lambda_{m-1}}} (X_{1m}^{\lambda_m})^{\frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}}} (X_{11}^{\lambda_1})^{\frac{1}{2\lambda_1}} \\
 & \quad + \left\{ \prod_{j=1}^{m-2} \left[ (X_{2j}^{\lambda_j})^{\frac{1}{2\lambda_j}} (X_{2(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_j}} (X_{2(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \right\} \\
 & \quad \times (X_{2(m-1)}^{\lambda_{m-1}})^{\frac{1}{2\lambda_{m-1}}} (X_{2m}^{\lambda_m})^{\frac{1}{2\lambda_{m-1}}} (X_{2m}^{\lambda_m})^{\frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}}} (X_{21}^{\lambda_1})^{\frac{1}{2\lambda_1}} \\
 & \quad + \dots \\
 & \quad + \left\{ \prod_{j=1}^{m-2} \left[ (X_{nj}^{\lambda_j})^{\frac{1}{2\lambda_j}} (X_{n(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_j}} (X_{n(j+1)}^{\lambda_{j+1}})^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \right\} \\
 & \quad \times (X_{n(m-1)}^{\lambda_{m-1}})^{\frac{1}{2\lambda_{m-1}}} (X_{nm}^{\lambda_m})^{\frac{1}{2\lambda_{m-1}}} (X_{nm}^{\lambda_m})^{\frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}}} (X_{n1}^{\lambda_1})^{\frac{1}{2\lambda_1}} \\
 & \quad + \left\{ \prod_{j=1}^{m-2} \left[ \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_j}} \left( 1 - \sum_{i=1}^n X_{ij}^{\lambda_j} \right)^{\frac{1}{2\lambda_j}} \left( 1 - \sum_{i=1}^n X_{i(j+1)}^{\lambda_{j+1}} \right)^{\frac{1}{2\lambda_{j+1}} - \frac{1}{2\lambda_j}} \right] \right\} \\
 & \quad \times \left( 1 - \sum_{i=1}^n X_{im}^{\lambda_m} \right)^{\frac{1}{2\lambda_{m-1}}} \left( 1 - \sum_{i=1}^n X_{i(m-1)}^{\lambda_{m-1}} \right)^{\frac{1}{2\lambda_{m-1}}} \\
 & \quad \times \left( 1 - \sum_{i=1}^n X_{im}^{\lambda_m} \right)^{\frac{1}{\lambda_m} - \frac{1}{2\lambda_{m-1}}} \left( 1 - \sum_{i=1}^n X_{i1}^{\lambda_1} \right)^{\frac{1}{2\lambda_1}} \\
 & = \prod_{j=1}^m X_j + \prod_{j=1}^m (1 - X_j^{\lambda_j})^{\frac{1}{\lambda_j}}. \tag{9}
 \end{aligned}$$

The proof of Lemma 2.3 is completed.  $\square$

LEMMA 2.4. [5] If  $x > -1$ ,  $\alpha > 1$  or  $\alpha < 0$ , then

$$(1+x)^\alpha \geq 1 + \alpha x. \tag{10}$$

The inequality is reversed for  $0 \leq \alpha \leq 1$ .

Next, we present new improvements of inequalities (2), (3) and (4).

THEOREM 2.5. Let  $A_{ij} \geq 0$  ( $i = 1, 2, \dots, n$ ,  $j = 1, 2, \dots, m$ ), let  $s$  be any given natural number ( $1 \leq s \leq n$ ), and let  $m \geq 2$ ,  $\lambda_{m+1} = \lambda_1$ ,  $A_{i(m+1)} = A_{i1}$ .

(a) If  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m > 0$ ,  $\sum_{j=1}^m \frac{1}{\lambda_j} \geq 1$ , then

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\leq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^m \left[ 1 - \left( \frac{A_{sj}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{s(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\leq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{11}$$

(b) If  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_m < 0$ , then

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\geq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^m \left[ 1 - \left( \frac{A_{sj}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{s(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{12}$$

(c) If  $\lambda_m > 0$ ,  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{m-1} < 0$ , and if  $\sum_{j=1}^m \frac{1}{\lambda_j} \leq 1$ , then

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\geq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^{m-1} \left[ 1 - \left( \frac{A_{sj}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{s(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{13}$$

*Proof.* (a). For any given natural number  $s$  ( $1 \leq s \leq n$ ), from the following substitution:

$$X_{ij} = \left( \frac{A_{ij}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} \right)^{\frac{1}{\lambda_j}} \quad (i = 1, 2, \dots, n, j = 1, 2, \dots, m), \tag{14}$$

we have

$$0 < X_{ij} < 1, \quad 1 - \sum_{1 \leq i \leq n, i \neq s} X_{ij}^{\lambda_j} > 0.$$

Consequently, by using the substitution (14) and inequality (5), we find

$$\prod_{j=1}^m \left[ 1 - \sum_{1 \leq i \leq n, i \neq s} \left( \frac{A_{ij}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} \right)^{\frac{1}{\lambda_j}} \right]^{\frac{1}{\lambda_j}} + \sum_{1 \leq i \leq n, i \neq s} \left( \prod_{j=1}^m \frac{A_{ij}}{\left( \sum_{k=1}^n A_{kj}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}} \right) \leq \prod_{j=1}^m \left\{ 1 - \left[ \sum_{1 \leq i \leq n, i \neq s} \left( \frac{A_{ij}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} \right) - \sum_{1 \leq i \leq n, i \neq s} \left( \frac{A_{i(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right) \right]^2 \right\}^{\frac{1}{2\lambda_j}}, \tag{15}$$

and then we obtain

$$\frac{\prod_{j=1}^m A_{sj}}{\prod_{j=1}^m \left( \sum_{k=1}^n A_{kj}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}} + \frac{\sum_{1 \leq i \leq n, i \neq s} \prod_{j=1}^m A_{ij}}{\prod_{j=1}^m \left( \sum_{k=1}^n A_{kj}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}} \leq \prod_{j=1}^m \left[ 1 - \left( \frac{A_{sj}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{s(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}}, \tag{16}$$

which implies

$$\frac{\sum_{i=1}^n \left( \prod_{j=1}^m A_{ij} \right)}{\prod_{j=1}^m \left( \sum_{k=1}^n A_{kj}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}} \leq \prod_{j=1}^m \left[ 1 - \left( \frac{A_{sj}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{s(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}}. \tag{17}$$

So, the desired inequality (11) is valid.

(b). In view of

$$X_{ij} = \left( \frac{A_{ij}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} \right)^{\frac{1}{\lambda_j}} > 1$$

for all  $\lambda_j < 0$  ( $j = 1, 2, \dots, m$ ), and

$$1 - \sum_{1 \leq i \leq n, i \neq s} X_{ij}^{\lambda_j} > 0,$$

using similar reasoning as in Case (a), but applying Lemma 2.2 in place of Lemma 2.1, we immediately obtain inequality (12).

(c). Noting that

$$0 < X_{im} = \left( \frac{A_{im}^{\lambda_m}}{\sum_{k=1}^n A_{km}^{\lambda_m}} \right)^{\frac{1}{\lambda_m}} < 1, \quad X_{ij} = \left( \frac{A_{ij}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} \right)^{\frac{1}{\lambda_j}} > 1$$

for  $\lambda_m > 0, \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{m-1} < 0$ , and

$$1 - \sum_{1 \leq i \leq n, i \neq s} X_{ij}^{\lambda_j} > 0,$$

making the similar methods as in Case (a), but using Lemma 2.3 in place of Lemma 2.1, we get inequality (13).

The proof of Theorem 2.5 is completed.  $\square$

Letting  $s = 1$  in (11), (12) and (13), respectively, we obtain the following corollary.

**COROLLARY 2.6.** *Let  $A_{ij} \geq 0$  ( $i = 1, 2, \dots, n, j = 1, 2, \dots, m$ ), let  $m \geq 2$ , and let  $\lambda_{m+1} = \lambda_1, A_{i(m+1)} = A_{i1}$ .*

(a) *If  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m > 0$ , and if  $\sum_{j=1}^m \frac{1}{\lambda_j} \geq 1$ , then*

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\leq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^m \left[ 1 - \left( \frac{A_{1j}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{1(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\leq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{18}$$

(b) *If  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_m < 0$ , then*

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\geq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^m \left[ 1 - \left( \frac{A_{1j}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{1(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{19}$$

(c) *If  $\lambda_m > 0, \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{m-1} < 0, \sum_{j=1}^m \frac{1}{\lambda_j} \leq 1$ , then*

$$\begin{aligned} \sum_{i=1}^n \prod_{j=1}^m A_{ij} &\geq \left[ \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}} \right] \times \prod_{j=1}^{m-1} \left[ 1 - \left( \frac{A_{1j}^{\lambda_j}}{\sum_{k=1}^n A_{kj}^{\lambda_j}} - \frac{A_{1(j+1)}^{\lambda_{j+1}}}{\sum_{k=1}^n A_{k(j+1)}^{\lambda_{j+1}}} \right)^2 \right]^{\frac{1}{2\lambda_j}} \\ &\geq \prod_{j=1}^m \left( \sum_{i=1}^n A_{ij}^{\lambda_j} \right)^{\frac{1}{\lambda_j}}. \end{aligned} \tag{20}$$

In particular, putting  $m = 2, s = 1, \lambda_1 = p, \lambda_2 = q, A_{i1} = a_i, A_{i2} = b_i$  in Corollary 2.6, we have the new improvements of Hölder’s inequality as follows.

**COROLLARY 2.7.** *Let  $a_i \geq 0, b_i \geq 0 (i = 1, 2, \dots, n)$ . If  $p \geq q > 0, \frac{1}{p} + \frac{1}{q} = 1$ , then*

$$\begin{aligned} \sum_{i=1}^n a_i b_i &\leq \left( \sum_{i=1}^n a_i^p \right)^{\frac{1}{p}} \left( \sum_{i=1}^n b_i^q \right)^{\frac{1}{q}} \\ &\quad \times \left[ 1 - \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{b_1^q}{\sum_{k=1}^n b_k^q} \right)^2 \right]^{\frac{1}{2p}} \left[ 1 - \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{b_1^q}{\sum_{k=1}^n b_k^q} \right)^2 \right]^{\frac{1}{2q}} \end{aligned}$$

$$\begin{aligned}
 &= \left( \sum_{i=1}^n a_i^p \right)^{\frac{1}{p}} \left( \sum_{i=1}^n b_i^q \right)^{\frac{1}{q}} \left[ 1 - \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{b_1^q}{\sum_{k=1}^n b_k^q} \right)^2 \right]^{\frac{1}{2}} \\
 &\leq \left( \sum_{i=1}^n a_i^p \right)^{\frac{1}{p}} \left( \sum_{i=1}^n b_i^q \right)^{\frac{1}{q}}.
 \end{aligned} \tag{21}$$

If  $p < 0, q > 0, \frac{1}{p} + \frac{1}{q} = 1$ , then

$$\sum_{i=1}^n a_i b_i \geq \left( \sum_{i=1}^n a_i^p \right)^{\frac{1}{p}} \left( \sum_{i=1}^n b_i^q \right)^{\frac{1}{q}} \left[ 1 - \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{b_1^q}{\sum_{k=1}^n b_k^q} \right)^2 \right]^{\frac{1}{2p}}. \tag{22}$$

Moreover, from Corollary 2.7, we get the following refinement of Minkowski inequality.

COROLLARY 2.8. Let  $a_k > 0, b_k > 0 (k = 1, 2, \dots, n)$ .

(I) If  $p > 1$ , then

$$\begin{aligned}
 \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{\frac{1}{p}} &\leq \left( \sum_{k=1}^n a_k^p \right)^{\frac{1}{p}} + \left( \sum_{k=1}^n b_k^p \right)^{\frac{1}{p}} \\
 &\quad - \frac{1}{2} \left( \sum_{k=1}^n a_k^p \right)^{\frac{1}{p}} \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \\
 &\quad - \frac{1}{2} \left( \sum_{k=1}^n b_k^p \right)^{\frac{1}{p}} \left( \frac{b_1^p}{\sum_{k=1}^n b_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2.
 \end{aligned} \tag{23}$$

(II) If  $0 < p < 1$ , then

$$\begin{aligned}
 \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{1/p} &\geq \left( \sum_{k=1}^n a_k^p \right)^{1/p} + \left( \sum_{k=1}^n b_k^p \right)^{1/p} \\
 &\quad + \frac{1-p}{2p} \left( \sum_{k=1}^n a_k^p \right)^{1/p} \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \\
 &\quad + \frac{1-p}{2p} \left( \sum_{k=1}^n b_k^p \right)^{1/p} \left( \frac{b_1^p}{\sum_{k=1}^n b_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2.
 \end{aligned} \tag{24}$$

Proof. (I). For  $p > 1$ , we write

$$\sum_{k=1}^n (a_k + b_k)^p = \sum_{k=1}^n a_k (a_k + b_k)^{p-1} + \sum_{k=1}^n b_k (a_k + b_k)^{p-1},$$



and using inequality 21, we have

$$\begin{aligned} \sum_{k=1}^n (a_k + b_k)^p &\leq \left( \sum_{k=1}^n a_k^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{\frac{p-1}{p}} \\ &\quad \times \left[ 1 - \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \right]^{\frac{1}{2}} \\ &\quad + \left( \sum_{k=1}^n b_k^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{\frac{p-1}{p}} \\ &\quad \times \left[ 1 - \left( \frac{b_1^p}{\sum_{k=1}^n b_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \right]^{\frac{1}{2}}. \end{aligned} \quad (25)$$

Then, by using Lemma 2.4, we obtain

$$\begin{aligned} \sum_{k=1}^n (a_k + b_k)^p &\leq \left( \sum_{k=1}^n a_k^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{\frac{p-1}{p}} \\ &\quad \times \left[ 1 - \frac{1}{2} \left( \frac{a_1^p}{\sum_{k=1}^n a_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \right] \\ &\quad + \left( \sum_{k=1}^n b_k^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^n (a_k + b_k)^p \right]^{\frac{p-1}{p}} \\ &\quad \times \left[ 1 - \frac{1}{2} \left( \frac{b_1^p}{\sum_{k=1}^n b_k^p} - \frac{(a_1 + b_1)^p}{\sum_{k=1}^n (a_k + b_k)^p} \right)^2 \right]. \end{aligned}$$

Dividing both sides by  $\left( \sum_{k=1}^n (a_k + b_k)^p \right)^{\frac{p-1}{p}}$ , we obtain the desired inequality (23).

(II). For  $0 < p < 1$ , by inequality (22) and in the same way, we obtained the desired inequality (24). The proof of Corollary 2.8 is completed.  $\square$

### 3. Applications

In this section, we provide two applications of our new inequalities. Firstly, we give here a refined and generalized version of Chung's inequality.

The famous Chung inequality [6] states that if  $a_1 \geq a_2 \geq \dots \geq a_n \geq 0$  and  $\sum_{j=1}^k a_j \leq \sum_{j=1}^k b_j$  for  $k = 1, 2, \dots, n$ , then

$$\sum_{j=1}^n a_j^2 \leq \sum_{j=1}^n b_j^2. \quad (26)$$

**THEOREM 3.1.** *Let  $a_1 \geq a_2 \geq \dots \geq a_n \geq 0$ , and let  $\sum_{j=1}^k a_j \leq \sum_{j=1}^k b_j$  for  $k = 1, 2, \dots, n$ . Then, for  $p > 1$  we have*

$$\sum_{j=1}^n a_j^p \leq \left( \sum_{j=1}^n b_j^p \right) \left[ 1 - \left( \frac{a_1^p}{\sum_{j=1}^n a_j^p} - \frac{b_1^p}{\sum_{j=1}^n b_j^p} \right)^2 \right]^{\frac{p}{2}} \leq \sum_{j=1}^n b_j^2. \tag{27}$$

*Proof.* From the hypotheses in Theorem 3.1 we find that

$$\begin{aligned} \sum_{j=1}^n a_j^p &= \sum_{k=1}^{n-1} \left[ \left( a_k^{p-1} - a_{k+1}^{p-1} \right) \sum_{j=1}^k a_j \right] + a_n^{p-1} \sum_{j=1}^n a_j \\ &\leq \sum_{k=1}^{n-1} \left[ \left( a_k^{p-1} - a_{k+1}^{p-1} \right) \sum_{j=1}^k b_j \right] + a_n^{p-1} \sum_{j=1}^n b_j \\ &= \sum_{j=1}^n a_j^{p-1} b_j. \end{aligned} \tag{28}$$

By using inequality (21), we have

$$\sum_{j=1}^n a_j^{p-1} b_j \leq \left( \sum_{j=1}^n a_j^p \right)^{\frac{p-1}{p}} \left( \sum_{j=1}^n b_j^p \right)^{\frac{1}{p}} \left[ 1 - \left( \frac{a_1^p}{\sum_{j=1}^n a_j^p} - \frac{b_1^p}{\sum_{j=1}^n b_j^p} \right)^2 \right]^{\frac{1}{2}}. \tag{29}$$

Hence, combining inequalities (28) and (29) yields inequality (27). The proof of Theorem 3.1 is completed.  $\square$

Nextly, we present a new refinement of Beckenbach-type inequality proposed by Wang. The famous Beckenbach inequality [4] has been generalized and refined in several directions (e.g. [8]). In [17], Wang obtained an interesting Beckenbach-type inequality as follows.

**THEOREM B.** *Let  $f(x)$ ,  $g(x)$  be positive integrable functions defined on  $[0, T]$ , and let  $a, b, c$  be positive numbers. If  $p \geq q > 0$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , then*

$$\frac{\left( a + c \int_0^T \varphi^p(x) dx \right)^{\frac{1}{p}}}{b + c \int_0^T \varphi(x) g(x) dx} \leq \frac{\left( a + c \int_0^T f^p(x) dx \right)^{\frac{1}{p}}}{b + c \int_0^T f(x) g(x) dx}, \tag{30}$$

where  $\varphi(x) = \left( \frac{ag(x)}{b} \right)^{\frac{q}{p}}$ .

**THEOREM 3.2.** *Let  $f(x)$ ,  $g(x)$  be positive integrable functions defined on  $[0, T]$ , and let  $a, b, c$  be positive numbers. If  $p \geq q > 0$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , then*

$$\begin{aligned} \frac{\left( a + c \int_0^T \varphi^p(x) dx \right)^{\frac{1}{p}}}{b + c \int_0^T \varphi(x) g(x) dx} &\leq \frac{\left( a + c \int_0^T f^p(x) dx \right)^{\frac{1}{p}}}{b + c \int_0^T f(x) g(x) dx} \\ &\times \left[ 1 - \frac{1}{2} \left( \frac{a}{a + c \int_0^T f^p(x) dx} - \frac{a^{-\frac{q}{p}} b^q}{a^{-\frac{q}{p}} b^q + c \int_0^T g^q(x) dx} \right)^2 \right], \end{aligned} \tag{31}$$

where  $\varphi(x) = \left(\frac{ag(x)}{b}\right)^{\frac{q}{p}}$ .

*Proof.* A simple calculation gives

$$\frac{\left(a + c \int_0^T \varphi^p(x) dx\right)^{\frac{1}{p}}}{b + c \int_0^T \varphi(x)g(x) dx} = \left(a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx\right)^{-\frac{1}{q}}. \quad (32)$$

By using Hölder's inequality 1 and inequality 21 we get

$$\begin{aligned} b + c \int_0^T f(x)g(x) dx &\leq b + c \left(\int_0^T f^p(x) dx\right)^{\frac{1}{p}} \left(\int_0^T g^q(x) dx\right)^{\frac{1}{q}} \\ &= a^{\frac{1}{p}}(ba^{-\frac{1}{p}}) + c \left(\int_0^T f^p(x) dx\right)^{\frac{1}{p}} \left(\int_0^T g^q(x) dx\right)^{\frac{1}{q}} \\ &\leq \left(a + c \int_0^T f^p(x) dx\right)^{\frac{1}{p}} \left(a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx\right)^{\frac{1}{q}} \\ &\quad \times \left[1 - \left(\frac{a}{a + c \int_0^T f^p(x) dx} - \frac{a^{-\frac{q}{p}}b^q}{a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx}\right)^2\right]^{\frac{1}{2}}, \end{aligned} \quad (33)$$

which implies

$$\begin{aligned} &\left(a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx\right)^{-\frac{1}{q}} \\ &\leq \frac{\left(a + c \int_0^T f^p(x) dx\right)^{\frac{1}{p}}}{b + c \int_0^T f(x)g(x) dx} \\ &\quad \times \left[1 - \left(\frac{a}{a + c \int_0^T f^p(x) dx} - \frac{a^{-\frac{q}{p}}b^q}{a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx}\right)^2\right]^{\frac{1}{2}}. \end{aligned} \quad (34)$$

Moreover, from Lemma 2.4 we have

$$\begin{aligned} &\left(a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx\right)^{-\frac{1}{q}} \\ &\leq \frac{\left(a + c \int_0^T f^p(x) dx\right)^{\frac{1}{p}}}{b + c \int_0^T f(x)g(x) dx} \\ &\quad \times \left[1 - \frac{1}{2} \left(\frac{a}{a + c \int_0^T f^p(x) dx} - \frac{a^{-\frac{q}{p}}b^q}{a^{-\frac{q}{p}}b^q + c \int_0^T g^q(x) dx}\right)^2\right]. \end{aligned} \quad (35)$$

Hence, combining inequalities (32) and (35) yields inequality (31). The proof of Theorem 3.2 is completed.  $\square$

*Acknowledgements.* The authors would like to express their sincere thanks to the anonymous referees for their great efforts to improve this paper.

This work was supported by the NNSF of China (No. 11626079), the Application Basic Research Plan Key Basic Research Project of Hebei Province of China (No. 16964213D), the Scientific Research Project of Higher Education of Hebei Province (No. BJ2017031), the Fundamental Research Funds for the Central Universities (No. 2015ZD29) and the Higher School Science Research Funds of Hebei Province of China (No. Z2015137).

#### REFERENCES

- [1] H. AGAHI, R. MESIAR, *Probability inequalities for decomposition integrals*, J. Comput. Appl. Math. **315** (2017), 240–248.
- [2] H. AGAHI, Y. OUYANG, R. MESIAR, E. PAP, M. ŠTRBOJA, *Hölder and Minkowski type inequalities for pseudo-integral*, Appl. Math. Comput. **217** (2011), 8630–8639.
- [3] H. AGAHI, R. MESIAR, Y. OUYANG, *On some advanced type inequalities for Sugeno integral and  $T(S)$ -evaluators*, Inform. Sci. **190** (2012), 64–75.
- [4] E. F. BECKENBACH, *On Hölder's inequality*, J. Math. Anal. Appl. **15** (1966), 21–29.
- [5] E. F. BECKENBACH, R. BELLMAN, *Inequalities*, Springer-Verlag, Berlin, (1983).
- [6] K. L. CHUNG, *Proposed Problem E1025*, Amer. Math. Monthly **59** (1952), 407–408.
- [7] K. J. FALCONER, *A Hölder-type inequality on a regular rooted tree*, J. Math. Anal. Appl. **423** (2015), 913–923.
- [8] J. KUANG, *Applied Inequalities*, Shandong Science Press, Jinan, (2010).
- [9] S. MEDIĆ, T. GRBIĆ, A. PEROVIĆ, S. NIKOLIČIĆ, *Inequalities of Hölder and Minkowski type for pseudo-integrals with respect to interval-valued  $\oplus$ -measures*, Fuzzy Set. Syst. **304** (2016), 110–130.
- [10] J.-F. TIAN, *New property of a generalized Hölder's inequality and its applications*, Inform. Sci. **288** (2014), 45–54.
- [11] J.-F. TIAN, M.-H. HA, *Properties of generalized sharp Hölder's inequalities*, J. Math. Inequal. **11** (2017), 511–525.
- [12] J.-F. TIAN, M.-H. HA, *Properties and refinements of Aczél-type inequalities*, J. Math. Inequal. **12** (2018), 175–189.
- [13] J.-F. TIAN, M.-H. HA, *Extensions of Hölder-type inequalities on time scales and their applications*, J. Nonlinear Sci. Appl. **10** (2017), 937–953.
- [14] J.-F. TIAN, W. PEDRYCZ, *New refinements of generalized Hölder's inequality and their applications*, Math. Inequal. Appl. **19** (2016), 805–822.
- [15] J.-F. TIAN, ZH.-H. YANG, *Generalizations of Hu-type inequalities and their applications*, J. Nonlinear Sci. Appl. **10** (2017), 1971–1985.
- [16] P. M. VASIĆ, J. E. PEČARIĆ, *On the Jensen inequality for monotone functions*, An Univ Timișoara Ser Șt Matematică **17** (1979), 95–104.
- [17] C. L. WANG, *Characteristics of nonlinear positive functionals and their applications*, J. Math. Anal. Appl. **95** (1983), 564–574.
- [18] L. M. WU, J. SUN, X. YE, L. ZHU, *Hölder type inequality for Sugeno integral*, Fuzzy Set. Syst. **161** (2010), 2337–2347.
- [19] ZH.-H. YANG, J. TIAN, *Monotonicity and sharp inequalities related to gamma function*, J. Math. Inequal. **12** (2018), 1–22.
- [20] ZH.-H. YANG, J. TIAN, *Monotonicity and inequalities for the gamma function*, J. Inequal. Appl. (2017) 2017:317.
- [21] ZH.-H. YANG, J. TIAN, *Optimal inequalities involving power-exponential mean, arithmetic mean and geometric mean*, J. Math. Inequal. **11** (2017), 1169–1183.

- [22] C. ZHAO, W. CHEUNG, *Hölder's reverse inequality and its applications*, Publications de L'institut Mathématique **99** (2016), 211–216.
- [23] C. ZHAO, W. CHEUNG, *On reverse Minkowski-type inequalities*, Mediterr. J. Math. **12** (2015), 1085–1094.

(Received August 2, 2017)

*Jing-Feng Tian*  
College of Science and Technology  
North China Electric Power University  
Baoding, Hebei Province, 071051, P. R. China  
e-mail: tianjf@ncepu.edu.cn

*Ming-Hu Ha*  
School of Science  
Hebei University of Engineering  
Handan, Hebei Province, 056038, P. R. China  
e-mail: mhhbu@163.com

*Chao Wang*  
School of Management  
Hebei University of Engineering  
Handan, Hebei Province, 056038, P. R. China  
e-mail: wang222chao@163.com