

SOME k -FRACTIONAL ASSOCIATES OF HERMITE–HADAMARD’S INEQUALITY FOR QUASI-CONVEX FUNCTIONS AND APPLICATIONS TO SPECIAL MEANS

R. HUSSAIN, A. ALI, A. LATIF AND G. GULSHAN

(Communicated by M. Andrić)

Abstract. This article brings together some inequalities associated with Hermite-Hadamard’s inequality for quasi-convex functions by way of k -Riemann-Liouville fractional integrals of order α . The inequalities thus obtained are applied to some special means of real numbers.

1. Introduction

A function $f : I \subset \mathbb{R} \longrightarrow \mathbb{R}$ is said to be convex on I if for every $a, b \in I$ and $t \in [0, 1]$, we have

$$f(ta + (1-t)b) \leq tf(a) + (1-t)f(b).$$

An immediate consequence of convexity is the integrability of the function in the Riemann’s sense. Subsequently the lower and upper estimations for the integral average of a convex function defined on the compact interval $[a, b]$, involving the midpoint and the endpoints of the domain, are given by the following inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t)dt \leq \frac{f(a)+f(b)}{2}.$$

This is the celebrated Hermite-Hadamard’s inequality. Almost since the outset of this inequality in 1881 by Hermite (see [4]), it has been worked on. As a result many generalizations, refinements, extensions and counter part of this inequality are available in literature. So persisting the tradition of generalization, this inequality is generalized here for quasi-convex functions by means of remarkable k -Riemann-Liouville fractional integrals.

Mathematics subject classification (2010): 26D15, 26A51, 32F99, 41A17.

Keywords and phrases: Hermite-Hadamard inequality, quasi-convex function, k -Riemann-Liouville fractional integrals, Hölder’s integral inequality, power mean inequality.

DEFINITION 1. The function $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ is said to be quasi-convex if for every $a, b \in I$ and $t \in [0, 1]$, we have

$$f(ta + (1-t)b) \leq \max\{f(a), f(b)\},$$

(see [4]). Quasi-convexity is a weaker convexity, that is it generalizes the notion of convexity. Therefore every convex function is quasi-convex whereas there are quasi-convex functions which are not convex (see [7]).

In [3] Dragomir and Pearce proved the following result for quasi-convex function, connected with Hermite-Hadamard inequality:

LEMMA 1. Let $f : [a, b] \rightarrow \mathbb{R}$ be a quasi-convex function and $f \in L_1[a, b]$, we have the inequality

$$\frac{1}{b-a} \int_a^b f(t) dt \leq \max\{f(a), f(b)\}. \quad (1)$$

In [7] Ion proved the following two results connected with quasi-convex function:

THEOREM 1. Let $f : [a, b] \rightarrow \mathbb{R}$ be differentiable function on (a, b) . If $|f'|$ is quasi-convex on $[a, b]$, the subsequent inequality is valid

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{b-a}{4} \max\{|f'(a)|, |f'(b)|\}. \quad (2)$$

THEOREM 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be differentiable function on (a, b) . If $|f'|^{\frac{p}{p-1}}$ is quasi-convex on $[a, b]$ with $p > 1$, the subsequent inequality is valid

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{b-a}{2(p+1)^{\frac{1}{p}}} \left(\max\{|f'(a)|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}}\} \right)^{\frac{p-1}{p}}. \quad (3)$$

In [1] the following result connected with quasi-convex function is proved:

THEOREM 3. Let $f : I^\circ \subset \mathbb{R} \rightarrow \mathbb{R}$ be differentiable function on I° , $a, b \in I^\circ$ with $a < b$. If $|f'|^q$ is quasi-convex on $[a, b]$, $q \geq 1$, the subsequent inequality is valid

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{b-a}{4} \left(\max\{|f'(a)|^q, |f'(b)|^q\} \right)^{\frac{1}{q}}. \quad (4)$$

In [8] Mubeen and Habibullah introduced the following class k -fractional integrals:

DEFINITION 2. Let $f \in L_1[a, b]$, the k -Riemann-Liouville fractional integrals ${}_k J_{a^+}^\alpha f(u)$ and ${}_k J_{b^-}^\alpha f(u)$ of order $\alpha > 0$ with $a \geq 0$, $k > 0$, are defined by

$${}_k J_{a^+}^\alpha f(u) = \frac{1}{k\Gamma_k(\alpha)} \int_a^u (u-t)^{\frac{\alpha}{k}-1} f(t) dt, \quad 0 \leq a < u < b$$

and

$${}_k J_{b^-}^\alpha f(u) = \frac{1}{k\Gamma_k(\alpha)} \int_u^b (t-u)^{\frac{\alpha}{k}-1} f(t) dt, \quad 0 \leq a < u < b$$

respectively, where $\Gamma_k(\alpha)$ is the k -gamma function given as $\Gamma_k(\alpha) = \int_0^\infty t^{\alpha-1} e^{-\frac{t^k}{k}} dt$, see [2].

In fact k -Riemann-Liouville fractional integrals of order α are generalization of Riemann-Liouville fractional integrals of order α . If we take $k \rightarrow 1$, the k -Riemann-Liouville fractional integrals of order α turn out to be Riemann-Liouville fractional integrals of order α which are described in [5].

In [6] following useful identity related to k -fractional integrals is proved:

LEMMA 2. Let $f : [a, b] \rightarrow R$ be differentiable function on (a, b) . If $f' \in L[a, b]$, the following equality for k -fractional integrals is valid

$$\begin{aligned} & \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \\ &= \frac{b-a}{2} \int_0^1 [(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}] f'(ta + (1-t)b) dt. \end{aligned}$$

2. Main results

The main aim of the present paper is to establish new inequalities for quasi-convex functions via k -Riemann-Liouville fractional integrals. Starting with the following lemma.

LEMMA 3. Let $f : [a, b] \rightarrow R$ be positive function and $f \in L_1[a, b]$. If f is quasi-convex on $[a, b]$, the subsequent inequality for k -fractional integrals is valid

$$\frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \leq \max\{f(a), f(b)\}$$

with $\frac{\alpha}{k} > 0$.

Proof. Since f is quasi-convex on $[a, b]$, we have

$$f(ta + (1-t)b) \leq \max\{f(a), f(b)\}$$

and

$$f((1-t)a+tb) \leq \max\{f(a), f(b)\}$$

by adding these inequalities we get

$$\frac{1}{2}[f(ta+(1-t)b)+f((1-t)a+tb)] \leq \max\{f(a), f(b)\}$$

now multiplying both sides by $t^{\frac{\alpha}{k}-1}$ and then integrating the resulting inequality with respect to t over $[0, 1]$, we obtain

$$\begin{aligned} & \int_0^1 t^{\frac{\alpha}{k}-1} f(ta+(1-t)b) dt + \int_0^1 t^{\frac{\alpha}{k}-1} f((1-t)a+tb) dt \\ &= \int_b^a \left(\frac{b-u}{b-a}\right)^{\frac{\alpha}{k}-1} f(u) \frac{du}{a-b} + \int_a^b \left(\frac{v-a}{b-a}\right)^{\frac{\alpha}{k}-1} f(v) \frac{dv}{b-a} \\ &\leq \frac{2k}{\alpha} \max\{f(a), f(b)\} \end{aligned}$$

by using the definition of k -Riemann-Liouville fractional integrals, we get

$$\frac{\Gamma_k(\alpha+k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \leq \max\{f(a), f(b)\},$$

hence the proof is complete. \square

REMARK 1. If we choose $\alpha = k$ in Lemma 3, with properties of gamma function we have the inequality (1).

THEOREM 4. Let $f : [a, b] \rightarrow R$ be a differentiable function on (a, b) . If $|f'|$ is quasi-convex on $[a, b]$, $\alpha > 0$, the subsequent inequality for k -fractional integrals is valid

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{\Gamma_k(\alpha+k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{b-a}{\left(\frac{\alpha}{k}+1\right)} \left(1 - \frac{1}{2^{\frac{\alpha}{k}}}\right) \max\{|f'(a)|, |f'(b)|\}. \end{aligned}$$

Proof. Using Lemma 2, the fact that $|f'|$ is quasi-convex and properties of modulus, we have

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{\Gamma_k(\alpha+k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{b-a}{2} \int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| |f'(ta+(1-t)b)| dt \\ & \leq \frac{b-a}{2} \int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| \max\{|f'(a)|, |f'(b)|\} dt \end{aligned}$$

$$\begin{aligned}
&= \frac{b-a}{2} \left\{ \int_0^{\frac{1}{2}} \left[(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}} \right] dt + \int_{\frac{1}{2}}^1 \left[t^{\frac{\alpha}{k}} - (1-t)^{\frac{\alpha}{k}} \right] dt \right\} \max \{ |f'(a)|, |f'(b)| \} \\
&= \frac{b-a}{\left(\frac{\alpha}{k} + 1\right)} \left(1 - \frac{1}{2^{\frac{\alpha}{k}}} \right) \max \{ |f'(a)|, |f'(b)| \}.
\end{aligned}$$

Here we have used

$$\begin{aligned}
\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| dt &= \int_0^{\frac{1}{2}} \left[(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}} \right] dt + \int_{\frac{1}{2}}^1 \left[t^{\frac{\alpha}{k}} - (1-t)^{\frac{\alpha}{k}} \right] dt \\
&= \frac{2}{\left(\frac{\alpha}{k} + 1\right)} \left(1 - \frac{1}{2^{\frac{\alpha}{k}}} \right)
\end{aligned}$$

which completes the proof. \square

REMARK 2. If we choose $\alpha = k$ in Theorem 4, we have the inequality (2).

THEOREM 5. Let $f : [a, b] \rightarrow R$ be a differentiable function on (a, b) such that $f' \in L_1[a, b]$. If $|f'|^q$ is quasi-convex on $[a, b]$ and $q > 1$, the subsequent inequality for k -fractional integrals is valid

$$\begin{aligned}
&\left| \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\
&\leq \frac{b-a}{2\left(\frac{\alpha}{k}p + 1\right)^{\frac{1}{p}}} \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}
\end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$ and $\frac{\alpha}{k} \in [0, 1]$.

Proof. From Lemma 2 and using Hölder's inequality with properties of modulus, we have

$$\begin{aligned}
&\left| \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\
&\leq \frac{b-a}{2} \int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| |f'(ta + (1-t)b)| dt \\
&\leq \frac{b-a}{2} \left(\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}}.
\end{aligned}$$

We know that for $\frac{\alpha}{k} \in [0, 1]$ and for all $t_1, t_2 \in [0, 1]$, $\left| t_1^{\frac{\alpha}{k}} - t_2^{\frac{\alpha}{k}} \right| \leq |t_1 - t_2|^{\frac{\alpha}{k}}$, therefore

$$\begin{aligned}
\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}|^p dt &\leq \int_0^1 |1-2t|^{\frac{\alpha}{k}p} dt \\
&= \int_0^{\frac{1}{2}} |1-2t|^{\frac{\alpha}{k}p} dt + \int_{\frac{1}{2}}^1 |2t-1|^{\frac{\alpha}{k}p} dt \\
&= \frac{1}{\frac{\alpha}{k}p + 1}.
\end{aligned}$$

Since $|f'|^q$ is quasi-convex on $[a, b]$, we have

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{b-a}{2\left(\frac{\alpha}{k}p + 1\right)^{\frac{1}{p}}} \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}. \quad \square \end{aligned}$$

REMARK 3. If in Theorem 5, we choose $\alpha = k$, we have the inequality (3).

THEOREM 6. Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable function on (a, b) such that $f' \in L_1[a, b]$. If $|f'|^q$ is quasi-convex on $[a, b]$ and $q \geq 1$, the subsequent inequality for k -fractional integrals is valid

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{b-a}{\left(\frac{\alpha}{k} + 1\right)} \left(1 - \frac{1}{2^{\frac{\alpha}{k}}} \right) \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}} \end{aligned}$$

with $\frac{\alpha}{k} > 0$.

Proof. From Lemma 2, using power mean inequality with properties of modulus and using the fact that $|f'|^q$ is quasi-convex, we have

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{\Gamma_k(\alpha + k)}{2(b-a)^{\frac{\alpha}{k}}} [{}_k J_{a^+}^\alpha f(b) + {}_k J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{b-a}{2} \int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| |f'(ta + (1-t)b)| dt \\ & \leq \frac{b-a}{2} \left(\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| |f'(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b-a}{2} \left(\int_0^1 |(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}| dt \right) \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}} \\ & = \frac{b-a}{2} \left(\int_0^{\frac{1}{2}} [(1-t)^{\frac{\alpha}{k}} - t^{\frac{\alpha}{k}}] dt + \int_{\frac{1}{2}}^1 [t^{\frac{\alpha}{k}} - (1-t)^{\frac{\alpha}{k}}] dt \right) \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}} \\ & = \frac{b-a}{\left(\frac{\alpha}{k} + 1\right)} \left(1 - \frac{1}{2^{\frac{\alpha}{k}}} \right) \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}, \end{aligned}$$

which completes the proof. \square

REMARK 4. If in theorem 6, we choose $\alpha = k$, we have the inequality (4).

3. Applications to special means

We now consider special means of positive real numbers ξ, η ($\xi \neq \eta$), as follows

Arithmetic mean

$$A(\xi, \eta) = \frac{\xi + \eta}{2}.$$

Geometric Mean

$$G(\xi, \eta) = \sqrt{\xi\eta}.$$

Harmonic Mean

$$H(\xi, \eta) = \frac{2\xi\eta}{\xi + \eta}.$$

Logarithmic mean

$$L(\xi, \eta) = \frac{\eta - \xi}{\ln|\eta| - \ln|\xi|}.$$

Generalised Log-mean

$$L_n(\xi, \eta) = \left[\frac{\eta^{n+1} - \xi^{n+1}}{(n+1)(\eta - \xi)} \right]^{\frac{1}{n}}, \quad n \in \mathbb{Z} \setminus \{-1, 0\}.$$

Indentric Mean

$$I(\xi, \eta) = \frac{1}{e} \left(\frac{\eta^\eta}{\xi^\xi} \right)^{\frac{1}{\eta - \xi}}.$$

PROPOSITION 1. Let $\xi, \eta \in R^+$, $\xi < \eta$, we have an obvious result

$$A(\xi, \eta) \leq \max\{\xi, \eta\}, \quad (5)$$

$$L_n^n(\xi, \eta) \leq \max\{\xi^n, \eta^n\}. \quad (6)$$

Proof. The assertion (5) follows from Lemma 3 applied to the quasi-convex function $f(x) = x$, where $x \in R$ and taking $\alpha = k = 1$. The second assertion (6) also follows from Lemma 3 applied to the quasi-convex function $f(x) = x^n$ and $\alpha = k = 1$. Note that $f(x) = x^n$ is quasi-convex for; $x \in R$ when $n \in \mathbb{Z}$ is even and $x \in R^+$ when $n \in \mathbb{Z}$ is odd. \square

PROPOSITION 2. Let $\xi, \eta \in R^+$, $\xi < \eta$, and $n \in \mathbb{Z} \setminus \{-1, 0\}$, we have

$$|A(\xi^n, \eta^n) - L_n^n(\xi, \eta)| \leq \frac{\eta - \xi}{4} \max\{|n\xi^{n-1}|, |n\eta^{n-1}|\}.$$

Proof. The assertions follow from Theorem 4, applied to function $f(x) = x^n$ $\alpha = k = 1$. Note that if $f(x) = x^n$, then $|f'(x)|$ is quasi-convex for; $x \in R$ with $n \in Z$ is 0, 2 or odd, and $x \in R^+$ with $n \in Z \setminus \{0, 2\}$ is even. \square

PROPOSITION 3. Let $\xi, \eta \in R^+$, $\xi < \eta$, and $n \in Z$, we have

$$|A(\xi, \eta) - L(\xi, \eta)| \leq \frac{\ln \eta - \ln \xi}{4} \max\{\xi, \eta\}.$$

Proof. The assertions follow from Theorem 4, applied to the function $f(x) = e^x$ and $\alpha = k = 1$. Note that if $f(x) = e^x$ then $|f'(x)|$ is convex for all $x \in R$. \square

PROPOSITION 4. Let $\xi, \eta \in R^+$, $\xi < \eta$, $q > 1$ and $n \in Z$, we have

$$|G(\xi, \eta) - I(\xi, \eta)| \leq e^{-\frac{\eta - \xi}{2(\rho+1)^{\frac{1}{p}}}} \left(\max\left\{ \left| \frac{1}{\xi} \right|^q, \left| \frac{1}{\eta} \right|^q \right\} \right)^{\frac{1}{q}}.$$

Proof. The assertions follow from Theorem 5, applied the function $f(x) = \ln x$ and $\alpha = k = 1$. Note that if $f(x) = \ln x$ then $|f'(x)|^q$ is quasi-convex for $x \in R \setminus \{0\}$. \square

PROPOSITION 5. Let $\xi, \eta \in R^+$, $\xi < \eta$, $q \geq 1$ and $n \in Z$, we have

$$|H^{-1}(\xi, \eta) - L(\xi, \eta)| \leq \frac{\eta - \xi}{4} \left(\max\left\{ \left| \frac{1}{\xi} \right|^q, \left| \frac{1}{\eta} \right|^q \right\} \right)^{\frac{1}{q}}.$$

Proof. The assertions follow from Theorem 6, applied to the function $f(x) = \frac{1}{x}$ and $\alpha = k = 1$. Note that if $f(x) = \frac{1}{x}$ the function $|f'(x)|^q$ is convex for all $x \in R$. \square

4. Conclusion

In the present paper we have presented generalization of Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals of order α , given in [9], to the corresponding inequalities for k -Riemann-Liouville fractional integrals of order α .

REFERENCES

- [1] M. ALOMARI, M. DARUS AND S. S. DRAGOMIR, *Inequalities of Hermite-Hadamard's type for functions whose derivatives absolute values are quasi-convex*, RGMIA Res. Rep. Coll. **12**, Supplement, Article 14.
- [2] R. DIAZ AND E. PARIGUAN, *On hypergeometric functions and pochhammer k -symbol*, Divulgaciones Matemáticas **15**, 2 (2007), 179–192.
- [3] S. S. DRAGOMIR AND C. E. M. PEARCE, *Quasi-convex functions and Hadamard's inequality*, Bull. Austral. Math. Soc. **57** (1998), 377–385.
- [4] S. S. DRAGOMIR AND C. E. M. PEARCE, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria Universty, 2000.

- [5] R. GORENFLO AND F. MAINARDI, *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien, 1997, 223–276.
- [6] R. HUSSAIN, A. ALI, G. GULSHAN, A. LATIF AND M. MUDDASSAR, *Generalized co-ordinated integral inequalities for convex functions by way of k -fractional derivatives*, Miskolc Mathematical Notesa Publications of the university of Miskolc. (Submitted)
- [7] D. A. ION, *Some estimates on the Hermite-Hadamard inequality through quasi-convex functions*, Annals of University of Craiova, Math. Sci. Ser., **34** (2007), 82–87.
- [8] S. MUBEEN AND G. M. HABIBULLAH, *k -fractional integrals and applications*, Int. J. Contemp. Math. Sciences **7**, 2 (2012), 89–94.
- [9] E. SET AND B. ÇELİK, *Fractional Hermite-Hadamard Type Inequalities for Quasi-convex functions*, Ordu Univ. J. Sci. Tech. **6**, 1 (2016), 137–149.

(Received October 1, 2016)

R. Hussain
Department of Mathematics
Mirpur University of science and Technology (MUST)
Mirpur-10250 (AJK), Pakistan
e-mail: chairperson.maths@must.edu.pk

A. Ali
Department of Mathematics
Mirpur University of science and Technology (MUST)
Mirpur-10250 (AJK), Pakistan
e-mail: drali.math@must.edu.pk

A. Latif
Department of Mathematics
Mirpur University of science and Technology (MUST)
Mirpur-10250 (AJK), Pakistan
e-mail: asialatif.maths@must.edu.pk

G. Gulshan
Department of Mathematics
Mirpur University of science and Technology (MUST)
Mirpur-10250 (AJK), Pakistan
e-mail: ghazala.maths@must.edu.pk