

ON THE WEAK SOLUTION $u \in C_{1-\alpha}(I, E)$ OF A FRACTIONAL-ORDER WEIGHTED CAUCHY TYPE PROBLEM IN REFLEXIVE BANACH SPACES

A. M. A. EL-SAYED AND SH. A. ABD EL-SALAM

(Communicated by N. Vasylyeva)

Abstract. In this paper, we study the existence of a weak solution $u \in C_{1-\alpha}(I, E)$ of the nonlinear weighted Cauchy type problem of fractional-order.

1. Introduction

In this paper, we study the existence of solutions, in the Banach space $C_{1-\alpha}[I, E]$, for the nonlinear weighted Cauchy-type problem of the following type

$$\begin{cases} D^\alpha u(t) = f(t, u(t)), & t > 0, \quad \alpha \in (0, 1) \\ t^{1-\alpha} u(t)|_{t=0} = b, & b > 0. \end{cases} \quad (1)$$

This problem has been studied by many authors for example in ([4]), the author supposed that the function $f(t, u)$ is continuous on $R^+ \times R$, $|f(t, u)| \leq t^\mu e^{-\sigma t} \psi(t) |u|^m$, $\mu \geq 0$, $m > 1$, $\sigma > 0$, $\psi(t)$ is a continuous function on R^+ . Also; In ([2]–[3]) the author proved the existence of L_1 and L_p solution of the same problem respectively.

2. Preliminaries

Let $L_1(I)$ be the space of Lebesgue integrable functions on the interval $I = [0, 1]$. Unless otherwise stated, E is a reflexive Banach space with norm $\|\cdot\|$ and dual E^* . We will denote by E_w the space E endowed with the weak topology $\sigma(E, E^*)$ and denote by $C(I, E)$ the space of continuous functions defined on $I = [0, 1]$ with norm

$$\|u\|_C = \sup_{t \in [0, 1]} \|u(t)\|.$$

Also; define the space $C_{1-\alpha}(I, E)$ by

$$C_{1-\alpha}(I, E) = \{u : t^{1-\alpha} u(t) \text{ is continuous on } I = [0, 1]\},$$

Mathematics subject classification (2010): 34A12, 34A08, 35D30.

Keywords and phrases: Weighted Cauchy type problem, fractional-order, weak solution.

with norm

$$\|u\|_{C_{1-\alpha}} = \|t^{1-\alpha} u\|_C.$$

We recall that the fractional integral operator of order $\alpha > 0$ with left-hand point a is defined by (see [9], [14], [15] and [20])

$$I_a^\alpha u(t) = \int_a^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} u(s) ds.$$

DEFINITIONS. Let E be a Banach space and let $u : I \rightarrow E$. Then

- (1) $u(\cdot)$ is said to be weakly continuous (measurable) at $t_0 \in I$ if for every $\varphi \in E^*$ we have $\varphi(u(\cdot))$ continuous (measurable) at t_0 .
- (2) A function $h : E \rightarrow E$ is said to be weakly sequentially continuous if h takes weakly convergent sequences in E to weakly convergent sequences in E .

Note that:

- (1) If u is weakly continuous on I , then u is strongly measurable (see [7]), hence weakly measurable.
- (2) In reflexive Banach spaces weakly measurable functions are Pettis integrable (see [1], [7] and [13] for the definition) if and only if $\varphi(u(\cdot))$ is Lebesgue integrable on I for every $\varphi \in E^*$.

Now, we present some auxiliary results that will be needed in this paper. Firstly, we state O'Regan fixed point theorem ([12]).

THEOREM 2.1. *Let E be a Banach space with Q a nonempty, bounded, closed, convex, equicontinuous subset of $C[I, E]$. Suppose $T : Q \rightarrow Q$ is weakly sequentially continuous and assume $TQ(t)$ is weakly relatively compact in E for each $t \in I$, holds. Then the operator T has a fixed point in Q .*

The following theorems can be found in [5], [22] and [10] respectively:

THEOREM 2.2. (Dominated convergence theorem for Pettis integral) *Let $u : I \rightarrow E$. Suppose there is a sequence (u_n) of Pettis integrable functions from I into E such that $\lim_{n \rightarrow \infty} \varphi(u_n) = \varphi(u)$ a.e. for $\varphi \in E^*$. If there is a scalar function $\psi \in L_1(I)$ with $\|u_n(\cdot)\| < \psi(\cdot)$ a.e. for all n , then u is Pettis integrable and*

$$\int_J u_n(s) ds \rightarrow \int_J u(s) ds \text{ weakly } \forall t \in I.$$

THEOREM 2.3. *A subset of a reflexive Banach space is weakly compact if and only if it is closed in the weak topology and bounded in the norm topology.*

THEOREM 2.4. *Let Q be a weakly compact subset of $C[I, E]$. Then $Q(t)$ is weakly compact subset of E for each $t \in I$.*

Finally, we state some results which is an immediate consequence of the Hahn-Banach theorem.

THEOREM 2.5. *Let E be a normed space with $u_0 \neq 0$. then there exists a $\varphi \in E^*$ with $\|\varphi\| = 1$ and $\varphi(u_0) = \|u_0\|$.*

THEOREM 2.6. *If $u_0 \in E$ is such that $\varphi(u_0) = 0$ for every $\varphi \in E^*$, then $u_0 = 0$.*

Now consider the fractional-order integral equation

$$u(t) = b t^{\alpha-1} + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} f(s, u(s)) ds, \quad t \in [0, 1]. \quad (2)$$

In [12] the author studied the integral equation

$$y(t) = x_0 + \int_0^t f(s, y(s)) ds, \quad t \in [0, T], \quad x_0 \in E$$

where $E = (E, |\cdot|)$ is a real Banach space, under the assumptions that $f(t, \cdot)$ is weakly sequentially continuous for each $t \in [0, T]$ and $f(\cdot, y(\cdot))$ is Pettis integrable on $[0, T]$ for each continuous function $y: [0, T] \rightarrow E$ and $|f(t, y)| \leq h_r(t)$ for a.e. $t \in [0, T]$ and all $y \in E$ with $|y| \leq r, r > 0, h_r \in L_1[0, T]$.

Also, in [11] the author studied the Volterra-Hammerstein integral equation

$$y(t) = h(t) + \int_0^t k(t, s) f(s, y(s)) ds, \quad t \in [0, T], \quad T > 0,$$

under the assumptions that $f: [0, T] \times B \rightarrow B$ is weakly-weakly continuous and $h: [0, T] \rightarrow B$ is weakly continuous, where B is a reflexive Banach space.

Here we study the existence of weak solution of the fractional-order integral equation (2) such that the function $f: I \times B_r \rightarrow E$ satisfies the following conditions:

- (1) For each $t \in I, f_t = f(t, \cdot)$ is weakly sequentially continuous.
- (2) For each $u \in E_r, f(\cdot, u(\cdot))$ is weakly measurable on I .
- (3) for any $r > 0$, the weak closure of the range of $f(I \times B_r)$ is weakly compact in E (or equivalently; there exists an M_r such that $\|f(t, u)\| \leq M_r$ for all $(t, u) \in I \times B_r$).

EXAMPLE 2.1. Let T be the interval $[0, 1]$ and define $f: T \rightarrow L^\infty(T)$ by $f(t) = \chi_{[0, t]}$. This function is weakly measurable and for each $\phi \in L_\infty^*$, we have $\phi f \in L_1$ (each ϕf is a function of bounded variation). Thus, according to Lemma 3.2, $I^\alpha f$ exists. Also, the fractional order Pettis integral of f exists see [6, 16, 18].

DEFINITION 2.1. By a weak solution of (2) we mean a function $u \in C_{1-\alpha}[I, E]$ such that for all $\varphi \in E^*$

$$\varphi(u(t)) = b t^{\alpha-1} + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \varphi(f(s, u(s))) ds, \quad t \in [0, 1].$$

3. Fractional-order integrals in reflexive Banach spaces

Here, we define the fractional-order integral operator in reflexive Banach spaces. Definition given below is an extension of such a notion for real-valued functions.

DEFINITION 3.1. Let $u : I \rightarrow E$ be a weakly measurable function, such that $\varphi(u(\cdot)) \in L_1(I)$, and let $\alpha > 0$. Then the fractional (arbitrary) order Pettis integral (shortly FPI) $I^\alpha u(t)$ is defined by

$$I^\alpha u(t) = \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} u(s) ds.$$

In the above definition the sign " \int " denotes the Pettis integral.

LEMMA 3.1. [16] Let $u : I \rightarrow E$ be a weakly measurable function, such that $\varphi(u(\cdot)) \in L_1(I)$, and let $\alpha > 0$. The fractional (arbitrary) order Pettis integral

$$I^\alpha u(t) = \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} u(s) ds$$

exists for almost every $t \in I$ as a function from I into E and $\varphi(I^\alpha u(t)) = I^\alpha \varphi(u(t))$.

LEMMA 3.2. [17] Let $u : I \rightarrow E$ be weakly continuous function on $[0, 1]$. Then, FPI of u exists for almost every $t \in [0, 1]$ as a weakly continuous function from $[0, 1]$ to E . Moreover;

$$\varphi(I^\alpha u(t)) = I^\alpha \varphi(u(t)), \text{ for all } \varphi \in E^*.$$

DEFINITION 3.2. [13] Let $u : I \rightarrow E$. We define the fractional-Pseudo derivative (shortly FPD) of u of order $\alpha \in (n-1, n)$, $n \in \mathbb{N}$ by

$$\frac{d^\alpha}{dt^\alpha} u(t) = D^n I^{n-\alpha} u(t).$$

In the above definition the sign " D " denotes the Pseudo differential operator.

LEMMA 3.3. [21] Let $u : [0, 1] \rightarrow E$ be weakly continuous function on $[0, 1]$ such that the real-valued function $I^{n-\alpha} \varphi u$ is n -times differentiable. Then, the FPD of u of order $\alpha \in (n-1, n)$ exists.

DEFINITION 3.3. A function $u : I \rightarrow E$ is called Pseudo solution of (1) if $u \in C_{1-\alpha}[I, E]$ has FPD of order $\alpha \in (0, 1)$, $t^{1-\alpha} u(t)|_{t=0} = b$, $b > 0$ and satisfies

$$\frac{d}{dt} \varphi(I^{1-\alpha} u(t)) = \varphi(f(t, u(t))), \text{ a.e. on } [0, 1], \text{ for each } \varphi \in E^*.$$

Now, for the properties of the integrals of fractional-orders in reflexive spaces we have the following lemma (see [16]):

LEMMA 3.4. Let $u : I \rightarrow E$ be weakly measurable and $\varphi(u(\cdot)) \in L_1(I)$. If $\alpha, \beta \in (0, 1)$, we have:

- (1) $I^\alpha I^\beta u(t) = I^{\alpha+\beta} u(t)$ for a.e. $t \in I$.
- (2) $\lim_{\alpha \rightarrow 1} I^\alpha u(t) = I^1 u(t)$ weakly uniformly on I if only these integrals exist on I .
- (3) $\lim_{\alpha \rightarrow 0} I^\alpha u(t) = u(t)$ weakly in E for a.e. $t \in I$.
- (4) If, for a fixed $t \in I$, $\varphi(u(t))$ is bounded for each $\varphi \in E^*$, then $\lim_{t \rightarrow 0} I^\alpha u(t) = 0$.

4. Main result

In this section we present our main result by proving the existence of solution of equation (2) in $C_{1-\alpha}[I, E]$.

Let E be a reflexive Banach space. And let

$$E_r = \left\{ u \in C_{1-\alpha}[I, E] : \|u\|_{C_{1-\alpha}} < b + \frac{M_r}{\Gamma(1+\alpha)} \right\}.$$

We will consider the set

$$B_r = \{u(t) \in E : u \in E_r, t \in I\}.$$

Now, we are in a position to formulate and prove our main result.

THEOREM 4.1. Let the assumptions (1)–(3) are satisfied, then equation (2) has at least one weak solution $u \in C_{1-\alpha}[I, E]$.

Proof. Let us define the operator T as

$$Tu(t) = b t^{\alpha-1} + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} f(s, u(s)) ds, \quad t \in [0, 1].$$

We will solve equation (2) by finding a fixed point of the operator T .

We will prove that

$$T : C_{1-\alpha}[I, E] \rightarrow C_{1-\alpha}[I, E].$$

First note that from assumption (2), we get that for each $u \in C_{1-\alpha}[I, E]$, $f(\cdot, u(\cdot))$ is weakly measurable on I . Since f has weakly compact range, then $\varphi(f(\cdot, u(\cdot)))$ is Lebesgue integrable on I for every $\varphi \in E^*$ and thus the operator T is well defined. Now, we show that if $u \in C_{1-\alpha}[I, E]$, then $Tu \in C_{1-\alpha}[I, E]$. Note that there exists $r > 0$ with $\|u\|_{C_{1-\alpha}} = \sup_{t \in I} \|t^{1-\alpha} u(t)\| < b + \frac{M_r}{\Gamma(1+\alpha)}$.

Now assumption (3) implies that

$$\|f(t, u(t))\| \leq M_r \quad \text{for } t \in [0, 1].$$

Let $t, \tau \in [0, 1]$ with $t > \tau$. Without loss of generality, assume $t^{1-\alpha} Tu(t) - \tau^{1-\alpha} Tu(\tau) \neq 0$. Then there exists (a consequence of Theorem 2.5) $\varphi \in E^*$ with $\|\varphi\| = 1$ and

$$\|t^{1-\alpha} Tu(t) - \tau^{1-\alpha} Tu(\tau)\| = \varphi(t^{1-\alpha} Tu(t) - \tau^{1-\alpha} Tu(\tau)).$$

Thus

$$\begin{aligned}
\|t^{1-\alpha}Tu(t) - \tau^{1-\alpha}Tu(\tau)\| &\leq \left| t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \varphi(f(s, u(s))) ds \right. \\
&\quad \left. - \tau^{1-\alpha} \int_0^\tau \frac{(\tau-s)^{\alpha-1}}{\Gamma(\alpha)} \varphi(f(s, u(s))) ds \right| \\
&\leq \left| \int_0^\tau \frac{t^{1-\alpha}(t-s)^{\alpha-1} - \tau^{1-\alpha}(\tau-s)^{\alpha-1}}{\Gamma(\alpha)} \varphi(f(s, u(s))) ds \right| \\
&\quad + \left| \int_\tau^t \frac{t^{1-\alpha}(t-s)^{\alpha-1}}{\Gamma(\alpha)} \varphi(f(s, u(s))) ds \right| \\
&\leq \frac{M_r}{\Gamma(\alpha)} \left(\int_0^\tau |t^{1-\alpha}(t-s)^{\alpha-1} - \tau^{1-\alpha}(\tau-s)^{\alpha-1}| ds \right. \\
&\quad \left. + \int_\tau^t |t^{1-\alpha}(t-s)^{\alpha-1}| ds \right) \\
&\leq \frac{M_r}{\Gamma(1+\alpha)} \left(2(t-\tau)^\alpha + |t-\tau| \right). \tag{3}
\end{aligned}$$

which proves that $Tu \in C_{1-\alpha}[I, E]$.

Now, let

$$Q = \left\{ u \in E_r : (\forall t, \tau \in I) \|t^{1-\alpha}u(t) - \tau^{1-\alpha}u(\tau)\| \leq \frac{M_r}{\Gamma(1+\alpha)} \left(2(t-\tau)^\alpha + |t-\tau| \right) \right\},$$

Note that Q is nonempty, closed, bounded, convex and equicontinuous subset of $C_{1-\alpha}[I, E]$. Now, we claim that $T : Q \rightarrow Q$ and is weakly sequentially continuous. If this is true then according to Theorem 2.3, TQ is bounded in $C_{1-\alpha}[I, E]$ (hence, Theorem 2.4, implies $TQ(t)$ is weakly relatively compact in E for each $t \in I$) and the result follows immediately from Theorem 2.1. It remains to prove our claim. First we show that T maps Q into Q . To see this, note that the inequality (3) shows that TQ is norm continuous. Now, take $u \in Q$; without loss of generality, we may assume that $t^{1-\alpha}I^\alpha f(t, u(t)) \neq 0$, then, by Theorem 2.5, there exists $\varphi \in E^*$ with $\|\varphi\| = 1$ and $\|t^{1-\alpha}I^\alpha f(t, u(t))\| = \varphi(t^{1-\alpha}I^\alpha f(t, u(t)))$. Thus

$$\|t^{1-\alpha}Tu(t)\| \leq b + \frac{M_r}{\Gamma(1+\alpha)}, \tag{4}$$

therefore

$$\|Tu\|_{C_{1-\alpha}} < b + \frac{M_r}{\Gamma(1+\alpha)}.$$

Thus $T : Q \rightarrow Q$. Finally, we will show that T is weakly sequentially continuous. To see this, let $\{u_n\}_{n=1}^\infty$ be a sequence in Q and let $u_n(t) \rightarrow u(t)$ in E_w for each $t \in [0, 1]$. Recall [10] that a sequence $\{u_n\}_{n=1}^\infty$ is weakly convergent in $C[I, E]$ if and only if it is weakly pointwise convergent in E . Fix $t \in I$. From the weak sequential continuity of $f(t, \cdot)$, the Lebesgue dominated convergence theorem (see assumption (3)) for the

Pettis integral [5] implies for each $\varphi \in E^*$ that $\varphi(Tu_n(t)) \rightarrow \varphi(Tu(t))$ a.e. on I , $Tu_n(t) \rightarrow Tu(t)$ in E_w . So $T : Q \rightarrow Q$ is weakly sequentially continuous. The proof is complete. \square

Now, we are looking for sufficient conditions to ensure the existence of Pseudo solution to the nonlinear weighted Cauchy-type problem (1).

Note that, the following theorem is a generalization of the results of §3.3 in [8]:

THEOREM 4.2. *If $f : I \times B_r \rightarrow E$ satisfies the assumptions of Theorem 4.1, then the nonlinear weighted Cauchy-type problem (1) has a fractional-Pseudo derivative (FPD) $u \in C_{1-\alpha}[I, E]$.*

Proof. Let us remark, that by assumptions (2), (3) the FPI of f of order $\alpha > 0$ exists and

$$\varphi(I^\alpha f(t, u(t))) = I^\alpha \varphi(f(t, u(t))), \text{ for all } \varphi \in E^*.$$

Let u be a solution of equation (2), then

$$u(t) = bt^{\alpha-1} + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} f(s, u(s)) ds, \quad t \in [0, 1].$$

It is clear that

$$t^{1-\alpha} u(t)|_{t=0} = b.$$

Furthermore, we have

$$u(t) = b t^{\alpha-1} + I^\alpha f(t, u(t)) \tag{5}$$

since $u \in C_{1-\alpha}[I, E]$, then $\varphi(I^{1-\alpha} u(t)) = I^{1-\alpha} \varphi(u(t))$, for all $\varphi \in E^*$ (see Lemma 3.2). From equation (5), we deduce that

$$\varphi(u(t)) = b t^{\alpha-1} + \varphi(I^\alpha f(t, u(t))). \tag{6}$$

Operating by $I^{1-\alpha}$ on both sides of the equation (6) and using the properties of fractional calculus in the space $L_1[0, 1]$ (see [19] and [20]) result in

$$I^{1-\alpha} \varphi(u(t)) = b_1 + I \varphi(f(t, u(t))).$$

Therefore,

$$\varphi(I^{1-\alpha} u(t)) = b_1 + I \varphi(f(t, u(t))).$$

Thus

$$\frac{d}{dt} \varphi(I^{1-\alpha} u(t)) = \varphi(f(t, u(t))) \text{ a.e. on } [0, 1].$$

That is u has the FPD of order $\alpha \in (0, 1)$ and u is a solution of the differential equation (1). Conversely, let $u(t)$ be a solution of (1), integrate both sides, then

$$I^{1-\alpha} \varphi(u(t)) - I^{1-\alpha} \varphi(u(t))|_{t=0} = I \varphi(f(t, u(t))),$$

operating by I^α on both sides of the last equation, then

$$I\varphi(u(t)) - I^\alpha C = I^{1+\alpha} \varphi(f(t, u(t))),$$

differentiate both sides, then

$$\varphi(u(t)) - C_1 t^{\alpha-1} = I^\alpha \varphi(f(t, u(t))),$$

from the initial condition, we find that $C_1 = b$, then we obtain (2), i.e. Problem (1) and equation (2) are equivalent to each other. \square

REFERENCES

- [1] J. DIESTEL AND J. J. UHL, JR., *Vector measures*, Math. Surveys Monogr. **15**, Amer. Math. Soc., Providence, R. I., (1977).
- [2] A. M. A. EL-SAYED AND SH. A. ABD EL-SALAM, *Weighted Cauchy-type problem of a functional differ-integral equation*, EJQTDE **30** (2007) 1–9.
- [3] A. M. A. EL-SAYED AND SH. A. ABD EL-SALAM, *L_p -solution of weighted Cauchy-type problem of a differ-integral functional equation*, Inter. J. of Nonlinear Sci. **5** (2008).
- [4] K. M. FURATI AND N. E. TATAR, *Power-type estimates for a nonlinear fractional differential equation*, Nonlinear analysis **62** (2005), 1025–1036.
- [5] R. F. GEITZ, *Pettis integration*, Proc. Amer. Math. Soc. **82** (1981), 81–86.
- [6] R. F. GEITZ, *Geometry and the Pettis integral*, Trans. Am. Math. Soc. **1982**, 169(2) 535–548.
- [7] E. HILLE AND R. S. PHILLIPS, *Functional Analysis and Semi-groups*, Amer. Math. Soc. colloq. Publ. **31**, Amer. Math. Soc., Providence, R. I., (1957).
- [8] A. A. KILBAS, H. M. SRIVASTAVA AND J. J. TRUJILLO, *Theory and applications of fractional differential equations*, North-Holland Mathematical Studies, Vol. 204, Elsevier (North-Holland) Science Publishers, Amsterdam, London and New York, (2006).
- [9] K. S. MILLER AND B. ROSS, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley, New York (1993).
- [10] A. R. MITCHELL AND CH. SMITH, *An existence theorem for weak solutions of differential equations in reflexive Banach spaces*, Nonlinear equations in abstract spaces. Proc. Int. Symp., Arlington, (1977), (1978), 387–403.
- [11] D. O'REGAN, *Integral equations in reflexive Banach spaces and weak topologies*, Amer. Math. Soc. vol. 124, no. 2 (1996), 607–614.
- [12] D. O'REGAN, *Weak solutions of ordinary differential equations in Banach spaces*, Applied Math. Letters **12** (1999), 101–105.
- [13] B. J. PETTIS, *On integration in vector spaces*, Trans. Amer. Math. Soc. **44** (1938), 277–304.
- [14] I. PODLUBNY AND A. M. A. EL-SAYED, *On two definitions of fractional calculus*, Preprint UEF 03-96 (ISBN 80-7099-252-2), Slovak Academy of Science-Institute of Experimental phys. (1996).
- [15] I. PODLUBNY, *Fractional Differential Equations*, Acad. press, San Diego-New York-London (1999).
- [16] H. A. H. SALEM AND A. M. A. EL-SAYED, *Weak solution for fractional order integral equations in reflexive Banach spaces*, Mathematica Slovaca, (2005), no. 2, 169–181.
- [17] H. A. H. SALEM, A. M. A. EL-SAYED AND O. L. MOUSTAFA, *A note on the fractional calculus in Banach spaces*, Studia Sci. Math. Hungar. **42** (2) (2005), 115–130.
- [18] H. A. H. SALEM, *On the nonlinear Hammerstien integral equations in Banach spaces and application to the boundary value problem of fractional order*, Mathematical and computer modelling **48** (2008), 1178–1190.
- [19] H. A. H. SALEM AND M. VĀTH, *An abstract Gronwall lemma and application to global existence results for functional differential and integral equations of fractional order*, J. Integral Equations Appl. **16** (2004), 411–439.

- [20] S. SAMKO, A. KILBAS AND O. L. MARICHEV, *Fractional Integrals and Derivatives*, Gordon and Breach Science Publisher, (1993).
- [21] G. F. STEFANSSON, *Pettis integrability*, Trans. Am. Math. Soc. 1993, 401–417.
- [22] A. SZEP, *Existence theorem for weak solutions of differential equations in Banach spaces*, Studia Sci. Math. Hungar. **6** (1971), 197–203.

(Received October 5, 2017)

A. M. A. El-Sayed
Faculty of Science
Alexandria University
Alexandria, Egypt

e-mail: amasayed@alexu.edu.eg

Sh. A. Abd El-Salam
Faculty of Science
Damanhour University
Damanhour, Egypt

e-mail: shrnahmed@sci.dmu.edu.eg