

ON HAYAJNEH AND KITTANEH'S CONJECTURE ON UNITARILY INVARIANT NORM

JUN-TONG LIU, QING-WEN WANG AND FANG-FANG SUN

(Communicated by Y. Seo)

Abstract. In this short note, we give an affirmative answer to a conjecture posed by Hayajneh and Kittaneh on a unitarily invariant norm inequality.

1. Introduction

Let \mathbb{M}_n be the set of $n \times n$ complex matrices. In [4], Bourin posed the following questions. Let $A, B \in \mathbb{M}_n$ be positive semidefinite and $p, q > 0$. Then

$$\|A^{p+q} + B^{p+q}\| \leq \|(A^p + B^p)(A^q + B^q)\| \quad (1)$$

holds or not for any unitarily invariant norm. Further whether

$$\|A^{p+q} + B^{p+q}\| \leq \|(A^p + B^p)^{1/2}(A^q + B^q)(A^p + B^p)^{1/2}\| \quad (2)$$

holds or not for any unitarily invariant norm.

Hayajneh and Kittaneh proved that the above inequalities (1) and (2) hold for the trace norm and the Hilbert-Schmidt norm. Meanwhile, they posed the following conjectures (see [5]).

CONJECTURE 1. Let $A_1, A_2, B_1, B_2 \in \mathbb{M}_n$ be positive semidefinite with $A_1A_2 = A_2A_1$ and $B_1B_2 = B_2B_1$. Then

$$\|A_1A_2 + B_1B_2\| \leq \|(A_1 + B_1)(A_2 + B_2)\|$$

holds for any unitarily invariant norm.

CONJECTURE 2. Let $A_1, A_2, B_1, B_2 \in \mathbb{M}_n$ be positive semidefinite with $A_1A_2 = A_2A_1$ and $B_1B_2 = B_2B_1$. Then

$$\|A_1A_2 + B_1B_2\| \leq \|(A_1 + B_1)^{1/2}(A_2 + B_2)(A_1 + B_1)^{1/2}\|$$

holds for any unitarily invariant norm.

Mathematics subject classification (2010): 15A60, 47A30.

Keywords and phrases: Inequality, positive semidefinite, unitarily invariant norm.

In [1], Audenaert proved that if $A_i, B_i \in \mathbb{M}_n, (i = 1, 2, \dots, k)$ be positive semidefinite such that A_i commutes with B_i for each i , then

$$\left\| \sum_{i=1}^k A_i B_i \right\| \leq \left\| \left(\sum_{i=1}^k A_i^{1/2} B_i^{1/2} \right)^2 \right\| \leq \left\| \left(\sum_{i=1}^k A_i \right) \left(\sum_{i=1}^k B_i \right) \right\| \tag{3}$$

holds for any unitarily invariant norm.

Particularly, this result confirms the Conjecture 1 and answers a question of Bourin. Recently Lin [7] gave another proof of the above inequalities (3).

Recall that a norm $\|\cdot\|$ on \mathbb{M}_n is unitarily invariant if $\|UAV\| = \|A\|$ for any $A \in \mathbb{M}_n$ and unitary matrices $U, V \in \mathbb{M}_n$. In the sequel, let $\|\cdot\|$ denote any unitarily invariant norm.

In this short paper, We shall give an affirmative answer to the above Conjecture 2 by proving the following theorem.

THEOREM 1. *For $i = 1, 2, \dots, k$, let $A_i, B_i \in \mathbb{M}_n$ be positive semidefinite such that A_i commutes with B_i for each i . Then*

$$\left\| \sum_{i=1}^k A_i B_i \right\| \leq \left\| \left(\sum_{i=1}^k A_i^{1/2} B_i^{1/2} \right)^2 \right\| \leq \left\| \left(\sum_{i=1}^k A_i \right)^{1/2} \left(\sum_{i=1}^k B_i \right) \left(\sum_{i=1}^k A_i \right)^{1/2} \right\|.$$

It is readily seen that the Conjecture 2 is a special case (i.e., $k = 2$) of Theorem 1. Thus, Theorem 1 confirms the Conjecture 2 and answers another question of Bourin.

COROLLARY 1. *Let $A, B \in \mathbb{M}_n$ be positive semidefinite and $p, q > 0$. Then*

$$\|A^{p+q} + B^{p+q}\| \leq \|(A^p + B^p)^{1/2} (A^q + B^q) (A^p + B^p)^{1/2}\|.$$

2. Proof of Theorem 1

For $X \in \mathbb{M}_n$, the conjugate transpose of X is denoted by X^* . Let the singular values of $X \in \mathbb{M}_n$ denote by $\sigma_1(X) \geq \sigma_2(X) \geq \dots \geq \sigma_n(X)$. The block matrix $\begin{bmatrix} A & X \\ X^* & B \end{bmatrix}$, where $A, B, X \in \mathbb{M}_n$, is positive partial transpose (i.e., PPT) if $\begin{bmatrix} A & X \\ X^* & B \end{bmatrix}$ and $\begin{bmatrix} A & X^* \\ X & B \end{bmatrix}$ are both positive semidefinite.

In order to prove Theorem 1, we need to show the following lemma, which is a variation of Theorem 4.1 in [6].

LEMMA 1. *Let $\begin{bmatrix} A & X \\ X^* & B \end{bmatrix}$ be PPT, where $A, B, X \in \mathbb{M}_n$. Then*

$$\|X^* X\| \leq \|A^{1/2} B A^{1/2}\|.$$

Proof. Since $\begin{bmatrix} A & X \\ X^* & B \end{bmatrix}$ is positive semidefinite, we have $X = A^{1/2}CB^{1/2}$ for some contraction $C \in \mathbb{M}_n$ (see [2, p. 13]). Similarly, $X^* = A^{1/2}DB^{1/2}$ for some contraction $D \in \mathbb{M}_n$. Hence, for $k = 1, \dots, n$,

$$\begin{aligned} \prod_{i=1}^k \sigma_i(X^*X) &= \prod_{i=1}^k \sigma_i(A^{1/2}DB^{1/2}A^{1/2}CB^{1/2}) \\ &\leq \prod_{i=1}^k \sigma_i(DB^{1/2}A^{1/2}CB^{1/2}A^{1/2}) \\ &\leq \prod_{i=1}^k \sigma_i(B^{1/2}A^{1/2})\sigma_i(B^{1/2}A^{1/2}) \\ &= \prod_{i=1}^k \sigma_i(A^{1/2}BA^{1/2}), \end{aligned}$$

where the first inequality is by the fact (see [3, p. 253]) that $\prod_{j=1}^k \sigma_j(XY) \leq \prod_{j=1}^k \sigma_j(YX)$, $k = 1, 2, \dots$, whenever XY is normal; the second one is due to the fact that $\sigma(AB) \prec_{\log} \sigma(A) \circ \sigma(B)$ (see [8, p. 353]). Therefore, we have $\sigma(X^*X) \prec_{w\log} \sigma(A^{1/2}BA^{1/2})$, then $\sigma(X^*X) \prec_w \sigma(A^{1/2}BA^{1/2})$. Consequently, the assertion follows. \square

Now, we are in a position to present the proof of Theorem 1.

Proof of Theorem 1. The first inequality in Theorem 1 follows from the result of Audenaert, we next prove the second inequality in Theorem 1. Since A_i commutes with B_i , then $A_i^{1/2}$ also commutes with $B_i^{1/2}$, hence, A_iB_i and $A_i^{1/2}B_i^{1/2}$ are both positive semidefinite. It is easy to see that

$$\begin{bmatrix} A_i & A_i^{1/2}B_i^{1/2} \\ A_i^{1/2}B_i^{1/2} & B_i \end{bmatrix}, \quad i = 1, \dots, k,$$

are positive semidefinite, by using the Schur complement, so they are all PPT. Consequently

$$\begin{bmatrix} \sum_{i=1}^k A_i & \sum_{i=1}^k A_i^{1/2}B_i^{1/2} \\ \sum_{i=1}^k A_i^{1/2}B_i^{1/2} & \sum_{i=1}^k B_i \end{bmatrix}$$

is also PPT. By Lemma 1, we get

$$\left\| \left(\sum_{i=1}^k A_i^{1/2}B_i^{1/2} \right)^2 \right\| \leq \left\| \left(\sum_{i=1}^k A_i \right)^{1/2} \left(\sum_{i=1}^k B_i \right) \left(\sum_{i=1}^k A_i \right)^{1/2} \right\|,$$

hence, the second inequality follows in Theorem 1. \square

Acknowledgements. Special thanks to the anonymous referee and the editor for their helpful comments and suggestions. This research was supported by National Natural Science Foundation Project of China (No. 11571220), An-hui Provincial Natural

Science Foundation (No. 1508085MA12), Key projects of University excellent talent support program, An-hui Provincial (No. gxyqZD2016188) and Natural Scientific Research Project of Fuyang Normal College (No. 2016FSKJ20).

REFERENCES

- [1] K. AUDENAERT, *A norm inequality for pairs of commuting positive semidefinite matrices*, Electron. J. Linear Algebra. **30** (2015), 80–84.
- [2] R. BHATIA, *Positive Definite Matrices*, Princeton University Press, Princeton, 2007.
- [3] R. BHATIA, *Matrix Analysis*, GTM 169, Springer-Verlag, New York, 1997.
- [4] J. C. BOURIN, *Matrix subadditivity inequalities and block matrices*, Internet. J. Math. **20** (2009), 679–691.
- [5] S. HAYAJNEH, F. KITTANEH, *Trace inequalities and a question of bourin*, Bull. Aust. Math. Soc. **88** (2013), 384–389.
- [6] M. LIN, *Inequalities related to 2×2 block PPT matrices*, Oper. Matrices. **9** (2015), 917–924.
- [7] M. LIN, *Remarks on two recent results of Audenaert*, Linear Algebra Appl. **489** (2016), 24–29.
- [8] F. ZHANG, *Matrix Theory, Basic results and techniques*, Second Edition, Springer, 2011.

(Received March 5, 2016)

Jun-Tong Liu
 Department of Mathematics
 Shanghai University
 Shanghai, 200444, China
 and
 School of Mathematics and Statistics
 Fuyang Normal College
 Fuyang, 236041, China
 e-mail: juntongliu82@163.com

Corresponding author:
Qing-Wen Wang
 Department of Mathematics
 Shanghai University
 Shanghai, 200444, China
 e-mail: wqw@t.shu.edu.cn, wqw369@yahoo.com

Fang-Fang Sun
 Department of Mathematics
 Shanghai University
 Shanghai, 200444, China
 e-mail: SFangHwu@139.com