

## BOUNDEDNESS OF SOME SUBLINEAR OPERATORS ON WEIGHTED VARIABLE HERZ–MORREY SPACES

LIWEI WANG AND LISHENG SHU

(Communicated by R. Oinarov)

*Abstract.* A new class of generalized Herz–Morrey spaces with weight and variable exponent is defined. In addition, the boundedness of some sublinear operators on such spaces is also considered. The approach is based on the Muckenhoupt theory with variable exponent and function decomposition.

### 1. Introduction

Following the fundamental work of Kováčik and Rákosník [15] in the early 1990s, function spaces with variable exponents were investigated by a significant number of authors, see [1, 2, 6, 7, 8, 16, 19, 24, 25, 26, 27, 28] and the references therein. The theory of these spaces had a remarkable development in part due to its useful applications. For instance, they appear in the modeling of electrorheological fluids, in the study of image processing and in PDE with non-standard growth conditions, for an overview we refer to [5, 9, 20].

On the other hand, there do exist many different properties between the variable function spaces and the classical cases. For instance, the variable exponent Lebesgue spaces  $L^{p(\cdot)}(\mathbb{R}^n)$  are not translation invariant. More precisely, if  $p(\cdot)$  is non-constant in  $\mathbb{R}^n$ , then there always exists  $f \in L^{p(\cdot)}(\mathbb{R}^n)$  and  $h \in \mathbb{R}^n$ , such that  $f(x+h)$  is not in  $L^{p(\cdot)}(\mathbb{R}^n)$ . In addition, if  $p(\cdot)$  is unbounded, then  $L^{p(\cdot)}(\mathbb{R}^n)$  is no longer separable and it can happen that  $L^\infty(E) \subset L^{p(\cdot)}(E)$  when the set  $E \subset \mathbb{R}^n$  has infinite measure, see [2] for further details. Therefore, apart from useful application considerations, the motivation to study such spaces have an intrinsic interest.

Herz spaces  $\dot{K}_{p(\cdot)}^{\alpha,q}(\mathbb{R}^n)$  and Herz–Morrey spaces  $M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\mathbb{R}^n)$  with variable exponent  $p(\cdot)$  were introduced first by Izuki [12, 13]. Very recently, the boundedness of some important operators in harmonic analysis, such as the Hardy–Littlewood maximal operator, Marcinkiewicz integrals and some sublinear operators and so on, were obtained on these spaces, see [14, 22, 23]. In 2016, Izuki and Noi [10, 11] defined new generalized Herz spaces having weight and variable exponent, namely, weighted Herz spaces with variable exponent. Under proper assumptions on each exponent and weight,

*Mathematics subject classification* (2010): 42B20, 42B25.

*Keywords and phrases:* Herz–Morrey space, variable exponent, Muckenhoupt weight, sublinear operator.

The research is supported by NSF of China (Grant No. 11471033).

they proved the boundedness of fractional integral operators and intrinsic square functions on those spaces. Inspired of their work [10, 13, 23], the aim of this paper is to define weighted Herz-Morrey spaces with variable exponent and study the boundedness of some sublinear operators on these spaces under certain weak size conditions, which are similar to those introduced by Soria and Weiss in [21].

We usually denote the ball with radius  $r$  and center  $x$  by  $B$ .  $f_B$  denotes the integral average of  $f$  on  $B$ , i.e.  $f_B = \frac{1}{|B|} \int_B f(x) dx$ .  $p'(\cdot)$  means the conjugate exponent defined by  $1/p(\cdot) + 1/p'(\cdot) = 1$ . The letter  $C$  stands for a positive constant, which may vary from line to line.

## 2. Preliminaries and lemmas

### 2.1. Lebesgue spaces with variable exponent

Let  $p(\cdot)$  be a measurable function with values in  $[1, \infty)$ . The set  $\mathcal{P}(\mathbb{R}^n)$  consists of all variable exponents  $p(\cdot)$  satisfying

$$1 < p_- \leq p(x) \leq p_+ < \infty,$$

where  $p_- := \text{ess inf}\{p(x) : x \in \mathbb{R}^n\}$ ,  $p_+ := \text{ess sup}\{p(x) : x \in \mathbb{R}^n\}$ . The variable exponent Lebesgue space  $L^{p(\cdot)}(\mathbb{R}^n)$  is the class of all measurable functions  $f$  on  $\mathbb{R}^n$  such that

$$I_{p(\cdot)}(f) := \int_{\mathbb{R}^n} |f(x)|^{p(x)} dx < \infty.$$

This set becomes a Banach space when equipped with the norm

$$\|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} = \inf\{\lambda > 0 : I_{p(\cdot)}(f/\lambda) \leq 1\}.$$

It is easy to see that the variable Lebesgue exponent norm has the following property

$$\|f^\sigma\|_{L^{p(\cdot)}(\mathbb{R}^n)} = \|f\|_{L^{\sigma p(\cdot)}(\mathbb{R}^n)}^\sigma, \quad \sigma \geq 1/p_-.$$

Given an open set  $\Omega \subset \mathbb{R}^n$ . For all compact subsets  $F \subset \Omega$ , the space  $L_{loc}^{p(\cdot)}(\Omega)$  is defined by

$$L_{loc}^{p(\cdot)}(\Omega) = \{f : f \in L^{p(\cdot)}(F)\}.$$

A measurable function  $g(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$  is called globally log-Hölder continuous if it satisfies

$$|g(x) - g(y)| \leq \frac{-C}{\log(|x - y|)}, \quad |x - y| \leq 1/2, \quad x, y \in \mathbb{R}^n, \quad (2.1)$$

$$|g(x) - g_\infty| \leq \frac{C}{\log(e + |x|)}, \quad x \in \mathbb{R}^n, \quad (2.2)$$

for some real constant  $g_\infty$ . The set of  $p(\cdot)$  satisfying (2.1) and (2.2) is denoted by  $LH(\mathbb{R}^n)$ . As is well-known, if  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$ , then the Hardy-Littlewood maximal operator  $M$ , which is defined by

$$Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy,$$

is bounded on  $L^{p(\cdot)}(\mathbb{R}^n)$ , see [3].

### 2.2. Weighted function spaces with variable exponent

Let  $\omega$  be a weight function on  $\mathbb{R}^n$ , that is,  $\omega$  is real-valued, non-negative and locally integrable. The weighted variable exponent Lebesgue space  $L^{p(\cdot)}(\omega)$  is defined by

$$L^{p(\cdot)}(\omega) := \{f : f\omega^{\frac{1}{p(\cdot)}} \in L^{p(\cdot)}(\mathbb{R}^n)\}.$$

The space  $L^{p(\cdot)}(\omega)$  is a Banach space equipped with the norm

$$\|f\|_{L^{p(\cdot)}(\omega)} = \|f\omega^{\frac{1}{p(\cdot)}}\|_{L^{p(\cdot)}}.$$

A weight is said to be a Muckenhoupt  $A_1$  weight if

$$M\omega(x) \leq C\omega(x) \quad a.e. x \in \mathbb{R}^n.$$

For  $1 < p < \infty$ , we say that  $\omega$  is an  $A_p$  weight if

$$\sup_B \left( \frac{1}{|B|} \int_B \omega(x) dx \right) \left( \frac{1}{|B|} \int_B \omega(x)^{1-p'} dx \right)^{p-1} < \infty.$$

The Muckenhoupt  $A_p$  class with constant exponent  $p \in (1, \infty)$  was recently generalized by Izuki and Noi [10, 11] as follows.

DEFINITION 2.1. Let  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ . A weight is said to be an  $A_{p(\cdot)}$  weight if

$$\sup_B \frac{1}{|B|} \|\omega^{1/p(\cdot)} \chi_B\|_{L^{p(\cdot)}} \|\omega^{-1/p(\cdot)} \chi_B\|_{L^{p'(\cdot)}} < \infty.$$

REMARK 2.1. If  $p(\cdot) \equiv p \in (1, \infty)$ , then we see immediately that the definition reduces to the classical Muckenhoupt  $A_p$  class. Cruz-Uribe et al. [4] showed that  $\omega \in A_{p(\cdot)}$  if and only if the Hardy-Littlewood maximal operator  $M$  is bounded on the space  $L^{p(\cdot)}(\omega)$ .

REMARK 2.2. Suppose that  $p(\cdot), q(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$  and  $p(\cdot) \leq q(\cdot)$ , then we have  $A_1 \subset A_{p(\cdot)} \subset A_{q(\cdot)}$ , see [11].

DEFINITION 2.2. Let  $0 < \beta < n$ , and  $p_1(\cdot), p_2(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  such that  $1/p_2(x) \equiv 1/p_1(x) - \beta/n$ . A weight  $\omega$  is said to be an  $A(p_1(\cdot), p_2(\cdot))$  weight if

$$\|\omega \chi_B\|_{L^{p_2(\cdot)}} \|\omega^{-1} \chi_B\|_{L^{p_1(\cdot)}} \leq C|B|^{1-\frac{\beta}{n}}.$$

REMARK 2.3. Let  $p_1(\cdot), p_2(\cdot)$  and  $\omega$  be as in Definition 2.2, we then have

$$\omega \in A(p_1(\cdot), p_2(\cdot)) \Leftrightarrow \omega^{p_2(\cdot)} \in A_{1+p_2(\cdot)/p_1(\cdot)}.$$

Let  $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ ,  $R_k = B_k \setminus B_{k-1}$  and  $\chi_k = \chi_{R_k}$  be the characteristic function of the set  $R_k$  for  $k \in \mathbb{Z}$ .

DEFINITION 2.3. Let  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ ,  $0 < q \leq \infty$  and  $\alpha \in \mathbb{R}$ . The homogeneous weighted Herz space  $\dot{K}_{p(\cdot)}^{\alpha, q}(\omega)$  is defined as the set of all  $f \in L_{loc}^{p(\cdot)}(\mathbb{R}^n \setminus \{0\}, \omega)$  such that

$$\|f\|_{\dot{K}_{p(\cdot)}^{\alpha, q}(\omega)} := \left( \sum_{k \in \mathbb{Z}} 2^{\alpha k q} \|f \chi_k\|_{L^{p(\cdot)}(\omega)}^q \right)^{1/q} < \infty,$$

with the usual modification when  $q = \infty$ .

DEFINITION 2.4. Let  $0 \leq \lambda < \infty$ ,  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ ,  $0 < q \leq \infty$  and  $\alpha \in \mathbb{R}$ . The homogeneous weighted Herz-Morrey space  $M\dot{K}_{q, p(\cdot)}^{\alpha, \lambda}(\omega)$  is defined as the set of all  $f \in L_{loc}^{p(\cdot)}(\mathbb{R}^n \setminus \{0\}, \omega)$  such that

$$\|f\|_{M\dot{K}_{q, p(\cdot)}^{\alpha, \lambda}(\omega)} := \sup_{L \in \mathbb{Z}} 2^{-L\lambda} \left( \sum_{k=-\infty}^L 2^{\alpha k q} \|f \chi_k\|_{L^{p(\cdot)}(\omega)}^q \right)^{1/q} < \infty,$$

with the usual modification when  $q = \infty$ .

REMARK 2.4. Clearly,  $M\dot{K}_{q, p(\cdot)}^{\alpha, 0}(\omega) = \dot{K}_{p(\cdot)}^{\alpha, q}(\omega)$ . In the case  $\omega \equiv 1$  and  $p(\cdot) \equiv p$ , then  $\dot{K}_{p(\cdot)}^{\alpha, q}(\omega) = \dot{K}_q^{\alpha, p}(\mathbb{R}^n)$  and  $M\dot{K}_{q, p(\cdot)}^{\alpha, \lambda}(\omega) = M\dot{K}_{p, q}^{\alpha, \lambda}(\mathbb{R}^n)$  are the classical Herz spaces and Herz-Morrey spaces in [18] and [17], respectively.

### 2.3. Key lemmas

In order to prove our main results, we need the following lemmas.

LEMMA 2.1. Let  $\mathcal{X}$  be a Banach function space on  $\mathbb{R}^n$ . If  $f \in \mathcal{X}$  and  $g \in \mathcal{X}'$ , then we have

$$\|fg\|_{L^1(\mathbb{R}^n)} \leq \|f\|_{\mathcal{X}} \|g\|_{\mathcal{X}'},$$

where  $\mathcal{X}'$  denotes the associated space of  $\mathcal{X}$ .

We remark that Lemma 2.1 is the well-known generalized Hölder inequality. As a direct application of Lemma 2.1, we obtain the following result:

LEMMA 2.2. If  $\mathcal{X}$  is a Banach function space on  $\mathbb{R}^n$ , then we have

$$|B| \|\chi_B\|_{\mathcal{X}}^{-1} \|\chi_B\|_{\mathcal{X}'}^{-1} \leq 1.$$

Lemmas 2.3 and 2.4 below have been proved by Izuki and Noi in [10, 11].

LEMMA 2.3. Let  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$  and  $\omega \in A_{rp(\cdot)}$ ,  $1/p_- < r < 1$ . Then there exists a constant  $0 < \delta < 1$  such that for all  $k, j \in \mathbb{Z}$ ,

$$\frac{\|\chi_{B_k}\|_{L^{p(\cdot)}(\omega)}}{\|\chi_{B_j}\|_{L^{p(\cdot)}(\omega)}} \leq C2^{(k-j)n\delta}, \quad k \leq j,$$

and

$$\frac{\|\chi_{B_k}\|_{L^{p(\cdot)}(\omega)}}{\|\chi_{B_j}\|_{L^{p(\cdot)}(\omega)}} \leq C2^{(k-j)nr}, \quad k > j.$$

LEMMA 2.4. Let  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$  and  $\omega^{p(\cdot)} \in A_1$ . Then there exists constants  $0 < \delta_1, \delta_2 < 1$  such that

$$\frac{\|\chi_S\|_{L^{p(\cdot)}(\omega^{p(\cdot)})}}{\|\chi_B\|_{L^{p(\cdot)}(\omega^{p(\cdot)})}} \leq C \left( \frac{|S|}{|B|} \right)^{\delta_1}, \quad \frac{\|\chi_S\|_{(L^{p(\cdot)}(\omega^{p(\cdot)}))'}}{\|\chi_B\|_{(L^{p(\cdot)}(\omega^{p(\cdot)}))'}} \leq C \left( \frac{|S|}{|B|} \right)^{\delta_2},$$

for all balls  $B$  and all measurable sets  $S \subset B$ .

REMARK 2.5. We remark that  $(L^{p(\cdot)}(\omega^{p(\cdot)}))' = L^{p'(\cdot)}(\omega^{-p'(\cdot)})$  provided that  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$  and  $\omega^{p(\cdot)} \in A_1$ . The proof can be found in [11].

### 3. Main results and their proofs

In this section, we prove the boundedness of some sublinear operators under certain weak size conditions on weighted Herz-Morrey spaces with variable exponent. We consider only  $0 < \lambda < \infty$ , the arguments are similar in the case  $\lambda = 0$ .

Our main results can be stated as follows.

THEOREM 3.1. Let  $0 \leq \lambda < \infty$ ,  $p(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$ ,  $0 < q < \infty$ ,  $1/p_- < r < 1$ ,  $\omega \in A_{rp(\cdot)}$  and  $-n\delta + \lambda < \alpha < n(1-r) + \lambda$ , where  $0 < \delta < 1$  is the constant appearing in Lemma 2.3. Suppose that a sublinear operator  $T$  satisfies the size conditions

$$|Tf(x)| \leq C\|f\|_{L^1(\mathbb{R}^n)}/|x|^n, \tag{3.1}$$

when  $\text{supp} f \subseteq R_k$  and  $|x| \geq 2^{k+1}$  with  $k \in \mathbb{Z}$ , and

$$|Tf(x)| \leq C2^{-kn}\|f\|_{L^1(\mathbb{R}^n)}, \tag{3.2}$$

when  $\text{supp} f \subseteq R_k$  and  $|x| \leq 2^{k-2}$  with  $k \in \mathbb{Z}$ . Then, if  $T$  is bounded on  $L^{p(\cdot)}(\omega)$ ,  $T$  is also bounded on  $M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)$ .

COROLLARY 3.1. Let  $\lambda, p(\cdot), q, r, \omega$  and  $\alpha$  be as in Theorem 3.1, if a sublinear operator  $T$  satisfies the size conditions

$$|Tf(x)| \leq C \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^n} dy, \quad x \notin \text{supp} f, \tag{3.3}$$

for any integrable and compactly supported functions  $f$  and  $T$  is bounded on  $L^{p(\cdot)}(\omega)$ , then  $T$  is bounded on  $M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)$ .

REMARK 3.1. We note that (3.3) is satisfied by many important operators in harmonic analysis, such as the Hardy-Littlewood maximal operator, Calderón-Zygmund operators and Bochner-Riesz means at the critical index and so on, see [21].

*Proof of Theorem 3.1.* Let  $f \in M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)$ . We decompose

$$f(x) = \sum_{j=-\infty}^{\infty} f(x)\chi_j(x) := \sum_{j=-\infty}^{\infty} f_j(x).$$

Then, we have

$$\begin{aligned} \|T(f)\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q &= \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \|T(f)\chi_k\|_{L^{p(\cdot)}(\omega)}^q \\ &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \left( \sum_{j=-\infty}^{k-2} \|T(f_j)\chi_k\|_{L^{p(\cdot)}(\omega)} \right)^q \\ &\quad + C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \left( \sum_{j=k-1}^{k+1} \|T(f_j)\chi_k\|_{L^{p(\cdot)}(\omega)} \right)^q \\ &\quad + C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \left( \sum_{j=k+2}^{\infty} \|T(f_j)\chi_k\|_{L^{p(\cdot)}(\omega)} \right)^q \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

We first estimate  $I_1$ . Noting that  $j \leq k-2$  and  $x \in R_k$ , by (3.1), Lemmas 2.1 and 2.2, we get

$$\begin{aligned} |T(f_j)(x)| &\leq C 2^{-kn} \|f_j\|_{L^1(\mathbb{R}^n)} \\ &\leq C 2^{-kn} \|f \omega^{\frac{1}{p(\cdot)}} \chi_j\|_{L^{p(\cdot)}(\mathbb{R}^n)} \|\omega^{-\frac{1}{p(\cdot)}} \chi_j\|_{L^{p'(\cdot)}(\mathbb{R}^n)} \\ &\leq C 2^{-kn} \|f \omega^{\frac{1}{p(\cdot)}} \chi_j\|_{L^{p(\cdot)}(\mathbb{R}^n)} \|\omega^{-\frac{1}{p(\cdot)}} \chi_{B_j}\|_{L^{p'(\cdot)}(\mathbb{R}^n)} \\ &\leq C 2^{-kn} \|f \chi_j\|_{L^{p(\cdot)}(\omega)} \|\chi_{B_j}\|_{(L^{p(\cdot)}(\omega))'} \\ &\leq C 2^{(j-k)n} \|f \chi_j\|_{L^{p(\cdot)}(\omega)} \|\chi_{B_j}\|_{L^{p(\cdot)}(\omega)}^{-1}, \end{aligned} \tag{3.4}$$

which combining with Lemma 2.3 yields

$$\begin{aligned} &\|T(f_j)\chi_k\|_{L^{p(\cdot)}(\omega)} \\ &\leq C 2^{(j-k)n} \|f \chi_j\|_{L^{p(\cdot)}(\omega)} \|\chi_{B_j}\|_{L^{p(\cdot)}(\omega)}^{-1} \|\chi_{B_k}\|_{L^{p(\cdot)}(\omega)} \\ &\leq C 2^{(j-k)n(1-r)} \|f \chi_j\|_{L^{p(\cdot)}(\omega)}. \end{aligned} \tag{3.5}$$

On the other hand, we see that

$$\begin{aligned}
 \|f\chi_j\|_{L^{p(\cdot)}(\omega)} &= 2^{-j\alpha} \left( 2^{j\alpha q} \|f\chi_j\|_{L^{p(\cdot)}(\omega)}^q \right)^{1/q} \\
 &\leq C 2^{-j\alpha} \left( \sum_{l=-\infty}^j 2^{l\alpha q} \|f\chi_l\|_{L^{p(\cdot)}(\omega)}^q \right)^{1/q} \\
 &= C 2^{j(\lambda-\alpha)} \left( 2^{-j\lambda} \left( \sum_{l=-\infty}^j 2^{l\alpha q} \|f\chi_l\|_{L^{p(\cdot)}(\omega)}^q \right)^{1/q} \right) \\
 &\leq C 2^{j(\lambda-\alpha)} \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}.
 \end{aligned} \tag{3.6}$$

Hence, in view of  $\alpha < n(1-r) + \lambda$ , combining (3.5) and (3.6), we have

$$\begin{aligned}
 I_1 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \left( \sum_{j=-\infty}^{k-2} 2^{(j-k)n(1-r)} \|f\chi_j\|_{L^{p(\cdot)}(\omega)} \right)^q \\
 &\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\lambda q k} \left( \sum_{j=-\infty}^{k-2} 2^{(j-k)(n(1-r)+\lambda-\alpha)} \right)^q \\
 &\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \left( \sum_{k=-\infty}^L 2^{\lambda q k} \right) \\
 &\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q.
 \end{aligned}$$

For  $I_2$ , using the boundedness of  $T$  on  $L^{p(\cdot)}(\omega)$ , we derive the estimate

$$I_2 \leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \|f\chi_k\|_{L^{p(\cdot)}(\omega)}^q \leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q.$$

We proceed now to estimate  $I_3$ . As argued before, we apply the size condition (3.2) and obtain

$$\begin{aligned}
 |T(f_j)(x)| &\leq C 2^{-jn} \|f_j\|_{L^1(\mathbb{R}^n)} \\
 &\leq C 2^{-jn} \|f\omega^{\frac{1}{p(\cdot)}}\chi_j\|_{L^{p(\cdot)}(\mathbb{R}^n)} \|\omega^{-\frac{1}{p(\cdot)}}\chi_j\|_{L^{p'(\cdot)}(\mathbb{R}^n)} \\
 &\leq C \|f\chi_j\|_{L^{p(\cdot)}(\omega)} \|\chi_{B_j}\|_{L^{p'(\cdot)}(\omega)}^{-1}.
 \end{aligned} \tag{3.7}$$

An application of Lemma 2.3 gives

$$\begin{aligned}
 &\|T(f_j)\chi_k\|_{L^{p(\cdot)}(\omega)} \\
 &\leq C \|f\chi_j\|_{L^{p(\cdot)}(\omega)} \|\chi_{B_j}\|_{L^{p(\cdot)}(\omega)}^{-1} \|\chi_{B_k}\|_{L^{p(\cdot)}(\omega)} \\
 &\leq C 2^{(k-j)n\delta} \|f\chi_j\|_{L^{p(\cdot)}(\omega)}.
 \end{aligned} \tag{3.8}$$

By (3.6) and (3.8), since  $\alpha > -n\delta + \lambda$ , we arrive at

$$\begin{aligned}
I_3 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\alpha q k} \left( \sum_{j=k+2}^{\infty} 2^{(k-j)n\delta} \|f\chi_j\|_{L^{p(\cdot)}(\omega)} \right)^q \\
&\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \sum_{k=-\infty}^L 2^{\lambda q k} \left( \sum_{j=k+2}^{\infty} 2^{(k-j)(n\delta + \alpha - \lambda)} \right)^q \\
&\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q \sup_{L \in \mathbb{Z}} 2^{-L\lambda q} \left( \sum_{k=-\infty}^L 2^{\lambda q k} \right) \\
&\leq C \|f\|_{M\dot{K}_{q,p(\cdot)}^{\alpha,\lambda}(\omega)}^q.
\end{aligned}$$

Consequently, the proof of Theorem 3.1 is complete.  $\square$

For the fractional singular integrals, we have a theorem similar to Theorem 3.1.

**THEOREM 3.2.** *Let  $0 \leq \lambda < \infty$ ,  $p_2(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap LH(\mathbb{R}^n)$ ,  $\omega^{p_2(\cdot)} \in A_1$ ,  $0 < q_1 \leq q_2 < \infty$ ,  $0 < \delta_1, \delta_2 < 1$  be the constants appearing in Lemma 2.4,  $0 < \gamma < \min\{n(\delta_1 + \delta_2), n\}$  and  $-n\delta_1 + \lambda < \alpha < n\delta_2 - \gamma + \lambda$ . Define  $p_1(\cdot)$  by  $1/p_2(\cdot) = 1/p_1(\cdot) - \gamma/n$ . Suppose that a sublinear operator  $T_\gamma$  satisfies*

$$|T_\gamma f(x)| \leq C \|f\|_{L^1(\mathbb{R}^n)} / |x|^{n-\gamma}, \quad (3.9)$$

when  $\text{supp} f \subseteq R_k$  and  $|x| \geq 2^{k+1}$  with  $k \in \mathbb{Z}$ , and

$$|T_\gamma f(x)| \leq C 2^{-k(n-\gamma)} \|f\|_{L^1(\mathbb{R}^n)}, \quad (3.10)$$

when  $\text{supp} f \subseteq R_k$  and  $|x| \leq 2^{k-2}$  with  $k \in \mathbb{Z}$ . Then, if  $T_\gamma$  is bounded from  $L^{p_1(\cdot)}(\omega^{p_1(\cdot)})$  to  $L^{p_2(\cdot)}(\omega^{p_2(\cdot)})$ ,  $T_\gamma$  is also bounded from  $M\dot{K}_{q_1,p_1(\cdot)}^{\alpha,\lambda}(\omega^{p_1(\cdot)})$  to  $M\dot{K}_{q_2,p_2(\cdot)}^{\alpha,\lambda}(\omega^{p_2(\cdot)})$ .

**COROLLARY 3.2.** *Let  $\lambda, p_1(\cdot), p_2(\cdot), q_1, q_2, \delta_1, \delta_2, \gamma, \omega$  and  $\alpha$  be as in Theorem 3.2, if a sublinear operator  $T_\gamma$  satisfies the size conditions*

$$|T_\gamma f(x)| \leq C \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^{n-\gamma}} dy, \quad x \notin \text{supp} f, \quad (3.11)$$

for any integrable and compactly supported functions  $f$  and  $T_\gamma$  is bounded from  $L^{p_1(\cdot)}(\omega^{p_1(\cdot)})$  to  $L^{p_2(\cdot)}(\omega^{p_2(\cdot)})$ , then  $T_\gamma$  is also bounded from  $M\dot{K}_{q_1,p_1(\cdot)}^{\alpha,\lambda}(\omega^{p_1(\cdot)})$  to  $M\dot{K}_{q_2,p_2(\cdot)}^{\alpha,\lambda}(\omega^{p_2(\cdot)})$ .

*Proof of Theorem 3.2.* Let  $f \in M\dot{K}_{q_1,p_1(\cdot)}^{\alpha,\lambda}(\omega^{p_1(\cdot)})$ . As in the proof of Theorem



3.1, we can write

$$\begin{aligned}
 \|T_\gamma(f)\|_{MK_{q_2, p_2(\cdot)}^{\alpha, \lambda}(\omega^{p_2(\cdot)})}^{q_1} &= \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \left( \sum_{k=-\infty}^L 2^{\alpha q_2 k} \|T_\gamma(f)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}^{q_2} \right)^{q_1/q_2} \\
 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \|T_\gamma(f)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}^{q_1} \\
 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=-\infty}^{k-2} \|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \right)^{q_1} \\
 &\quad + C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=k-1}^{k+1} \|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \right)^{q_1} \\
 &\quad + C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=k+2}^{\infty} \|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \right)^{q_1} \\
 &=: J_1 + J_2 + J_3.
 \end{aligned}$$

For  $J_1$ . Noting that  $j \leq k - 2$  and  $x \in R_k$ , by (3.9) and Lemma 2.1, we get

$$\begin{aligned}
 |T_\gamma(f_j)(x)| &\leq C|x|^{\gamma-n} \|f_j\|_{L^1(\mathbb{R}^n)} \\
 &\leq C2^{k(\gamma-n)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'}.
 \end{aligned} \tag{3.12}$$

Based on the fact that

$$\|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \leq C2^{j(n-\gamma)} \|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}^{-1},$$

see [10, p.11]. From (3.12), Lemma 2 and Lemma 4, it follows that

$$\begin{aligned}
 &\|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \\
 &\leq C2^{k(\gamma-n)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_k}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \\
 &\leq C2^{k\gamma} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_k}\|_{(L^{p_2(\cdot)}(\omega^{p_2(\cdot)}))'}^{-1} \\
 &\leq C2^{k\gamma} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_j}\|_{(L^{p_2(\cdot)}(\omega^{p_2(\cdot)}))'}^{-1} \frac{\|\chi_{B_j}\|_{(L^{p_2(\cdot)}(\omega^{p_2(\cdot)}))'}}{\|\chi_{B_k}\|_{(L^{p_2(\cdot)}(\omega^{p_2(\cdot)}))'}} \\
 &\leq C2^{(k-j)\gamma} 2^{(j-k)n\delta_2} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} 2^{jn} \|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}^{-1} \|\chi_{B_j}\|_{(L^{p_2(\cdot)}(\omega^{p_2(\cdot)}))'}^{-1} \\
 &\leq C2^{(k-j)(\gamma-n\delta_2)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}.
 \end{aligned} \tag{3.13}$$

A simple caculation shows that

$$\begin{aligned}
 \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} &= 2^{-j\alpha} \left( 2^{j\alpha q} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}^q \right)^{1/q} \\
 &\leq C2^{-j\alpha} \left( \sum_{l=-\infty}^j 2^{l\alpha q} \|f_l\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}^q \right)^{1/q} \\
 &= C2^{j(\lambda-\alpha)} \left( 2^{-j\lambda} \left( \sum_{l=-\infty}^j 2^{l\alpha q} \|f_l\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}^q \right)^{1/q} \right) \\
 &\leq C2^{j(\lambda-\alpha)} \|f\|_{MK_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}.
 \end{aligned} \tag{3.14}$$

Hence, in view of  $\alpha < n\delta_2 - \beta + \lambda$ , combining (3.13) and (3.14), we get

$$\begin{aligned}
J_1 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=-\infty}^{k-2} 2^{(k-j)(\gamma-n\delta_2)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \right)^{q_1} \\
&\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=-\infty}^{k-2} 2^{(k-j)(\gamma-n\delta_2)} 2^{j(\lambda-\alpha)} \right)^{q_1} \\
&\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\lambda q_1 k} \left( \sum_{j=-\infty}^{k-2} 2^{(k-j)(\gamma-n\delta_2+\alpha-\lambda)} \right)^{q_1} \\
&\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \left( \sum_{k=-\infty}^L 2^{\lambda q_1 k} \right) \\
&\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1}.
\end{aligned}$$

For  $J_2$ , by the boundedness of  $T_\gamma$  from  $L^{p_1(\cdot)}(\omega^{p_1(\cdot)})$  to  $L^{p_2(\cdot)}(\omega^{p_2(\cdot)})$ , one can easily obtain

$$J_2 \leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \|f\chi_k\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}^{q_1} \leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1}.$$

For  $J_3$ . Noting that  $j \geq k+2$  and  $x \in R_k$ , Lemma 2.1 yields

$$\begin{aligned}
|T_\gamma(f_j)(x)| &\leq C 2^{j(\gamma-n)} \|f_j\|_{L^1(\mathbb{R}^n)} \\
&\leq C 2^{j(\gamma-n)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'}.
\end{aligned} \tag{3.15}$$

Consequently, we have

$$\begin{aligned}
&\|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \\
&\leq C 2^{j(\gamma-n)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_j\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_k}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \\
&\leq C 2^{j(\gamma-n)} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_{B_j}\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \frac{\|\chi_{B_k}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}}{\|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}} \\
&\leq C 2^{j(\gamma-n)} 2^{(k-j)n\delta_1} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \|\chi_{B_j}\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})}.
\end{aligned} \tag{3.16}$$

Since  $\omega^{p_2(\cdot)} \in A_1 \subset A_{1+p_2(\cdot)/p_1'(\cdot)}$ , so  $\omega^{p_2(\cdot)} \in A(p_1(\cdot), p_2(\cdot))$ , then we obtain

$$\begin{aligned}
&\|\chi_{B_j}\|_{(L^{p_1(\cdot)}(\omega^{p_1(\cdot)}))'} \|\chi_{B_j}\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \\
&\leq C \|\omega^{-1}\chi_{B_j}\|_{L^{p_1'(\cdot)}} \|\omega\chi_{B_j}\|_{L^{p_2(\cdot)}} \\
&\leq C 2^{j(n-\gamma)}.
\end{aligned} \tag{3.17}$$

This combined with (3.16) shows that

$$\|T_\gamma(f_j)\chi_k\|_{L^{p_2(\cdot)}(\omega^{p_2(\cdot)})} \leq C 2^{(k-j)n\delta_1} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})}. \tag{3.18}$$

From (3.14) and (3.18), since  $\alpha > -n\delta_1 + \lambda$ , we get

$$\begin{aligned} J_3 &\leq C \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=k+2}^{\infty} 2^{(k-j)n\delta_1} \|f_j\|_{L^{p_1(\cdot)}(\omega^{p_1(\cdot)})} \right)^{q_1} \\ &\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\alpha q_1 k} \left( \sum_{j=k+2}^{\infty} 2^{(k-j)n\delta_1} 2^{j(\lambda-\alpha)} \right)^{q_1} \\ &\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{\lambda q_1 k} \left( \sum_{j=k+2}^{\infty} 2^{(k-j)(n\delta_1 + \alpha - \lambda)} \right)^{q_1} \\ &\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1} \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \left( \sum_{k=-\infty}^L 2^{\lambda q_1 k} \right) \\ &\leq C \|f\|_{M\dot{K}_{q_1, p_1(\cdot)}^{\alpha, \lambda}(\omega^{p_1(\cdot)})}^{q_1}. \end{aligned}$$

Thus, the proof of Theorem 3.2 is complete.  $\square$

*Acknowledgement.* The authors would like to express their deep gratitude to the referee for giving many valuable comments and suggestions.

REFERENCES

- [1] A. ALMEIDA AND D. DRIHEM, *Maximal, potential and singular type operators on Herz spaces with variable exponents*, J. Math. Anal. Appl., **394** (2012), 781–795.
- [2] D. CRUZ-URIBE AND A. FIORENZA, *Variable Lebesgue Spaces: Foundations and Harmonic Analysis, Applied and Numerical Harmonic Analysis*, Birkhäuser, Basel, 2013.
- [3] D. CRUZ-URIBE, A. FIORENZA AND C. NEUGEBAUER, *The maximal function on variable  $L^p$  spaces*, Ann. Acad. Sci. Fenn. Math., **28** (2003), 223–238.
- [4] D. CRUZ-URIBE, A. FIORENZA AND C. NEUGEBAUER, *Weighted norm inequalities for the maximal operator on variable Lebesgue spaces*, J. Math. Anal. Appl., **394** (2012), 744–760.
- [5] Y. CHEN, S. LEVINE AND R. RAO, *Variable exponent, linear growth functionals in image restoration*, SIAM J. Appl. Math., **66** (2006), 1383–1406.
- [6] D. CRUZ-URIBE, SFO, A. FIORENZA, J. MARTELL AND C. PÉREZ, *The boundedness of classical operators on variable  $L^p$  spaces*, Ann. Acad. Sci. Fenn. Math., **31** (2006), 239–264.
- [7] L. DIENING, P. HÄSTÖ AND S. ROUDENKO, *Function spaces of variable smoothness and integrability*, J. Funct. Anal., **256** (2009), 1731–1768.
- [8] X. FAN AND D. ZHAO, *On the spaces  $L^{p(x)}(\Omega)$  and  $W^{m, p(x)}(\Omega)$* , J. Math. Anal. Appl., **263** (2001), 424–446.
- [9] P. HARJULEHTO, P. HÄSTÖ, Ú. V. LÊ AND M. NUORTIO, *Overview of differential equations with non-standard growth*, Nonlinear Anal., **72** (2010), 4551–4574.
- [10] M. IZUKI AND T. NOI, *Boundedness of fractional integrals on weighted Herz spaces with variable exponent*, J. Inequal. Appl., **2016**, Article ID 199 (2016).
- [11] M. IZUKI AND T. NOI, *An intrinsic square function on weighted Herz spaces with variable exponent*, arXiv: 1606.01019.
- [12] M. IZUKI, *Herz and amalgam spaces with variable exponent, the Haar wavelets and greediness of the wavelet system*, East J. Approx., **15** (2009), 87–109.
- [13] M. IZUKI, *Fractional integrals on Herz-Morrey spaces with variable exponent*, Hiroshima Math. J., **40** (2010), 343–355.
- [14] M. IZUKI, *Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization*, Anal. Math., **36** (2010), 33–50.
- [15] O. KOVÁČIK AND J. RÁKOSNÍK, *On spaces  $L^{p(x)}$  and  $W^{k, p(x)}$* , Czech. Math. J., **41** (1991), 592–618.

- [16] V. KOKILASHVILI AND S. SAMKO, *Maximal and fractional operators in weighted  $L^{p(x)}$  spaces*, Rev. Mat. Iberoam., **20** (2004), 495–517.
- [17] S. LU AND L. XU, *Boundedness of rough singular intergral operators on the homogeneous Morrey-Herz spaces*, Hokkaido Math. J., **34** (2005), 299–314.
- [18] S. LU, D. YANG AND G. HU, *Herz type spaces and their applications*, Science Press, Beijing, 2008.
- [19] Y. LU AND Y. ZHU, *Boundedness of multilinear Calderón-Zygmund singular operators on Morrey-Herz spaces with variable exponents*, Acta Math. Sin. (Engl. Ser.), **30** (2014), 1180–1194.
- [20] M. RŮŽIČKA, *Electrorheological fluids: modeling and mathematical theory*, Springer-Verlag, Berlin, 2000.
- [21] F. SORIA AND G. WEISS, *A remark on singular integrals and power weights*, Indiana Univ. Math. J., **43** (1994), 187–204.
- [22] H. WANG, *Commutators of Marcinkiewicz integrals on Herz spaces with variable exponent*, Czech. Math. J., **66** (2016), 251–269.
- [23] J. WU, *Boundedness of some sublinear operators on Herz-Morrey spaces with variable exponent*, Georgian Math. J., **21** (2014), 101–111.
- [24] J. XU, *Variable Besov and Triebel-Lizorkin spaces*, Ann. Acad. Sci. Fenn. Math., **33** (2008), 511–522.
- [25] D. YANG, C. ZHUO AND E. NAKAI, *Characterizations of variable exponent Hardy spaces via Riesz transforms*, Rev. Mat. Complut., **29** (2016), 245–270.
- [26] D. YANG, C. ZHUO AND W. YUAN, *Triebel-Lizorkin type spaces with variable exponents*, Banach J. Math. Anal., **9** (2015), 146–202.
- [27] D. YANG, C. ZHUO AND W. YUAN, *Besov-type spaces with variable smoothness and integrability*, J. Funct. Anal., **269** (2015), 1840–1898.
- [28] P. ZHANG AND J. WU, *Commutators for the maximal function on Lebesgue spaces with variable exponent*, Math. Inequal. Appl., **17** (2014), 1375–1386.

(Received September 21, 2016)

Liwei Wang  
 School of Mathematics and Physics  
 Anhui Polytechnic University  
 Wuhu 241000, The People's Republic of China  
 e-mail: wangliwei@ahpu.edu.cn

Lisheng Shu  
 School of Mathematics and Computer Science  
 Anhui Normal University  
 Wuhu 241003, The People's Republic of China  
 e-mail: shulsh@mail.ahnu.edu.cn