

## NECESSARY AND SUFFICIENT CONDITIONS FOR THE VALIDITY OF HILBERT TYPE INTEGRAL INEQUALITIES WITH A CLASS OF QUASI-HOMOGENEOUS KERNELS AND ITS APPLICATION IN OPERATOR THEORY

YONG HONG, BING HE\* AND BICHENG YANG

(Communicated by M. Krnić)

*Abstract.* By using real analysis technique and the method of weight functions, the necessary and sufficient conditions for the validity of Hilbert type integral inequalities with a class of quasi-homogeneous kernels and the best constant factors are obtained, and its applications in operator theory are discussed.

### 1. Introduction and preliminary knowledge

Let  $r > 1$ , and  $\alpha$  be a constant. Set

$$L_{\alpha}^r(0, +\infty) = \left\{ f(x) \geq 0 : \|f\|_{r,\alpha} = \left( \int_0^{+\infty} x^{\alpha} f^r(x) dx \right)^{1/r} < +\infty \right\}.$$

If  $\frac{1}{p} + \frac{1}{q} = 1$  ( $p > 1$ ),  $M$  is a nonnegative constant,  $K(x, y) \geq 0$ ,  $f(x) \in L_{\alpha}^p(0, +\infty)$ ,  $g(y) \in L_{\beta}^q(0, +\infty)$ , then the Hilbert type integral inequality is of the form

$$\int_0^{+\infty} \int_0^{+\infty} K(x, y) f(x) g(y) dx dy \leq M \|f\|_{p,\alpha} \|g\|_{q,\beta}, \quad (1)$$

Define the singular integral operator  $T$  by

$$T(f)(y) = \int_0^{+\infty} K(x, y) f(x) dx, f(x) \in L_{\alpha}^p(0, +\infty). \quad (2)$$

Then (1) can be equivalently written as

$$\int_0^{+\infty} g(y) T(f)(y) dy \leq M \|f\|_{p,\alpha} \|g\|_{q,\beta}. \quad (3)$$

*Mathematics subject classification* (2010): 26D15, 47A07.

*Keywords and phrases:* Quasi-homogeneous kernel, Hilbert's integral inequality, necessary and sufficient conditions, best constant factor, bounded operator, operator norm.

This work is supported by the NNSF of China (Nos. 61370186, 61640222) and project for Distinctive Innovation of Ordinary University of Guangdong Province (Nos. 2015KTSCX097, 2016KTSCX093, 2016KTSCX094).

\* The corresponding author.

An operator  $T$  is called the bounded operator from  $L^p_\alpha(0, +\infty)$  to  $L^p_\gamma(0, +\infty)$ , if  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$ . At this point, the operator norm of  $T$  is defined by

$$\|T\| = \inf_{f \in L^p_\alpha(0, +\infty)} \frac{\|T(f)\|_{p,\gamma}}{\|f\|_{p,\alpha}}.$$

In particular,  $T$  is called a bounded operator in  $L^p_\alpha(0, +\infty)$ , if the bounded operator  $T$  is from  $L^p_\alpha(0, +\infty)$  to itself. For the sake of convenience, denote by  $\|f\|_{p,0} = \|f\|_p$ ,  $L^p_0(0, +\infty) = L^p(0, +\infty)$ .

One can know from (3) that, the boundedness of the operator is closely related to the Hilbert type inequality. Although the study of Hilbert type inequality has made many achievements (cf. [1]–[16]), but most of them are discussed for a specific integral kernel. In this paper, we will study the necessary and sufficient conditions for the validity of Hilbert type inequality with a class of quasi-homogeneous kernels, and discuss its applications in operator theory.

DEFINITION 1. A function  $K(x, y)$  is said to be quasi-homogeneous of order  $(\lambda_1, \lambda_2)$  if for  $t > 0$ ,

$$K(tx, y) = t^{\lambda_1} K(x, t^{-\frac{\lambda_1}{\lambda_2}} y), \quad K(x, ty) = t^{\lambda_2} K(t^{-\frac{\lambda_2}{\lambda_1}} x, y),$$

where  $\lambda_1$  and  $\lambda_2$  are nonzero constants. When  $\lambda_1 = \lambda_2 = \lambda$ ,  $K(x, y)$  becomes a homogeneous function of order  $\lambda$ .

LEMMA 1. Let  $T$  be as in (2). Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $\gamma = \beta(1 - p)$ . Then Hilbert type inequality (1) is equivalent to  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$ .

*Proof. Necessity.* If (1) holds, set

$$g(y) = y^{\beta(1-p)} (T(f)(y))^{p-1} = y^\gamma (T(f)(y))^{p-1},$$

then

$$\begin{aligned} \|T(f)\|_{p,\gamma}^p &= \int_0^{+\infty} y^\gamma (T(f)(y))^p dy = \int_0^{+\infty} \int_0^{+\infty} K(x, y) f(x) g(y) dx dy \\ &\leq M\|f\|_{p,\alpha} \|g\|_{q,\beta} = M\|f\|_{p,\alpha} \left( \int_0^{+\infty} y^\beta g^q(y) dy \right)^{1/q} \\ &= M\|f\|_{p,\alpha} \left( \int_0^{+\infty} y^\gamma (T(f)(y))^p dy \right)^{1/q} = M\|f\|_{p,\alpha} \|T(f)\|_{p,\gamma}^{p/q}. \end{aligned}$$

If  $\|T(f)\|_{p,\gamma} = 0$ , then inequality  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$  is naturally valid; if  $\|T(f)\|_{p,\gamma} = \infty$ , then inequality  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$  is impossible. Thus, in view of the above results, we have  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$ .

*Sufficiency.* If  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$ , then it is not difficult to derive (1), whence (1) and the inequality  $\|T(f)\|_{p,\gamma} \leq M\|f\|_{p,\alpha}$  are equivalent.  $\square$

LEMMA 2. Assume that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1\lambda_2 > 0$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $\frac{\lambda_2\alpha - \lambda_1}{p} + \frac{\lambda_1\beta - \lambda_2}{q} = \lambda_1\lambda_2$ ,  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(\lambda_1, \lambda_2)$ , set

$$W_1 = \int_0^{+\infty} u^{-\frac{\beta+1}{q}} K(1, u) du, W_2 = \int_0^{+\infty} u^{-\frac{\alpha+1}{p}} K(u, 1) du,$$

then  $\lambda_1 W_2 = \lambda_2 W_1$ , and

$$\omega_1(x) = \int_0^{+\infty} y^{-\frac{\beta+1}{q}} K(x, y) dy = x^{\lambda_1 - \frac{\lambda_1}{\lambda_2} \left(\frac{\beta+1}{q} - 1\right)} W_1,$$

$$\omega_2(y) = \int_0^{+\infty} x^{-\frac{\alpha+1}{p}} K(x, y) dx = y^{\lambda_2 - \frac{\lambda_2}{\lambda_1} \left(\frac{\alpha+1}{p} - 1\right)} W_2.$$

*Proof.* Since  $K(x, y)$  is a quasi-homogeneous function of order  $(\lambda_1, \lambda_2)$ , and  $\frac{\lambda_2\alpha - \lambda_1}{p} + \frac{\lambda_1\beta - \lambda_2}{q} = \lambda_1\lambda_2 > 0$ , we have

$$\begin{aligned} -\frac{\lambda_1}{\lambda_2} \left( \lambda_2 - \frac{\beta + 1}{q} \right) &= -\frac{1}{\lambda_2} \left( \lambda_1 \lambda_2 - \frac{\lambda_1 \beta - \lambda_2}{q} - \frac{\lambda_1 + \lambda_2}{q} \right) \\ &= -\frac{1}{\lambda_2} \left( \lambda_1 \lambda_2 + \frac{\lambda_2 \alpha - \lambda_1}{p} - \lambda_1 \lambda_2 - \frac{\lambda_1 + \lambda_2}{q} \right) \\ &= -\frac{\alpha}{p} + \frac{\lambda_1}{p \lambda_2} + \frac{\lambda_1 + \lambda_2}{q \lambda_2} = -\frac{\alpha + 1}{p} + 1 + \frac{\lambda_1}{\lambda_2}, \end{aligned}$$

and

$$\begin{aligned} W_1 &= \int_0^{+\infty} u^{\lambda_2 - \frac{\beta+1}{q}} K(u^{-\frac{\lambda_2}{\lambda_1}}, 1) du \\ &= \frac{\lambda_1}{\lambda_2} \int_0^{+\infty} t^{-\frac{\lambda_1}{\lambda_2} \left(\lambda_2 - \frac{\beta+1}{q}\right) - \frac{\lambda_1}{\lambda_2} - 1} K(t, 1) dt \\ &= \frac{\lambda_1}{\lambda_2} \int_0^{+\infty} t^{-\frac{\alpha+1}{p}} K(t, 1) dt = \frac{\lambda_1}{\lambda_2} W_2, \end{aligned}$$

it follows that  $\lambda_2 W_1 = \lambda_1 W_2$ . Substituting  $x^{-\frac{\lambda_1}{\lambda_2}} y = u$ , then

$$\begin{aligned} \omega_1(x) &= \int_0^{+\infty} x^{\lambda_1} y^{-\frac{\beta+1}{q}} K(1, x^{-\frac{\lambda_1}{\lambda_2}} y) dy \\ &= x^{\lambda_1 - \frac{\lambda_1}{\lambda_2} \left(\frac{\beta+1}{q} - 1\right)} \int_0^{+\infty} u^{-\frac{\beta+1}{q}} K(1, u) du \\ &= x^{\lambda_1 - \frac{\lambda_1}{\lambda_2} \left(\frac{\beta+1}{q} - 1\right)} W_1. \end{aligned}$$

Similarly, we also have  $\omega_2(y) = y^{\lambda_2 - \frac{\lambda_2}{\lambda_1} \left(\frac{\alpha+1}{p} - 1\right)} W_2$ .  $\square$

2. Main results

**THEOREM 1.** *Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 \lambda_2 > 0$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(\lambda_1, \lambda_2)$ ,  $K(1, u) (K(u, 1)) > 0$  a.e. in  $(0, 1) ((1, \infty))$ , and*

$$W_1 = \int_0^{+\infty} u^{-\frac{\beta+1}{q}} K(1, u) du, W_2 = \int_0^{+\infty} u^{-\frac{\alpha+1}{p}} K(u, 1) du$$

are both convergent. Then

(i) *there exists a constant  $M$ , for any  $f \in L^p_\alpha(0, +\infty)$ ,  $g \in L^q_\beta(0, +\infty)$ , the necessary and sufficient conditions for Hilbert type inequality*

$$\int_0^{+\infty} \int_0^{+\infty} K(x, y) f(x) g(y) dx dy \leq M \|f\|_{p, \alpha} \|g\|_{q, \beta} \tag{4}$$

to be true is that  $\frac{\lambda_2 \alpha - \lambda_1}{p} + \frac{\lambda_1 \beta - \lambda_2}{q} = \lambda_1 \lambda_2$ ;

(ii) *if (4) holds, then its best possible constant factor is  $M = \frac{W}{|\lambda_1|^{1/q} |\lambda_2|^{1/p}}$ , where  $W = |\lambda_1| W_2 = |\lambda_2| W_1$ .*

*Proof.* (i) *Necessity.* Assume that there exists a constant  $M$ , such that (4) holds. Set  $c = \frac{\lambda_2 \alpha - \lambda_1}{p} + \frac{\lambda_1 \beta - \lambda_2}{q} - \lambda_1 \lambda_2$ . First consider the case of  $\lambda_1 > 0, \lambda_2 > 0$ . If  $c > 0$ , for sufficiently small  $\varepsilon > 0$  ( $\varepsilon < \frac{c}{\lambda_1 \lambda_2}$ ), take

$$f(x) = \begin{cases} x^{(-\alpha-1+\lambda_1\varepsilon)/p}, & 0 < x \leq 1, \\ 0, & x > 1, \end{cases}$$

$$g(y) = \begin{cases} y^{(-\beta-1+\lambda_2\varepsilon)/q}, & 0 < y \leq 1, \\ 0, & y > 1. \end{cases}$$

Then

$$\|f\|_{p, \alpha} \|g\|_{q, \beta} = \left( \int_0^1 x^{-1+\lambda_1\varepsilon} dx \right)^{1/p} \left( \int_0^1 y^{-1+\lambda_2\varepsilon} dy \right)^{1/q} = \frac{1}{\varepsilon \lambda_1^{1/p} \lambda_2^{1/q}}, \tag{5}$$

$$\begin{aligned} & \int_0^{+\infty} \int_0^{+\infty} K(x, y) f(x) g(y) dx dy \\ &= \int_0^1 x^{(-\alpha-1+\lambda_1\varepsilon)/p} \left( \int_0^1 K(x, y) y^{(-\beta-1+\lambda_2\varepsilon)/q} dy \right) dx \\ &= \int_0^1 x^{\lambda_1 + (-\alpha-1+\lambda_1\varepsilon)/p} \left( \int_0^1 K(1, x^{-\lambda_1/\lambda_2} y) y^{(-\beta-1+\lambda_2\varepsilon)/q} dy \right) dx \\ &= \int_0^1 x^{-1+\lambda_1\varepsilon - \frac{c}{\lambda_2}} \left( \int_0^{x^{-\lambda_1/\lambda_2}} K(1, t) t^{(-\beta-1+\lambda_2\varepsilon)/q} dt \right) dx \\ &\geq \int_0^1 x^{-1+\lambda_1\varepsilon - \frac{c}{\lambda_2}} dx \int_0^1 K(1, t) t^{(-\beta-1+\lambda_2\varepsilon)/q} dt. \end{aligned} \tag{6}$$

It follows from (4), (5) and (6) that

$$\int_0^1 x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} dx \int_0^1 K(1,t)t^{(-\beta-1+\lambda_2\varepsilon)/q} dt \leq \frac{M}{\varepsilon\lambda_1^{1/p}\lambda_2^{1/q}}. \tag{7}$$

Since  $c > 0$ ,  $\lambda_2 > 0$ ,  $\lambda_1\varepsilon - \frac{c}{\lambda_2} < 0$  and  $\int_0^1 x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} dx$  diverges to  $+\infty$ . Whence it is a contradiction of (7). In other words, it is not valid for  $c > 0$ .

If  $c < 0$ , for sufficiently small  $\varepsilon > 0$  ( $\varepsilon < \frac{-c}{\lambda_1\lambda_2}$ ), take

$$f(x) = \begin{cases} x^{(-\alpha-1-\lambda_1\varepsilon)/p}, & x \geq 1, \\ 0, & 0 < x < 1, \end{cases}$$

$$g(y) = \begin{cases} y^{(-\beta-1-\lambda_2\varepsilon)/q}, & y \geq 1, \\ 0, & 0 < y < 1. \end{cases}$$

We get

$$\|f\|_{p,\alpha} \|g\|_{q,\beta} = \left( \int_1^{+\infty} x^{-1-\lambda_1\varepsilon} dx \right)^{1/p} \left( \int_1^{+\infty} y^{-1-\lambda_2\varepsilon} dy \right)^{1/q} = \frac{1}{\varepsilon\lambda_1^{1/p}\lambda_2^{1/q}},$$

$$\begin{aligned} & \int_0^{+\infty} \int_0^{+\infty} K(x,y)f(x)g(y) dx dy \\ &= \int_1^{+\infty} y^{(-\beta-1-\lambda_2\varepsilon)/q} \left( \int_1^{+\infty} K(x,y)x^{(-\alpha-1-\lambda_1\varepsilon)/p} dx \right) dy \\ &= \int_1^{+\infty} y^{\lambda_2+(-\beta-1-\lambda_2\varepsilon)/q} \left( \int_1^{+\infty} K(xy^{-\lambda_2/\lambda_1}, 1)x^{(-\alpha-1-\lambda_1\varepsilon)/p} dx \right) dy \\ &= \int_1^{+\infty} y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} \left( \int_{y^{-\lambda_2/\lambda_1}}^{+\infty} K(t, 1)t^{(-\alpha-1-\lambda_1\varepsilon)/p} dt \right) dy \\ &\geq \int_1^{+\infty} y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} dy \int_1^{+\infty} K(t, 1)t^{(-\alpha-1-\lambda_1\varepsilon)/p} dt. \end{aligned}$$

Therefore

$$\int_1^{+\infty} y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} dy \int_1^{+\infty} K(t, 1)t^{(-\alpha-1-\lambda_1\varepsilon)/p} dt \leq \frac{M}{\varepsilon\lambda_1^{1/p}\lambda_2^{1/q}}.$$

Since  $c < 0$ ,  $\lambda_1 > 0$ ,  $-\lambda_2\varepsilon - \frac{c}{\lambda_1} > 0$  and  $\int_1^{+\infty} y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} dy$  diverges to  $+\infty$ , which contradicts the above inequality. Hence it does not hold for  $c < 0$ .

To sum up, when  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ , we have  $c = 0$ , that is  $\frac{\lambda_2\alpha-\lambda_1}{p} + \frac{\lambda_1\beta-\lambda_2}{q} = \lambda_1\lambda_2$ .

Now let us consider the case of  $\lambda_1 < 0$ ,  $\lambda_2 < 0$ . If  $c > 0$ , for sufficiently small  $\varepsilon > 0$  ( $\varepsilon < \frac{c}{\lambda_1\lambda_2}$ ), take

$$f(x) = \begin{cases} x^{(-\alpha-1+\lambda_1\varepsilon)/p}, & x \geq 1, \\ 0, & 0 < x < 1, \end{cases}$$

$$g(y) = \begin{cases} y^{(-\beta-1+\lambda_2\varepsilon)/q}, & y \geq 1, \\ 0, & 0 < y < 1. \end{cases}$$

Then

$$\begin{aligned} \|f\|_{p,\alpha} \|g\|_{q,\beta} &= \left( \int_1^{+\infty} x^{-1+\lambda_1\varepsilon} dx \right)^{1/p} \left( \int_1^{+\infty} y^{-1+\lambda_2\varepsilon} dy \right)^{1/q} \\ &= \frac{1}{\varepsilon(-\lambda_1)^{1/p}(-\lambda_2)^{1/q}}, \\ &= \int_0^{+\infty} \int_0^{+\infty} K(x,y) f(x) g(y) dx dy \\ &= \int_1^{+\infty} x^{\lambda_1+(-\alpha-1+\lambda_1\varepsilon)/p} \left( \int_1^{+\infty} K(1,x^{-\lambda_1/\lambda_2}y)^{(-\beta-1+\lambda_2\varepsilon)/q} dy \right) dx \\ &= \int_1^{+\infty} x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} \left( \int_{x^{-\lambda_1/\lambda_2}}^{+\infty} K(1,t)t^{(-\beta-1+\lambda_2\varepsilon)/q} dt \right) dx \\ &\geq \int_1^{+\infty} x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} dx \int_1^{+\infty} K(1,t)t^{(-\beta-1+\lambda_2\varepsilon)/q} dt. \end{aligned}$$

Therefore

$$\int_1^{+\infty} x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} dx \int_1^{+\infty} K(1,t)t^{(-\beta-1+\lambda_2\varepsilon)/q} dt \leq \frac{M}{\varepsilon(-\lambda_1)^{1/p}(-\lambda_2)^{1/q}}.$$

Since  $c > 0$ ,  $\lambda_2 < 0$ ,  $\lambda_1\varepsilon - \frac{c}{\lambda_2} > 0$  and  $\int_1^{+\infty} x^{-1+\lambda_1\varepsilon-\frac{c}{\lambda_2}} dx$  diverges to  $+\infty$ . Thus it is also a contradiction of the above inequality. That is, it does not hold for  $c > 0$ .

If  $c < 0$ , for sufficiently small  $\varepsilon > 0$  ( $\varepsilon < \frac{-c}{\lambda_1\lambda_2}$ ), take

$$f(x) = \begin{cases} x^{(-\alpha-1-\lambda_1\varepsilon)/p}, & 0 < x \leq 1, \\ 0, & x > 1, \end{cases}$$

$$g(y) = \begin{cases} y^{(-\beta-1-\lambda_2\varepsilon)/q}, & 0 < y \leq 1, \\ 0, & y > 1. \end{cases}$$

Similarly, one can get

$$\int_0^1 y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} dy \int_0^1 K(t,1)t^{(-\alpha-1-\lambda_1\varepsilon)/p} dt \leq \frac{M}{\varepsilon(-\lambda_1)^{1/p}(-\lambda_2)^{1/q}}.$$

Since  $c < 0$ ,  $\lambda_1 < 0$ ,  $\lambda_2\varepsilon - \frac{c}{\lambda_1} < 0$  and  $\int_0^1 y^{-1-\lambda_2\varepsilon-\frac{c}{\lambda_1}} dy$  diverges to  $+\infty$ , which contradicts the above inequality. It does not hold for  $c < 0$ .

To sum up, when  $\lambda_1 < 0$ ,  $\lambda_2 < 0$ , we also get  $c = 0$ , that is  $\frac{\lambda_2\alpha-\lambda_1}{p} + \frac{\lambda_1\beta-\lambda_2}{q} = \lambda_1\lambda_2$ .

*Sufficiency.* If  $\frac{\lambda_2\alpha-\lambda_1}{p} + \frac{\lambda_1\beta-\lambda_2}{q} = \lambda_1\lambda_2$ , let  $a = \frac{\alpha}{pq} + \frac{1}{pq}$ ,  $b = \frac{\beta}{pq} + \frac{1}{pq}$ . By Hölder's inequality and Lemma 2, we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_0^{+\infty} K(x,y)f(x)g(y)dx dy \\ &= \int_0^{+\infty} \int_0^{+\infty} \left[ f(x) \frac{x^a}{y^b} \right] \left[ g(y) \frac{y^b}{x^a} \right] K(x,y)dx dy \\ &\leq \left( \int_0^{+\infty} \int_0^{+\infty} f^p(x) \frac{x^{ap}}{y^{bp}} K(x,y)dx dy \right)^{1/p} \\ &\quad \times \left( \int_0^{+\infty} \int_0^{+\infty} g^q(y) \frac{y^{bq}}{x^{aq}} K(x,y)dx dy \right)^{1/q} \\ &= \left( \int_0^{+\infty} x^{\frac{\alpha+1}{q}} f^p(x) \omega_1(x)dx \right)^{1/p} \left( \int_0^{+\infty} y^{\frac{\beta+1}{p}} g^q(y) \omega_2(y)dy \right)^{1/q} \\ &= \left( \int_0^{+\infty} x^{\frac{\alpha+1}{q} + \lambda_1 - \frac{\lambda_1}{\lambda_2} \left( \frac{\beta+1}{q} - 1 \right)} f^p(x) W_1 dx \right)^{1/p} \\ &\quad \times \left( \int_0^{+\infty} y^{\frac{\beta+1}{p} + \lambda_2 - \frac{\lambda_2}{\lambda_1} \left( \frac{\alpha+1}{p} - 1 \right)} g^q(y) W_2 dy \right)^{1/q} \\ &= W_1^{1/p} W_2^{1/q} \left( \int_0^{+\infty} x^{\frac{1}{\lambda_2} \left( \lambda_2\alpha - \frac{\lambda_2\alpha-\lambda_1}{p} - \frac{\lambda_1\beta-\lambda_2}{q} + \lambda_1\lambda_2 \right)} f^p(x)dx \right)^{1/p} \\ &\quad \times \left( \int_0^{+\infty} y^{\frac{1}{\lambda_1} \left( \lambda_1\beta - \frac{\lambda_2\alpha-\lambda_1}{p} - \frac{\lambda_1\beta-\lambda_2}{q} + \lambda_1\lambda_2 \right)} g^q(y)dy \right)^{1/q} \\ &= W_1^{1/p} W_2^{1/q} \left( \int_0^{+\infty} x^\alpha f^p(x)dx \right)^{1/p} \left( \int_0^{+\infty} y^\beta g^q(y)dy \right)^{1/q} \\ &= W_1^{1/p} W_2^{1/q} \|f\|_{p,\alpha} \|g\|_{q,\beta}, \end{aligned}$$

thus (4) holds when taking any constant  $M \geq W_1^{1/p} W_2^{1/q}$ .

(ii) Suppose that (4) holds. If the constant factor  $\frac{W}{|\lambda_1|^{1/q} |\lambda_2|^{1/p}}$  is not the best, then there exists a constant  $M_0 < \frac{W}{|\lambda_1|^{1/q} |\lambda_2|^{1/p}}$ , for  $f(x) \in L^p_\alpha(0, +\infty)$ ,  $g(y) \in L^q_\beta(0, +\infty)$ , such that

$$\int_0^{+\infty} \int_0^{+\infty} K(x,y)f(x)g(y)dx dy \leq M_0 \|f\|_{p,\alpha} \|g\|_{q,\beta}. \tag{8}$$

For sufficiently small  $\varepsilon > 0$  and  $\delta > 0$ , take

$$\begin{aligned} f(x) &= \begin{cases} x^{(-\alpha-1-|\lambda_1|\varepsilon)/p}, & x \geq \delta, \\ 0, & 0 < x < \delta, \end{cases} \\ g(y) &= \begin{cases} y^{(-\beta-1-|\lambda_2|\varepsilon)/q}, & y \geq 1, \\ 0, & 0 < y < 1. \end{cases} \end{aligned}$$

Then

$$\begin{aligned} \|f\|_{p,\alpha} \|g\|_{q,\beta} &= \left( \int_{\delta}^{+\infty} x^{-1-|\lambda_1|\varepsilon} dx \right)^{1/p} \left( \int_1^{+\infty} y^{-1-|\lambda_2|\varepsilon} dy \right)^{1/q} \\ &= \frac{1}{\varepsilon |\lambda_1|^{1/p} |\lambda_2|^{1/q}} \left( \frac{1}{\delta^{|\lambda_1|\varepsilon}} \right)^{1/p}. \end{aligned} \tag{9}$$

In view of  $\frac{\lambda_2\alpha-\lambda_1}{p} + \frac{\lambda_1\beta-\lambda_2}{q} = \lambda_1\lambda_2$ , we have

$$\begin{aligned} &\int_0^{+\infty} \int_0^{+\infty} K(x,y)f(x)g(y)dx dy \\ &= \int_1^{+\infty} y^{-\frac{\beta+1+|\lambda_2|\varepsilon}{q}} \left( \int_{\delta}^{+\infty} K(x,y)x^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dx \right) dy \\ &= \int_1^{+\infty} y^{\lambda_2-\frac{\beta+1+|\lambda_2|\varepsilon}{q}} \left( \int_{\delta}^{+\infty} K(xy^{-\lambda_2/\lambda_1}, 1)x^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dx \right) dy \\ &= \int_1^{+\infty} y^{\lambda_2-\frac{\beta+1+|\lambda_2|\varepsilon}{q}-\frac{\lambda_2}{\lambda_1}\frac{\alpha+1+|\lambda_1|\varepsilon}{p}+\frac{\lambda_2}{\lambda_1}} \left( \int_{\delta y^{-\lambda_2/\lambda_1}}^{+\infty} K(t,1)t^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dt \right) dy \\ &\geq \int_1^{+\infty} y^{\frac{1}{\lambda_1}\left(\lambda_1\lambda_2-\frac{\lambda_2\alpha-\lambda_1}{p}-\frac{\lambda_1\beta-\lambda_2}{q}-\lambda_1-\frac{|\lambda_1|\lambda_2\varepsilon}{p}-\frac{|\lambda_2|\lambda_1\varepsilon}{q}\right)} \\ &\quad \times \left( \int_{\delta}^{+\infty} K(t,1)t^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dt \right) dy \\ &= \int_1^{+\infty} y^{-1-|\lambda_2|\varepsilon} dy \int_{\delta}^{+\infty} K(t,1)t^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dt \\ &= \frac{1}{|\lambda_2|\varepsilon} \int_{\delta}^{+\infty} K(t,1)t^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dt, \end{aligned} \tag{10}$$

it follows from (8), (9) and (10) that

$$\frac{1}{|\lambda_2|} \int_{\delta}^{+\infty} K(t,1)t^{-\frac{\alpha+1+|\lambda_1|\varepsilon}{p}} dt \leq \frac{M_0}{|\lambda_1|^{1/p}|\lambda_2|^{1/q}},$$

consequently,

$$\frac{1}{|\lambda_2|} \int_{\delta}^{+\infty} K(t,1)t^{-\frac{\alpha+1}{p}} dt \leq \frac{M_0}{|\lambda_1|^{1/p}|\lambda_2|^{1/q}}$$

as  $\varepsilon \rightarrow 0^+$ . Let  $\delta \rightarrow 0^+$ , we get

$$\frac{W_2}{|\lambda_2|} = \frac{1}{|\lambda_2|} \int_0^{+\infty} K(t,1)t^{-\frac{\alpha+1}{p}} dt \leq \frac{M_0}{|\lambda_1|^{1/p}|\lambda_2|^{1/q}},$$

thus  $\frac{W}{|\lambda_1|^{1/q}|\lambda_2|^{1/p}} \leq M_0$ , this is a contradiction. Hence the constant factor  $\frac{W}{|\lambda_1|^{1/q}|\lambda_2|^{1/p}}$  is the best possible.  $\square$



### 3. Applications in operator theory

By using Lemma 1 and Theorem 1, we have

**THEOREM 2.** *Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 \lambda_2 > 0$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(\lambda_1, \lambda_2)$ ,  $K(1, u) (K(u, 1)) > 0$  a.e. in  $(0, 1) ((1, \infty))$ . Let the operator  $T$  be as in (2), and*

$$W_1 = \int_0^{+\infty} u^{-\frac{\beta+1}{q}} K(1, u) du, W_2 = \int_0^{+\infty} u^{-\frac{\alpha+1}{p}} K(u, 1) du,$$

are both convergent. Then

(i) the necessary and sufficient conditions for the operator  $T : L^p_\alpha(0, +\infty) \rightarrow L^p_{\beta(1-p)}(0, +\infty)$  to be bounded is  $\frac{\lambda_2 \alpha - \lambda_1}{p} + \frac{\lambda_1 \beta - \lambda_2}{q} = \lambda_1 \lambda_2$ ;

(ii) if  $T$  is bounded from  $L^p_\alpha(0, +\infty)$  to  $L^p_{\beta(1-p)}(0, +\infty)$ , then the operator norm of  $T$  is  $\|T\| = \frac{W}{|\lambda_1|^{1/q} |\lambda_2|^{1/p}}$ , where  $W = |\lambda_1| W_2 = |\lambda_2| W_1$ .

We get the following results by taking  $\alpha = \beta = 0$ .

**COROLLARY 1.** *Assume that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 < 0$ ,  $\lambda_2 < 0$ ,  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(\lambda_1, \lambda_2)$ ,  $K(1, u) (K(u, 1)) > 0$  a.e. in  $(0, 1) ((1, \infty))$ . Let the operator  $T$  be as in (2),  $W_1$  and  $W_2$  be as in Theorem 2. Then*

(i) the necessary and sufficient conditions for the operator  $T$  in  $L^p(0, +\infty)$  to be bounded is  $\frac{\lambda_1}{p} + \frac{\lambda_2}{q} + \lambda_1 \lambda_2 = 0$ .

(ii) if  $T$  is bounded in  $L^p(0, +\infty)$ , then the operator norm of  $T$  is  $\|T\| = \frac{W}{|\lambda_1|^{1/q} |\lambda_2|^{1/p}}$ , where  $W = |\lambda_2| \int_0^{+\infty} u^{-\frac{1}{q}} K(1, u) du$ .

**COROLLARY 2.** *Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ ,  $\frac{1}{\lambda_2 p} < 1$ , define the singular integral operator  $T$  by*

$$T(f)(y) = \int_0^{+\infty} \frac{1}{x^{\lambda_1} + y^{\lambda_2}} f(x) dx.$$

Then

(i)  $T$  is bounded in  $L^p(0, +\infty)$  if and only if  $\frac{\lambda_1}{p} + \frac{\lambda_2}{q} = \lambda_1 \lambda_2$ .

(ii) if  $T$  is bounded in  $L^p(0, +\infty)$ , then the operator norm of  $T$  is  $\|T\| = \frac{1}{\lambda_1^{1/q} \lambda_2^{1/p} \sin \frac{\pi}{\lambda_1 q}}$ .

*Proof.* Assume that  $K(x, y) = \frac{1}{x^{\lambda_1 + y^{\lambda_2}}}$ , then  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(-\lambda_1, -\lambda_2)$ . Since

$$\begin{aligned} \int_0^{+\infty} u^{-\frac{1}{q}} K(1, u) du &= \int_0^{+\infty} \frac{1}{1 + u^{\lambda_2}} u^{-\frac{1}{q}} du \\ &= \frac{1}{\lambda_2} \int_0^{+\infty} \frac{1}{1+t} t^{\frac{1}{\lambda_2 p} - 1} dt \\ &= \frac{1}{\lambda_2} B\left(\frac{1}{\lambda_2 p}, 1 - \frac{1}{\lambda_2 p}\right) \\ &= \frac{1}{\lambda_2} \frac{\pi}{\sin \frac{\pi}{\lambda_2 p}}, \end{aligned}$$

then according to Corollary 1, we know that Corollary 2 holds.  $\square$

**COROLLARY 3.** Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ ,  $\frac{1}{\lambda_2 p} < 1$ , define the singular integral operator  $T$  by

$$T(f)(y) = \int_0^{+\infty} \frac{1}{\max\{x^{\lambda_1}, y^{\lambda_2}\}} f(x) dx.$$

Then

(i)  $T$  is bounded in  $L^p(0, +\infty)$  if and only if  $\frac{\lambda_1}{p} + \frac{\lambda_2}{q} = \lambda_1 \lambda_2$ .

(ii) if  $T$  is bounded in  $L^p(0, +\infty)$ , then the operator norm of  $T$  is  $\|T\| = \frac{\lambda_1 q + \lambda_2 p}{\lambda_1^{1/q} \lambda_2^{1/p}}$ .

*Proof.* Set  $K(x, y) = \frac{1}{\max\{x^{\lambda_1}, y^{\lambda_2}\}}$ , then  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(-\lambda_1, -\lambda_2)$ . Since  $\lambda_2 > 0$  and  $\frac{1}{\lambda_2 p} < 1$ , then

$$\begin{aligned} \int_0^{+\infty} u^{-\frac{1}{q}} K(1, u) du &= \int_0^{+\infty} \frac{1}{\max\{1, u^{\lambda_2}\}} u^{-\frac{1}{q}} du \\ &= \frac{1}{\lambda_2} \int_0^{+\infty} \frac{1}{\max\{1, t\}} t^{\frac{1}{\lambda_2 p} - 1} dt \\ &= \frac{1}{\lambda_2} \int_0^1 t^{\frac{1}{\lambda_2 p} - 1} dt + \frac{1}{\lambda_2} \int_1^{+\infty} t^{\frac{1}{\lambda_2 p} - 2} dt \\ &= p - \frac{1}{\lambda_2} \frac{1}{\frac{1}{\lambda_2 p} - 1} = p - \frac{1}{\lambda_2} \frac{1}{-\frac{1}{\lambda_1 q}} \\ &= \frac{1}{\lambda_2} (\lambda_2 p + \lambda_1 q). \end{aligned}$$

It follows from Corollary 1 that Corollary 3 holds.  $\square$

**COROLLARY 4.** Suppose that  $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ ,  $\frac{1}{\lambda_1 q} < 1$ , define the singular integral operator  $T$  by

$$T(f)(y) = \int_0^{+\infty} \frac{\ln x^{\lambda_1} - \ln y^{\lambda_1}}{x^{\lambda_1} - y^{\lambda_2}} f(x) dx.$$

Then

(i)  $T$  is bounded in  $L^p(0, +\infty)$  if and only if  $\frac{\lambda_1}{p} + \frac{\lambda_2}{q} = \lambda_1 \lambda_2$ .

(ii) if  $T$  is bounded in  $L^p(0, +\infty)$ , then the operator norm of  $T$  is

$$\|T\| = \frac{1}{\lambda_1^{1/q} \lambda_2^{1/p}} \left( \frac{\pi}{\sin \frac{\pi}{\lambda_1 q}} \right)^2.$$

*Proof.* Set  $K(x, y) = \frac{\ln x^{\lambda_1} - \ln y^{\lambda_1}}{x^{\lambda_1} - y^{\lambda_1}}$ , then  $K(x, y)$  is a quasi-homogeneous non-negative measurable function of order  $(-\lambda_1, -\lambda_2)$ . Notice that  $\int_0^{+\infty} \frac{\ln t}{t-1} t^{\frac{1}{r}} dt = \left( \frac{\pi}{\sin \pi/r} \right)^2$  ( $r > 1$ ) (cf. [17]), we find

$$\begin{aligned} \int_0^{+\infty} u^{-\frac{1}{q}} K(1, u) du &= \int_0^{+\infty} \frac{-\ln u^{\lambda_2}}{1 - u^{\lambda_2}} u^{-\frac{1}{q}} du \\ &= \frac{1}{\lambda_2} \int_0^{+\infty} \frac{\ln t}{t-1} t^{\frac{1}{\lambda_2 p} - 1} dt \\ &= \frac{1}{\lambda_2} \int_0^{+\infty} \frac{\ln t}{t-1} t^{\frac{1}{\lambda_1 q}} dt \\ &= \frac{1}{\lambda_1} \left( \frac{\pi}{\sin \frac{\pi}{\lambda_1 q}} \right)^2. \end{aligned}$$

According to Corollary 1, we can see that Corollary 4 holds.  $\square$

#### REFERENCES

- [1] B. HE, J. F. CAO, B. C. YANG, *A brand new multiple Hilbert-type integral inequality*, Acta Mathematica Sinica, Chinese Series **58**, 4 (2015), 661–672.
- [2] Y. HONG, *A Hilbert-type integral inequality with quasi-homogeneous kernel and several functions*, Acta Mathematica Sinica, Chinese Series **57**, 5 (2014), 833–840.
- [3] I. PERIĆ, P. VUKOVIĆ, *Hardy-Hilbert's inequalities with a general homogeneous kernel*, Math. Inequal. Appl. **12** (2009), 525–536.
- [4] M. TH. RASSIAS, B. C. YANG, *On a Hardy-Hilbert-type inequality with a general homogeneous kernel*, Int. J. Nonlinear Anal. Appl. **7**, 1 (2016), 249–269.
- [5] Q. CHEN, Y. P. SHI, B. C. YANG, *A relation between two simple Hardy-Mulholland-type inequalities with parameters*, J. Inequal. Appl. **2016** (2016), 1–14.
- [6] Y. HONG, Y. M. WEN, *A necessary and sufficient condition of that Hilbert type series inequality with homogeneous kernel and the best constant factor*, Chin. Ann. Math. **37A**, 3 (2016), 329–336.
- [7] B. C. YANG, Q. CHEN, *On a more accurate Hardy-Mulholland-type inequality*, J. Inequal. Appl. **2016** (2016), 1–16.
- [8] M. Z. GAO, B. C. YANG, *On the extended Hilbert's inequality*, Proc. Amer. Math. Soc. **126**, 3 (1998), 751–759.
- [9] B. C. YANG, *On a more accurate multidimensional Hilbert-type inequality with parameters*, Math. Ineq. Appl. **18** (2015), 429–441.
- [10] D. M. XIN, B. C. YANG, Q. CHEN, *A discrete Hilbert-type inequality in the whole plane*, J. Inequal. Appl. **2016** (2016), 1–12.
- [11] J. C. KUANG, *Applied Inequalities*, Shangdong Science and Technology Press, Jinan, 2004.
- [12] B. C. YANG, Q. CHEN, *On a Hardy-Hilbert-type inequality with parameters*, J. Inequal. Appl. **2015** (2015), 1–18.

- [13] B. C. YANG, Q. CHEN, *A new extension of Hardy-Hilbert's inequality in the whole plane*, J. Funct. Space **2016** (2016), Art. ID 9197476, 1–8.
- [14] Y. HONG, *On multiple Hardy-Hilbert integral inequalities with some parameters*, J. Inequal. Appl. **2006** (2006), Art. ID 94960, 1–11.
- [15] Q. L. HUANG, B. C. YANG, *On a multiple Hilbert-type integral operator and applications*, J. Inequal. Appl. **2009** (2009), Art. ID 192197, 1–13.
- [16] B. C. YANG, Q. CHEN, *A more accurate multidimensional Hardy-Hilbert-type inequality with a general homogeneous kernel*, J. Math. Inequal. **12**, 1 (2018), 113–128.
- [17] G. H. HARDY, J. E. LITTLEWOOD, G. PÓLYA, *Inequalities*, Cambridge University Press, London, 1952.

(Received August 31, 2017)

*Yong Hong*  
*College of Mathematics and Statistics*  
*Guangdong University of Finance and Economics*  
*Guangzhou, Guangdong 510320, P. R. China*  
*e-mail: hongyonggdcc@yeah.net*

*Bing He*  
*Department of Mathematics*  
*Guangdong University of Education*  
*Guangzhou, Guangdong 510303, P. R. China*  
*e-mail: hzs314@163.com*

*Bicheng Yang*  
*Department of Mathematics*  
*Guangdong University of Education*  
*Guangzhou, Guangdong 510303, P. R. China*  
*e-mail: bcyang@gdei.edu.cn*