

SINGULAR MOSER–TRUDINGER INEQUALITY WITH THE EXACT GROWTH CONDITION IN \mathbb{R}^n

ZHAO LIU AND LU CHEN

(Communicated by J. Pečarić)

Abstract. In this paper, we establish the singular Moser-Trudinger inequality with exact growth condition. We prove that there exists a positive constant C_n such that

$$\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t(1+|u|)^{\frac{n}{n-1}}} dx \leq C_n \int_{\mathbb{R}^n} \frac{|u(x)|^n}{|x|^t} dx$$

for all $u \in W^{1,n}(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1$, where $\Phi(t) := e^t - \sum_{i=0}^{n-2} \frac{t^i}{i!}$. In order to avoid using symmetry and rearrangement, we employ the change of variables developed by Dong and Lu in [14] (see also Lam and Lu in [16]) to transform the singular Moser-Trudinger inequality with the exact growth condition to the corresponding non-singular case.

1. Introduction

Let $W_0^{1,p}(\Omega)$, $\Omega \subseteq \mathbb{R}^n$, be the usual Sobolev space, i.e, the completion of $C_c^\infty(\Omega)$ with the norm

$$\|u\|_{W_0^{1,p}(\Omega)} = \left(\int_{\Omega} |\nabla u|^p + |u|^p dx \right)^{1/p}.$$

It is well-known that

$$\begin{aligned} W_0^{1,p}(\Omega) &\subset L^{\frac{np}{n-p}}(\Omega), \quad 1 \leq p < n, \\ W_0^{1,p}(\Omega) &\subset L^\infty(\Omega), \quad p > n. \end{aligned}$$

The case $p = n$ is the limit case of these Sobolev embeddings and one can assert that $W_0^{1,n}(\Omega) \subset L^q(\Omega)$ for any $q \geq 1$. However, many examples have showed that $W_0^{1,n}(\Omega) \not\subset L^\infty(\Omega)$. In fact, it was established independently by Yudovich [36], Pohozaev [33] and Trudinger [35]. In 1971, Moser [32] proved the following inequality.

Mathematics subject classification (2010): 35A23, 42B37.

Keywords and phrases: Change of variables, Sharp constants, singular Moser-Trudinger inequality with the exact growth condition, Sobolev space.

The two authors' research were supported by the NNSF of China (No. 11371056). The first author was also partly supported by Scientific Research Fund of Jiangxi Provincial Education Department (No. GJJ160797).

THEOREM A. ([32]) *Let Ω be a domain with finite measure in \mathbb{R}^n , $n \geq 2$. Then there exists a positive constant C_n and a sharp constant $\alpha_n = n\omega_{n-1}^{\frac{1}{n-1}}$ such that*

$$\frac{1}{|\Omega|} \int_{\Omega} \exp(\alpha|u|^{\frac{n}{n-1}}) dx \leq C_n$$

for any $\alpha \leq \alpha_n$ and $u \in C_0^\infty(\Omega)$ with $\int_{\Omega} |\nabla u|^n dx \leq 1$, where ω_{n-1} is the area of the surface of the unit ball.

This result has been generalized in many directions. For instance, the singular Moser-Trudinger inequality which is an interpolation of Hardy inequality and Moser-Trudinger inequality was studied by Adimurthi and Sandeep [2]. The sharp constants for Moser-Trudinger inequalities on domains of finite measure on the Heisenberg group were established by Cohn and Lu [8] and Lam, Lu and Tang [21]. Lam, Lu and Tang [17, 20] further proved the singular Moser-Trudinger inequalities on the entire Heisenberg group. There has also been substantial progress for the Moser-Trudinger inequalities on spheres, CR spheres, or compact Riemannian manifolds, hyperbolic spaces, in Lorentz-Sobolev spaces, etc. We refer the interested reader to [5, 6, 9, 15, 23, 24, 27, 28] and the references therein. Moser-Trudinger inequalities also have many applications in geometric analysis and PDEs. Please see, for example, [10, 19, 25, 32, 34, 37] and the references therein.

When Ω has infinite volume, some versions of Moser-Trudinger type inequalities for unbounded domains were first proposed by Cao [7] when $n = 2$ and J. M. do Ó [11] for the general case $n \geq 2$. However, those inequalities are not sharp. Sharp versions of Moser-Trudinger type inequalities with best constants on unbounded domains were obtained by Adachi and Tanaka [3]. They proved that

THEOREM B. *Let $0 < \alpha < \alpha_n$, there exists a positive constant C_n such that*

$$\sup_{u \in W^{1,n}(\mathbb{R}^n), \int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1} \int_{\mathbb{R}^n} \Phi(\alpha|u(x)|^{\frac{n}{n-1}}) dx \leq C_n \int_{\mathbb{R}^n} |u(x)|^n dx,$$

where $\Phi(t) := e^t - \sum_{i=0}^{n-2} \frac{t^i}{i!}$. Moreover, the constant α_n is sharp in the sense that if $\alpha \geq \alpha_n$, the supremum will become infinite.

Ruf [34] (for the case $n = 2$), Li and Ruf [25] (for the general case $n \geq 2$) established a critical Moser-Trudinger type inequality for unbounded domains in Euclidean spaces. They obtained the following result.

THEOREM C. *There exists a positive constant C_n such that*

$$\sup_{u \in W^{1,n}(\mathbb{R}^n), \|u\|_{W^{1,n}(\mathbb{R}^n)} \leq 1} \int_{\mathbb{R}^n} \Phi(\alpha_n|u(x)|^{\frac{n}{n-1}}) dx \leq C_n.$$

Moreover, the constant α_n is sharp in the sense that if α_n is replaced by any $\alpha > \alpha_n$, the supremum will become infinite.

It has been recently shown that the critical Moser-Trudinger inequality in Theorem C and the subcritical Moser-Trudinger inequality are actually equivalent as shown by Lam, Lu and Zhang [22].

M. de Souza and J. M. Do Ó [12] studied the singular Moser-Trudinger inequality with the Dirichlet norm. They proved that

THEOREM D. *Let $0 < \beta < \alpha_n(1 - \frac{t}{n})$, there exists a positive constant C_n such that*

$$\int_{\mathbb{R}^n} \frac{\Phi(\beta |u|^{\frac{n}{n-1}})}{|x|^t} dx \leq C_n \int_{\mathbb{R}^n} \frac{|u(x)|^n}{|x|^t} dx, \quad \forall u \in W^{1,n}(\mathbb{R}^n) \text{ with } \int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1.$$

As we all know it, the above inequality does not hold if we replace β with $\alpha_n(1 - \frac{t}{n})$. So, it is natural to consider the singular Moser-Trudinger inequality with the exact growth condition. The exact growth condition for the Moser-Trudinger inequality was introduced by Masmoudi and Sani in [31] and the Moser-Trudinger inequality with the exact growth condition on hyperbolic space was established by Lu and Tang [26]. For Adams inequalities ([1]) with exact growth, we refer the reader to the works by Masmoudi and Sani [30] in dimension four and by Lu, Tang and Zhu in all dimensions $n \geq 3$ [29]. Recently, in an elegant paper of Dong and Lu in [14], they used very nicely the change of variables for non-radial functions to establish the singular Moser-Trudinger inequality. This new method has also been used by Lam and Lu in [16] to obtain sharp constants and maximizers for the Caffarelli-Kohn-Nirenberg type inequalities in a wide range of parameters. This change of variables could transform the singular inequality to the corresponding non-singular inequality. Motivated by their work, we establish the following theorem.

THEOREM 1. *For $n \geq 2$ and $0 \leq t < n$, there exists a positive constant C_n such that*

$$\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n}) |u|^{\frac{n}{n-1}})}{|x|^t (1 + |u|)^{\frac{n}{n-1}}} dx \leq C_n \int_{\mathbb{R}^n} \frac{|u(x)|^n}{|x|^t} dx, \quad \forall u \in W^{1,n}(\mathbb{R}^n), \quad \int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1. \quad (1)$$

We remark that the power $\frac{n}{n-1}$ is optimal. This can be justified by the following theorem.

THEOREM 2. *If the power $\frac{n}{n-1}$ in the denominator is replaced by any $p < \frac{n}{n-1}$, there exists a sequence of functions $\{u_k\}$ such that $\int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1$, but*

$$\left(\int_{\mathbb{R}^n} \frac{|u_k(x)|^n}{|x|^t} dx \right)^{-1} \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n}) |u_k|^{\frac{n}{n-1}})}{|x|^t (1 + |u_k|)^{\frac{n}{n-1}}} dx \rightarrow +\infty.$$

Adimurthi and Yang [4] proved that

THEOREM E. *There exists a positive constant C_n such that*

$$\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n}) |u|^{\frac{n}{n-1}})}{|x|^t} dx \leq C_n, \quad \forall u \in W^{1,n}(\mathbb{R}^n)$$

with

$$\int_{\mathbb{R}^n} |\nabla u|^n + |u|^n dx \leq 1.$$

As an application of Theorem 1, we also give some relation between Theorem E and Theorem 1.

COROLLARY 1. *Singular Moser-Trudinger inequality with the exact growth condition (Theorem 1) implies singular Moser-Trudinger inequality (Theorem E).*

This paper is organized as follows. In section 2, we use a way of change of variables to prove the singular Moser-Trudinger inequality with the exact growth condition. In section 3, we give the proof of the sharpness of the singular Moser-Trudinger inequality with the exact growth condition. In section 4, we prove some relation between singular Moser-Trudinger inequality and the singular Moser-Trudinger inequality with the exact growth condition.

2. The proof of Theorem 1

In this section, we will prove that singular Moser-Trudinger inequality with the exact growth condition. We need the following lemma.

LEMMA 1. ([31]) *Let $n \geq 2$, there exists a positive constant C_n such that*

$$\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n |u|^{\frac{n}{n-1}})}{(1+|u|)^{\frac{n}{n-1}}} dx \leq C_n \int_{\mathbb{R}^n} |u(x)|^n dx, \forall u \in W^{1,n}(\mathbb{R}^n) \text{ with } \|\nabla u\|_n \leq 1.$$

We start to prove Theorem 1. It is hard to transform the inequality (1) to the corresponding radial inequality by symmetry and rearrangement. In fact, It is very difficult to prove that

$$\frac{\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1-\frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t(1+|u|)^{\frac{n}{n-1}}} dx}{\int_{\mathbb{R}^n} \frac{|u|^n}{|x|^t} dx} \leq \frac{\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1-\frac{t}{n})|u^*|^{\frac{n}{n-1}})}{|x|^t(1+|u^*|)^{\frac{n}{n-1}}} dx}{\int_{\mathbb{R}^n} \frac{|u^*|^n}{|x|^t} dx},$$

where u^* denotes the rearrangement of u .

In the spirit of the work of Dong and Lu in [14] and Lam and Lu [16], we define a new function $v(x)$ corresponding to $u(x)$ which could keep the gradient norm less than 1, and eliminate the weights of integral at the same time.

Let $u(x) \in W^{1,n}(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} |\nabla u|^n dx \leq 1$, define the new function $v(x) \in W^{1,n}(\mathbb{R}^n)$ by the formula below,

$$v(x) = \left(\frac{n-t}{n}\right)^{\frac{n-1}{n}} u(|x|^{\frac{t}{n-t}}x).$$

Consider the vector-valued function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by $F(x) = |x|^{\frac{t}{n-t}}x$.

Direct calculations show that $\det(J_F) = \frac{n}{n-t} |x|^{\frac{nt}{n-t}}$. Then by the change of variables $y_i = |x|^{\frac{t}{n-t}}x_i$ ($1 \leq i \leq n$), we can obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |v(x)|^n dx &= \left(\frac{n-t}{n}\right)^{n-1} \int_{\mathbb{R}^n} |u(|x|^{\frac{t}{n-t}}x)|^n dx \\ &= \left(\frac{n-t}{n}\right)^n \int_{\mathbb{R}^n} \frac{|u(y)|^n}{|y|^t} dy \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}^n} |\nabla v(x)|^n dx &\leq \left(\frac{n}{n-t}\right) \int_{\mathbb{R}^n} |x|^{\frac{t}{n-t}} \left| \nabla u \left(|x|^{\frac{t}{n-t}} x \right) \right|^n dx \\ &= \int_{\mathbb{R}^n} |\nabla u(y)|^n dy. \end{aligned}$$

Then we apply Lemma 1 to derive that

$$\begin{aligned} &\int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u(y)|^{\frac{n}{n-t}})}{|y|^t(1 + |u(y)|)^{\frac{n}{n-t}}} dy \\ &= \sum_{j=n-1}^{\infty} \int_{\mathbb{R}^n} \frac{(\alpha_n(1 - \frac{t}{n})|u(y)|^{\frac{n}{n-t}})^j}{j!} \frac{1}{|y|^t(1 + |u(y)|)^{\frac{n}{n-t}}} dy \\ &= \sum_{j=n-1}^{\infty} \int_{\mathbb{R}^n} \frac{n}{n-t} \frac{(\alpha_n(1 - \frac{t}{n}))^j}{j!} |u(|x|^{\frac{t}{n-t}}x)|^{\frac{nj}{n-t}} \frac{1}{(1 + |u(|x|^{\frac{t}{n-t}}x)|)^{\frac{n}{n-t}}} dx \\ &\leq \sum_{j=n-1}^{\infty} \int_{\mathbb{R}^n} \left(\frac{n}{n-t}\right)^{j+1} \frac{(\alpha_n(1 - \frac{t}{n}))^j}{j!} |v(x)|^{\frac{nj}{n-t}} \frac{1}{(1 + |v(x)|)^{\frac{n}{n-t}}} dx \tag{2} \\ &= \frac{n}{n-t} \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n|v(x)|^{\frac{n}{n-t}})}{(1 + |v(x)|)^{\frac{n}{n-t}}} dx \\ &\leq \frac{nC_n}{n-t} \int_{\mathbb{R}^n} |v(x)|^n dx \\ &= C_n \left(\frac{n-t}{n}\right)^{n-1} \int_{\mathbb{R}^n} \frac{|u(y)|^n}{|y|^t} dy. \end{aligned}$$

Then we accomplish the proof of Theorem 1.

3. Sharpness of Theorem 1

In this section, we will prove Theorem 2. Namely, we will give the proof of the sharpness of Theorem 1. We will show that the inequality in Theorem 1 does not hold if the power $\frac{n}{n-1}$ in the denominator is replaced by any $p < \frac{n}{n-1}$.

Let $\{b_k\}_{k \geq 1} \subset \mathbb{R}^+$ and $b_k \rightarrow +\infty$ as $k \rightarrow +\infty$ and define

$$r_k := e^{-\omega_{\frac{n-1}{n}} b_k^{\frac{n}{n-1}}}.$$

For fixed $p < \frac{n}{n-1}$, we choose $\{u_k\}_{k=1}^{\infty}$ as follows:

$$u_k(x) := \begin{cases} b_k, & \text{if } |x| \leq r_k, \\ b_k \frac{|\log|x||}{|\log r_k|}, & \text{if } r_k < |x| \leq 1, \\ 0, & \text{if } |x| \geq 1. \end{cases}$$

The choice of this sequence is inspired by a similar sequence in [31].

For any $q > 0$, define

$$\tilde{u}_k(x) := u_k\left(\frac{x}{b_k}\right).$$

Then by calculation, we have

$$\int_{\mathbb{R}^n} \frac{|\tilde{u}_k(x)|^n}{|x|^t} dx \sim b_k^{(n-t)q - \frac{n}{n-1}}, \quad \int_{\mathbb{R}^n} |\nabla \tilde{u}_k|^n dx \leq 1.$$

and

$$\begin{aligned} & \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|\tilde{u}_k|^{\frac{n}{n-1}})}{|x|^t(1 + |\tilde{u}_k|)^p} dx \\ &= b_k^{(n-t)q} \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u_k|^{\frac{n}{n-1}})}{|x|^t(1 + |u_k|)^p} dx \\ &\geq b_k^{(n-t)q} \frac{\Phi(\alpha_n(1 - \frac{t}{n})b_k^{\frac{n}{n-1}})}{(1 + b_k)^p} \int_{|x| \leq r_k} \frac{1}{|x|^t} dx \\ &\geq b_k^{(n-t)q} \frac{\Phi(\alpha_n(1 - \frac{t}{n})b_k^{\frac{n}{n-1}})}{(1 + b_k)^p} e^{-\alpha_n(1 - \frac{t}{n})b_k^{\frac{n}{n-1}}}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \left(\int_{\mathbb{R}^n} \frac{|\tilde{u}_k(x)|^n}{|x|^t} dx \right)^{-1} \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|\tilde{u}_k|^{\frac{n}{n-1}})}{|x|^t(1 + |\tilde{u}_k|)^p} dx \\ &\gtrsim b_k^{\frac{n}{n-1} - p} \Phi\left(\alpha_n\left(1 - \frac{t}{n}\right)b_k^{\frac{n}{n-1}}\right) e^{-\alpha_n(1 - \frac{t}{n})b_k^{\frac{n}{n-1}}} \rightarrow \infty \end{aligned}$$

as $k \rightarrow +\infty$. So the power $\frac{n}{n-1}$ in the denominator can not be replaced by any $p < \frac{n}{n-1}$. This completes the proof of Theorem 2.

4. The proof of Corollary 1

In this section, we will show the singular Moser-Trudinger inequality with the exact growth condition implies the singular Moser-Trudinger inequality.

We will adapt an idea of using the level set of functions to derive the global inequality from the local one developed by Lam and Lu [17, 18] to our situation. We need to do some careful analysis and estimates to make this work here.

Let

$$A := \{x \in \mathbb{R}^n \mid |u(x)| \geq 1\}.$$

Since $u \in W^{1,n}(\mathbb{R}^n)$, it is obvious that A is a bounded domain. Now, we split the integral into two parts

$$\begin{aligned} & \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t} dx \\ &= \int_A \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t} dx + \int_{\mathbb{R}^n \setminus A} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t} dx. \end{aligned}$$

For $x \in \mathbb{R}^n \setminus A$, there exists a positive constant C_n such that

$$\Phi\left(\alpha_n\left(1 - \frac{t}{n}\right)|u(x)|^{\frac{n}{n-1}}\right) \leq C_n|u(x)|^n.$$

Then we have

$$\begin{aligned} & \int_{\mathbb{R}^n \setminus A} \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t} dx \\ & \leq C_n \int_{\mathbb{R}^n \setminus A} \frac{|u(x)|^n}{|x|^t} dx \\ & \leq C_n \int_{\{|u| \leq 1\} \cap \{|x| \leq 1\}} \frac{|u(x)|^n}{|x|^t} dx + C_n \int_{\{|u| \leq 1\} \cap \{|x| \geq 1\}} \frac{|u(x)|^n}{|x|^t} dx \\ & \leq C'_n. \end{aligned}$$

Since

$$\Phi^p(x) \leq \Phi(px), \quad \forall x \geq 0, \forall p \geq 1,$$

then one can apply the Hölder inequality to obtain

$$\begin{aligned} & \int_A \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t} dx \\ & = \int_A \frac{\Phi(\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})(1 + |u|)^{\frac{n}{p(n-1)}}}{|x|^{\frac{t}{p}}(1 + |u|)^{\frac{n}{p(n-1)}}|x|^{\frac{t(p-1)}{p}}} dx \\ & \leq \left(\int_A \frac{\Phi(p\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t(1 + |u|)^{\frac{n}{n-1}}} dx\right)^{\frac{1}{p}} \left(\int_A \frac{(1 + |u|)^{\frac{n}{(n-1)(p-1)}}}{|x|^t} dx\right)^{\frac{p-1}{p}} \\ & \leq \left(\int_A \frac{\Phi(p\alpha_n(1 - \frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t(1 + |u|)^{\frac{n}{n-1}}} dx\right)^{\frac{1}{p}} \left(\left(\int_A (1 + |u|)^{\frac{ns'}{(n-1)(p-1)}} dx\right)^{\frac{1}{s'}} \left(\int_A \frac{1}{|x|^{st}} dx\right)^{\frac{1}{s}}\right)^{\frac{p-1}{p}} \\ & =: I_1 \cdot I_2. \end{aligned}$$

For I_2 , we have

$$I_2 \leq \left(\int_A (1 + |u(x)|)^{\frac{ns'}{(n-1)(p-1)}} dx\right)^{\frac{(p-1)}{s'p}} \left(\int_A \frac{1}{|x|^{st}} dx\right)^{\frac{p-1}{sp}}.$$

One can choose s sufficiently close to 1 to obtain that $I_2 \leq C$.

For I_1 , let $\theta \in (0, 1)$ be such that $\|u\|_n^n = \theta$, then $\|\nabla u\|_n^n = 1 - \theta$. Let $\tilde{u} = p^{\frac{n-1}{n}}u$, we consider the following two cases:

Case 1: $\theta \geq \frac{n-1}{n}$,

Case 2: $\theta < \frac{n-1}{n}$.

For Case 1 and $1 < p < n^{\frac{1}{n-1}}$, we have

$$\|\nabla \tilde{u}\|_n^n = p^{n-1} \|\nabla u\|_n^n \leq p^{n-1}(1 - \theta) \leq \frac{p^{n-1}}{n} \leq 1.$$

For *Case 2*, let $p = \frac{n-1}{n-1-\theta}$, then $1 < p < \frac{n}{n-1} \leq n^{\frac{1}{n-1}}$. By construction, we have

$$\begin{aligned} \|\nabla \tilde{u}\|_n^n &= p^{n-1} \|\nabla u\|_n^n \leq \left(\frac{n-1}{n-1-\theta}\right)^{n-1} (1-\theta) \\ &= \left(\frac{(1-\theta)^{\frac{1}{n-1}}}{1-\frac{\theta}{n-1}}\right)^{n-1} \leq 1 \end{aligned}$$

since $(1-x)^q \leq 1-qx$ for any $x, q \in [0, 1]$. Then, we can apply singular Moser-Trudinger inequality with the exact growth condition to obtain

$$\begin{aligned} I_1 &= \left(\int_A \frac{\Phi(p\alpha_n(1-\frac{t}{n})|u|^{\frac{n}{n-1}})}{|x|^t(1+|u|)^{\frac{n}{n-1}}} dx\right)^{\frac{1}{p}} \\ &\leq \left(p \int_{\mathbb{R}^n} \frac{\Phi(\alpha_n(1-\frac{t}{n})|\tilde{u}|^{\frac{n}{n-1}})}{|x|^t(1+|\tilde{u}|)^{\frac{n}{n-1}}} dx\right)^{\frac{1}{p}} \\ &\leq C_n \left(p \int_{\mathbb{R}^n} \frac{|u(x)|^n}{|x|^t} dx\right)^{\frac{1}{p}} \\ &\leq C'_n. \end{aligned}$$

This completes the proof of Corollary 1.

We have recently learned that the authors of [13] have independently proved, among many other results, the following theorem.

THEOREM 3. *Let $0 \leq t \leq s < n$ and $0 \leq \alpha < \alpha_n$:*

$$\begin{aligned} SSTM(\alpha, s, t) &= \sup_{\|\nabla u\|_n \leq 1} \frac{1}{\left(\int_{\mathbb{R}^n} \frac{|u|^n}{|x|^t} dx\right)^{\frac{n-s}{n-t}}} \int_{\mathbb{R}^n} \frac{\Phi\left(\alpha\left(1-\frac{s}{n}\right)|u|^{\frac{n}{n-1}}\right)}{|x|^s} dx < \infty. \\ STME(s, t) &= \sup_{\|\nabla u\|_n \leq 1} \frac{1}{\left(\int_{\mathbb{R}^n} \frac{|u|^n}{|x|^t} dx\right)^{\frac{n-s}{n-t}}} \int_{\mathbb{R}^n} \frac{\phi_n\left(\alpha_n\left(1-\frac{s}{n}\right)|u|^{\frac{n}{n-1}}\right)}{\left(1+\left(1-\frac{s}{n}\right)^{\frac{n-s}{n-t}}|u|^{\frac{n}{n-1}\frac{n-s}{n-t}}\right)|x|^s} dx < \infty. \end{aligned} \tag{3}$$

The constant α_n is sharp. Moreover, the power $\frac{n}{n-1}$ in the denominator of (3) cannot be replaced by any $p < \frac{n}{n-1}$.

Moreover, $SSTM(\alpha, s, t)$, $0 < t \leq s < n$, can be attained and its extremizers are radially symmetric.

REFERENCES

- [1] D. ADAMS, *A sharp inequality of J. Moser for higher order derivatives*, Ann. of Math. **128**, 2 (1988), 385–398.
- [2] ADIMURTHI AND K. SANDEEP, *A singular Moser-Trudinger embedding and its applications*, NoDEA Nonlinear Differential Equations Appl. **13**, 5 (2007), 585–603.
- [3] S. ADACHI AND K. TANAKA, *Trudinger type inequalities in \mathbb{R}^N and their best exponents*, Proc. Amer. Math. Soc. **128**, 7 (2000), 2051–2057.
- [4] ADIMURTHI AND Y. YANG, *An interpolation of Hardy inequality and Trudinger-Moser inequality in \mathbb{R}^N and its applications*, Int. Math. Res. Not. **13** (2010), 2394–2426.
- [5] W. BECKNER, *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. **138**, 1 (1993), 213–242.
- [6] T. BRANSON, L. FONTANA AND C. MORPURGO, *Moser-Trudinger and Beckner-Onofri’s inequalities on the CR sphere*, Ann. of Math. **177**, 1 (2013), 1–52.
- [7] D. CAO, *Nontrivial solution of semilinear elliptic equation with critical exponent in \mathbb{R}^2* , Comm. Partial Differential Equations **17**, 3 (1992), 407–435.
- [8] W. S. COHN AND G. LU, *Best constants for Moser-Trudinger inequalities on the Heisenberg group*, Indiana Univ. Math. J. **50**, 4 (2001), 1567–1591.
- [9] W. S. COHN AND G. LU, *Sharp constants for Moser-Trudinger inequalities on spheres in complex space C^n* , Comm. Pure Appl. Math. **57**, 11 (2004), 1458–1493.
- [10] S. Y. A. CHANG AND P. YANG, *The inequality of Moser and Trudinger and applications to conformal geometry. Dedicated to the memory of Jürgen K. Moser*, Comm. Pure Appl. Math. **56**, 8 (2003), 1135–1150.
- [11] J. M. DO Ó, *N -Laplacian equations in \mathbb{R}^n with critical growth*, Abstr. Appl. Anal. **2**, 3 (1997), 301–315.
- [12] M. DE SOUZA AND J. M. DO Ó, *On singular Trudinger-Moser type inequalities for unbounded domains and their best exponents*, Potential Anal. **38**, 4 (2013), 1091–1101.
- [13] M. DONG, N. LAM AND G. LU, *Sharp weighted Trudinger-Moser and Caffarelli-Kohn-Nirenberg inequalities and their extremal functions*, preprint.
- [14] M. DONG AND G. LU, *Best constants and existence of maximizers for weighted Moser-Trudinger inequalities*, Calc. Var. Partial Differential Equations **55**, 4 (2016), 55–88.
- [15] L. FONTANA, *Sharp borderline Sobolev inequalities on compact Riemannian manifolds*, Comment. Math. Helv. **68**, 3 (1993), 415–454.
- [16] N. LAM AND G. LU, *Sharp constants and optimizers for a class of the Caffarelli-Kohn-Nirenberg inequalities*, arXiv:1510.01224, to appear in Advanced Nonlinear Studies.
- [17] N. LAM AND G. LU, *Sharp Moser-Trudinger inequality on the Heisenberg group at the critical case and applications*, Adv. Math. **231**, 6 (2012), 3259–3287.
- [18] N. LAM AND G. LU, *A new approach to sharp Moser-Trudinger and Adams type inequalities: a rearrangement-free argument*, J. Differential Equations **255**, 3 (2013), 298–325.
- [19] N. LAM AND G. LU, *N -Laplacian equations in \mathbb{R}^N with subcritical and critical growth without the Ambrosetti-Rabinowitz condition*, Adv. Nonlinear Stud. **13**, 2 (2013), 289–308.
- [20] N. LAM, G. LU AND H. TANG, *Sharp subcritical Moser-Trudinger inequalities on Heisenberg groups and subelliptic PDEs*, Nonlinear Anal. **95**, (2014), 77–92.
- [21] N. LAM, G. LU AND H. TANG, *On nonuniformly subelliptic equations of Q -sub-Laplacian type with critical growth in the Heisenberg group*, Adv. Nonlinear Stud. **12**, 3 (2012), 659–681.
- [22] N. LAM, G. LU AND L. ZHANG, *Equivalence of critical and subcritical sharp Trudinger-Moser-Adams inequalities*, arXiv:1504.04858, to appear in Revista Matemática Iberoamericana.
- [23] Y. X. LI, *Remarks on the extremal functions for the Moser-Trudinger inequality*, Acta Math. Sin. (Engl. Ser.) **22**, 2 (2006), 545–550.
- [24] Y. X. LI, *Moser-Trudinger inequality on compact Riemannian manifolds of dimension two*, J. Partial Differential Equations **14**, 2 (2001), 163–192.
- [25] Y. X. LI AND B. RUF, *A sharp Moser-Trudinger type inequality for unbounded domains in \mathbb{R}^n* , Indiana Univ. Math. J. **57**, 1 (2008), 451–480.
- [26] G. LU AND H. TANG, *Sharp Moser-Trudinger inequalities on hyperbolic spaces with exact growth condition*, J. Geom. Anal. **26**, 2 (2016), 837–857.

- [27] G. LU AND H. TANG, *Sharp singular Trudinger-Moser inequalities in Lorentz-Sobolev spaces*, Adv. Nonlinear Stud. **16**, 3 (2016), 581–601.
- [28] G. LU AND H. TANG, *Best constants for Moser-Trudinger inequalities on high dimensional hyperbolic spaces*, Adv. Nonlinear Stud. **13**, 4 (2013), 1035–1052.
- [29] G. LU, H. TANG AND M. ZHU, *Best constants for Adams' inequalities with the exact growth condition in \mathbb{R}^n* , Adv. Nonlinear Stud. **15**, 4 (2015), 763–788.
- [30] N. MASMOUDI AND F. SANI, *Adams' inequality with the exact growth condition in \mathbb{R}^4* , Comm. Pure Appl. Math. **67**, 8 (2014), 1307–1335.
- [31] N. MASMOUDI AND F. SANI, *Trudinger-Moser inequalities with the exact growth condition in \mathbb{R}^n and application*, Comm. Partial Differential Equations **40**, 8 (2015), 1408–1440.
- [32] J. MOSER, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J. **20**, (1970/71), 1077–1092.
- [33] S. I. POHOZAEV, *On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$* , (Russian), Dokl. Akad. Nauk SSSR. **165** (1965), 36–39.
- [34] B. RUF, *A sharp Moser-Trudinger type inequality for unbounded domains in \mathbb{R}^2* , J. Funct. Anal. **219**, 2 (2005), 340–367.
- [35] N. S. TRUDINGER, *On imbeddings into Orlicz spaces and some applications*, J. Math. Mech. **17** (1967), 473–483.
- [36] V. I. YUDOVICH, *Some estimates connected with integral operators and with solutions of elliptic equations*, (Russian), Dokl. Akad. Nauk SSSR. **138** (1961), 805–808.
- [37] J. ZHU, *Improved Moser-Trudinger inequality involving L^p norm in n -dimensions*, Adv. Nonlinear Stud. **14**, 2 (2014), 273–293.

(Received June 11, 2016)

Zhao Liu
School of Mathematics and Computer Science
Jiangxi Science and Technology Normal University
Nanchang 330038, P. R. China
e-mail: liuzhao2008tj@sina.com

Lu Chen
School of Mathematical Sciences
Beijing Normal University
Beijing 100875, P. R. China
e-mail: chenlu5818804@qq.com