

SOME NEW TWO-SIDED INEQUALITIES CONCERNING THE FOURIER TRANSFORM

AIGERIM KOPEZHANOVA, ERLAN NURSULTANOV AND LARS-ERIK PERSSON

(Communicated by I. Perić)

Abstract. The classical Hausdorff-Young and Hardy-Littlewood-Stein inequalities do not hold for $p > 2$. In this paper we prove that if we restrict to net spaces we can even derive a two-sided estimate for all $p > 1$. In particular, this result generalizes a recent result by Lifyand E. and Tikhonov S. [7] (MR 2464253)

1. Introduction

Let

$$\widehat{f}(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-itx} dx, \quad x \in \mathbb{R},$$

be the Fourier transform of a function $f \in L_1(\mathbb{R})$.

Let $1 < p < 2$, $p' = \frac{p}{p-1}$ and $0 < q \leq \infty$. Then we have the following inequalities

$$\|\widehat{f}\|_{L_{p',q}(\mathbb{R})} \leq c_1 \|f\|_{L_p(\mathbb{R})}, \quad (1)$$

$$\|\widehat{f}\|_{L_{p',q}(\mathbb{R})} \leq c_2 \|f\|_{L_{p,q}(\mathbb{R})}, \quad (2)$$

where $L_{p,q}(\mathbb{R})$ is the classical Lorentz space. These inequalities are called the Hausdorff-Young inequality and the Hardy-Littlewood-Stein inequality, respectively, (see e.g. [15] and [16]).

Note that these inequalities (1) and (2) hold with equality for $p = q = 2$ (Plancherel's theorem) but do not hold in general for $2 < p < \infty$.

Let $0 < p, q \leq \infty$, M be the set of the segments $[a, b]$ in \mathbb{R} and $|e| = b - a$.

The net space $N_{p',q}(M)$ is defined as the set of all measurable functions f such that the quasinorm

$$\|f\|_{N_{p',q}(M)} = \left(\int_0^\infty \left(t^{\frac{1}{p'}} \overline{f}(t, M) \right)^q \frac{dt}{t} \right)^{\frac{1}{q}} < \infty$$

Mathematics subject classification (2010): 46E30, 42A38.

Keywords and phrases: Inequalities, Hausdorff-Young's inequality, Hardy-Littlewood-Stein inequality, Fourier transform, Lorentz spaces, total variation, network spaces, two-sided estimates.

for $q < \infty$, and

$$\|f\|_{N_{p',\infty}(M)} = \sup_{t>0} t^{\frac{1}{p'}} \bar{f}(t, M) < \infty$$

for $q = \infty$, where

$$\bar{f}(t; M) := \sup_{\substack{|e| \geq t \\ e \in M}} \frac{1}{|e|} \left| \int_e f(x) dx \right|.$$

These spaces were introduced in [11] (see also [12] and [13]). In particular, the following result was proved:

THEOREM A. *Let $2 < p < \infty$, $0 < q \leq \infty$. Then*

$$\|\widehat{f}\|_{N_{p',q}(M)} \leq c_3 \|f\|_{L_{pq}(\mathbb{R})}. \tag{3}$$

The inequality (3) complements the Hardy-Littlewood-Stein inequality. Similar results for the Fourier transform in the periodic case were obtained in [10] and [5].

The main aims of this paper are to derive the sufficient condition so that the Fourier transform \widehat{f} belongs to L_p -space ($1 < p < \infty$) and to obtain conditions so that the norm of the Fourier transform \widehat{f} in L_p -space ($1 < p < \infty$) has both upper and lower estimates.

The main results are formulated in Section 3. The proofs can be found in Section 4 and in Section 2 we present some necessary preliminaries, including new lemmas of independent interest.

CONVENTIONS. The letter $c(c_1, c_2, etc.)$ means a constant which does not depend on the involved functions and it can be different in different occurrences. Moreover, for $A, B > 0$ the notation $A \asymp B$ means that there exists positive constants c_1 and c_2 such that $c_1 A \leq B \leq c_2 A$. For $1 < p < \infty$ we denote $p' = p/(p - 1)$.

2. Preliminaries

The total variation of the function f , defined on an interval $[a, b] \subset \mathbb{R}$ is the quantity

$$V_a^b(f) := \sup_{\mathfrak{P}} \sum_{i=0}^n |f(x_{i+1}) - f(x_i)|,$$

where the supremum is taken over all partitions of $[a, b]$:

$$\mathfrak{P}: a = x_0 < x_1 < \dots < x_n = b, \quad n \in \mathbb{Z}_+.$$

We say that the measurable function $f(x) \in V([a, b])$ if $V_a^b(f) < \infty$.

The total variations $V_a^\infty(f)$ and $V_{-\infty}^b(f)$ can be defined as follows:

$$V_a^\infty(f) := \lim_{b \rightarrow \infty} V_a^b(f)$$

and

$$V_{-\infty}^b(f) := \lim_{a \rightarrow -\infty} V_a^b(f).$$

We need the following lemma (see [3]).

LEMMA 1. Let $f(x)$ be a continuous function on a closed interval $[a, b]$ and $g = g(x) \in V([a, b])$. Then the integral

$$I = \int_a^b f(x)dg(x),$$

exists and

$$|I| \leq \max_{x \in [a, b]} |f(x)|V_a^b(g).$$

We also need the following lemmas of independent interest:

LEMMA 2. Let $1 < p < \infty$. Then

$$\|\widehat{f}\|_{N_{p',p}(M)} \leq c\|f\|_{L_p(\mathbb{R})}.$$

Proof. The statement of this Lemma 2 follows from Theorem A for the case $2 < p < \infty$. Moreover, taking into account that

$$\|\widehat{f}\|_{N_{p',p}(M)} \leq c_1\|\widehat{f}\|_{L_{p',p}(\mathbb{R})},$$

(see [13]) the statement of this Lemma 2 follows from (2) for the case $1 < p \leq 2$. The proof is complete. \square

Let $V_{2^k}(f) := V_{[2^k, 2^{k+1}] \cup [-2^{k+1}, -2^k]}(f)$ be the total variation of the function $f(x)$, defined on the set $[2^k, 2^{k+1}] \cup [-2^{k+1}, -2^k]$, $k \in \mathbb{Z}$.

LEMMA 3. Let $\alpha > 0$ and $1 < p < \infty$. If

$$\left(\sum_{k \in \mathbb{Z}} \left(2^{k\alpha} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}} < \infty,$$

then

$$V_1^\infty(f) < \infty \quad \text{and} \quad V_{-\infty}^{-1}(f) < \infty.$$

Proof. It is obvious that

$$V_1^\infty(f) = \sum_{k=0}^\infty V_{2^k}^{2^{k+1}}(f).$$

By using Hölder’s inequality, we obtain that

$$\begin{aligned} \sum_{k=0}^\infty V_{2^k}^{2^{k+1}}(f) &\leq \left(\sum_{k=0}^\infty \left(2^{k\alpha} V_{2^k}^{2^{k+1}}(f) \right)^p \right)^{\frac{1}{p}} \cdot \left(\sum_{k=0}^\infty 2^{-k\alpha \cdot p'} \right)^{\frac{1}{p'}} \\ &= c_{\alpha,p} \left(\sum_{k=0}^\infty \left(2^{k\alpha} V_{2^k}^{2^{k+1}}(f) \right)^p \right)^{\frac{1}{p}} \leq c_{\alpha,p} \left(\sum_{k=0}^\infty \left(2^{k\alpha} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}} < \infty. \end{aligned}$$

The second inequality for $V_{-\infty}^{-1}(f)$ can be proved in a similar way. The proof is complete. \square

LEMMA 4. Let $1 < p < \infty$ and $0 < q \leq \infty$. If $f \in N_{pq}(M)$, then the following equivalence

$$\|f\|_{N_{pq}} \asymp \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p}} \bar{f}(2^k, M) \right)^q \right)^{\frac{1}{q}}$$

holds, where $\bar{f}(2^k, M) = \sup_{\substack{|e| \geq 2^k \\ e \in M}} \frac{1}{|e|} \left| \int_e f(x) dx \right|$, M is the set of all segments $[a, b]$ in \mathbb{R} .

Proof. Since $f \in N_{pq}(M)$ we have that

$$I = \int_0^\infty \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t} < \infty.$$

This integral can be represented as follows

$$I = \sum_{k \in \mathbb{Z}} \int_{2^k}^{2^{k+1}} \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t}.$$

Taking into account that the function \bar{f} is monotone, we find that

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \int_{2^k}^{2^{k+1}} \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t} &\leq \sum_{k \in \mathbb{Z}} \left(2^{\frac{k+1}{p}} \bar{f}(2^k, M) \right)^q \int_{2^k}^{2^{k+1}} \frac{dt}{t} \\ &= c_1 \sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p}} \bar{f}(2^k, M) \right)^q, \end{aligned}$$

and

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \int_{2^k}^{2^{k+1}} \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t} &\geq 2^{-\frac{1}{p}} \sum_{k \in \mathbb{Z}} \left(2^{\frac{k+1}{p}} \bar{f}(2^{k+1}, M) \right)^q \int_{2^k}^{2^{k+1}} \frac{dt}{t} \\ &= c_2 \sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p}} \bar{f}(2^k, M) \right)^q. \end{aligned}$$

The proof is complete. \square

The statement in our Theorem 2 is related to a recent result by E. Liflyand and S. Tikhonov [7], where an extended solution of Boas' conjecture was proved (for original proof see [14]). In particular, they defined the class GM as follows:

DEFINITION 1. We say that the function f belongs to the class GM if for all $x \in (0, \infty)$ we have that

$$V_x^{2x}(f) \leq c \int_{\frac{x}{\beta}}^{\beta x} t^{-1} |f(t)| dt, \tag{4}$$

for some $\beta > 1$.

3. Main results

Our first main result reads as follows.

THEOREM 1. *Let $1 < p < \infty$ and $f \in L_1(\mathbb{R})$. If f satisfies the condition*

$$\left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}} < \infty, \tag{5}$$

then $\widehat{f} \in L_p(\mathbb{R})$ and the inequality

$$\|\widehat{f}\|_{L_p} \leq c \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}} \tag{6}$$

holds. Here the constant c does not depend on f .

In [2] estimates of the form (6) in terms of the Fourier coefficients are obtained but these estimates are obtained with additional conditions such as GM monotonicity and non-negativity of the Fourier coefficients.

Our next main result is

THEOREM 2. *Let $1 < p < \infty$. Assume that the function f satisfy that there exists $c > 0$ such that*

$$V_{2^k}(f) \leq c \sup_{\substack{|e| \geq 2^k \\ e \in M}} \frac{1}{|e|} \left| \int_e f(x) dx \right|, \quad k \in \mathbb{Z}. \tag{7}$$

Then $\|\widehat{f}\|_{L_p(\mathbb{R})} < \infty$ if and only if $\|f\|_{N_{p'/p}} < \infty$ and, moreover,

$$\|\widehat{f}\|_{L_p(\mathbb{R})} \asymp \|f\|_{N_{p'/p}(M)}.$$

REMARK 1. Necessary and sufficient conditions on the Fourier transform \widehat{f} for nonnegative functions from the class GM to belong to the space $L_p(\mathbb{R})$ ($1 < p < \infty$) was proved in [7] (see also e.g. [8]). In this connection we also refer to [17]. The following proposition shows that Theorem 2 in a sense generalizes the mentioned result.

PROPOSITION 1. (i) *If f is a non-negative function and $f \in GM$, then f satisfies condition (7).*

(ii) *The reversed implication does not hold in general, more precisely, there exists a function f satisfying (7) but not (4).*

REMARK 2. In connection to statement (i) we also refer to M. Dyachenko, E. Liflyand and S. Tikhonov [1]. Note that in [1] criteria of belonging of the cosine and the sine Fourier transforms in the Lebesgue spaces with the power weights are obtained for non-negative functions $f \in GM$.

REMARK 3. The class of GM functions was important in the study both of Fourier series and Fourier transforms. In connection to the results obtained above we also refer to the recent papers [6] and [9] by E. Liflyand and S. Tikhonov. The first one surveys GM functions, while in the second one more general weighted estimates are obtained than in [7]. Moreover, these results gave rise to some multivariate extensions, see [4].

The proofs of the statements in Theorem 1, Theorem 2 and Proposition 1 are given in the next Section.

4. Proofs of the main results

Proof of Theorem 1. By using Lemma 3, we find that $V_1^\infty(f) < \infty$ and $V_{-\infty}^{-1}(f) < \infty$. Then, for all sequences such that $x_k \rightarrow \infty$ we have that each sequence $\{f(x_k)\}$ is a fundamental sequence (satisfying (8) below). Indeed, since $V_1^\infty(f) < \infty$, then there exists N such that for all $k > N$ we have $V_{x_k}^\infty(f) < \varepsilon$. Hence,

$$|f(x_k) - f(x_{k+p})| \leq \sum_{j=k}^{k+p-1} |f(x_j) - f(x_{j+1})| = V_{x_k}^{x_{k+p}}(f) \leq V_{x_k}^\infty(f) < \varepsilon. \tag{8}$$

Thus, we have that $\lim_{k \rightarrow \infty} f(x_k)$ exists. Since the sequence $\{x_k\}$ is arbitrarily chosen we obtain that $\lim_{x \rightarrow +\infty} f(x) = a$. Therefore, due to the fact that $f \in L_1(\mathbb{R})$, we conclude that $\lim_{x \rightarrow +\infty} f(x) = 0$.

Let $x > 0$. Then

$$\int_x^{+\infty} df(y) = \lim_{b \rightarrow +\infty} (f(b) - f(x)) = -f(x).$$

By applying the duality representation of the norm of a function in the space $L_p(\mathbb{R})$, we obtain that

$$\|\widehat{f}\|_{L_p(\mathbb{R})} = \sup_{\|g\|_{L_{p'}(\mathbb{R})}=1-\infty} \int \widehat{f(x)} \overline{\widehat{g(x)}} dx = \sup_{\|g\|_{L_{p'}(\mathbb{R})}=1-\infty} \int f(x) \overline{\widehat{\widehat{g(x)}}} dx,$$

where $\overline{\widehat{\widehat{g(x)}}}$ is the conjugate function of $\widehat{g(x)}$.

This integral can be represented as follows

$$\|\widehat{f}\|_{L_p(\mathbb{R})} = \sup_{\|g\|_{L_{p'}(\mathbb{R})}=1} \left(\left| \int_{-\infty}^0 f(x) \overline{\widehat{\widehat{g(x)}}} dx \right| + \left| \int_0^\infty f(x) \overline{\widehat{\widehat{g(x)}}} dx \right| \right).$$

By using the fact that $f(x) = -\int_x^\infty df(y)$ we can consider the following integral

$$I := \left| \int_0^\infty f(x) \overline{\widehat{\widehat{g(x)}}} dx \right| = \left| \int_0^\infty \left(\int_x^\infty df(y) \right) \overline{\widehat{\widehat{g(x)}}} dx \right|.$$

Hence, by interchanging the order of integration and using Lemma 1, we find that

$$\begin{aligned} I &= \left| \int_0^\infty \int_0^y \overline{g(x)} dx df(y) \right| \\ &= \left| \sum_{k \in \mathbb{Z}} \int_{2^k}^{2^{k+1}} \int_0^y \overline{g(x)} dx df(y) \right| \\ &\leq \sum_{k \in \mathbb{Z}} \sup_{2^k \leq y \leq 2^{k+1}} \left| \int_0^y \widehat{g(x)} dx \right| \cdot V_{2^k}^{2^{k+1}}(f). \end{aligned}$$

Furthermore, taking into account that

$$\left| \int_0^y \widehat{g(x)} dx \right| \leq 2^{k+1} \sup_{\substack{|e| \geq 2^k \\ e \in M}} \frac{1}{|e|} \left| \int_e \widehat{g(x)} dx \right| = 2^{k+1} \overline{(\widehat{g})}(2^k, M) \text{ for } 2^k \leq y \leq 2^{k+1},$$

where $\overline{(\widehat{g})}(2^k; M)$ is the average function of \widehat{g} on the set M of the segments $[a, b]$, we obtain that

$$I \leq \sum_{k \in \mathbb{Z}} 2^{k+1} \overline{(\widehat{g})}(2^k, M) V_{2^k}^{2^{k+1}}(f) = c_1 \cdot \sum_{k \in \mathbb{Z}} 2^{\frac{k}{p}} \overline{(\widehat{g})}(2^k, M) V_{2^k}^{2^{k+1}}(f) 2^{\frac{k}{p'}}.$$

Next, by using Hölder’s inequality, we get that

$$I \leq c_1 \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p}} \overline{(\widehat{g})}(2^k, M) \right)^{p'} \right)^{\frac{1}{p'}} \cdot \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p}} V_{2^k}^{2^{k+1}}(f) \right)^p \right)^{\frac{1}{p}}.$$

Hence, by using Lemma 4 and Lemma 2, we obtain the following estimate:

$$I \leq c_2 \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{2^k}^{2^{k+1}}(f) \right)^p \right)^{\frac{1}{p}} \|g\|_{L_{p'}(\mathbb{R})}. \tag{9}$$

Similarly, we can estimate the integral $\int_{-\infty}^0 f(x) \overline{g(x)} dx$:

$$\left| \int_{-\infty}^0 f(x) \overline{g(x)} dx \right| \leq c_3 \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{-2^{k+1}}^{-2^k}(f) \right)^p \right)^{\frac{1}{p}} \|g\|_{L_{p'}(\mathbb{R})}. \tag{10}$$

By combining (9) and (10), we find that

$$\|\widehat{f}\|_{L_p(\mathbb{R})} \leq c \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}}.$$

The proof is complete. \square

Proof of Theorem 2. By using Lemma 2, we have that

$$\|f\|_{N_{p',p}(M)} \leq c_2 \|\widehat{f}\|_{L_p(\mathbb{R})}.$$

On the other hand, by using inequality (6) of Theorem 1, we find that

$$\|\widehat{f}\|_{L_p(\mathbb{R})} \leq c_3 \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} V_{2^k}(f) \right)^p \right)^{\frac{1}{p}}.$$

Therefore, in view of the fact that

$$V_{2^k}(f) \leq c \sup_{\substack{|e| \geq 2^k \\ e \in M}} \frac{1}{|e|} \left| \int_e f(x) dx \right|, \quad k \in \mathbb{Z},$$

it yields that

$$\|\widehat{f}\|_{L_p(\mathbb{R})} \leq c_4 \left(\sum_{k \in \mathbb{Z}} \left(2^{\frac{k}{p'}} \sup_{\substack{|e| \geq 2^k \\ e \in M}} \frac{1}{|e|} \left| \int_e f(x) dx \right| \right)^p \right)^{\frac{1}{p}} = c_4 \|f\|_{N_{p',p}(M)}.$$

The proof is complete. \square

Proof of Proposition 1. (i) Let f be a non-negative function and $f \in GM$. Then

$$\begin{aligned} V_{2^k}(f) &= V_{-2^{k+1}}^{-2^k}(f) + V_{2^k}^{2^{k+1}}(f) \leq c \left(\int_{-\frac{2^k}{\beta}}^{-\beta 2^k} t^{-1} |f(t)| dt + \int_{\frac{2^k}{\beta}}^{\beta 2^k} t^{-1} |f(t)| dt \right) \\ &\leq c \left(\int_{-\frac{2^k}{\beta^*}}^{-\beta^* 2^k} t^{-1} |f(t)| dt + \int_{\frac{2^k}{\beta^*}}^{\beta^* 2^k} t^{-1} |f(t)| dt \right), \end{aligned}$$

where $\beta^* = \max\{2, \beta\}$. Therefore,

$$\begin{aligned} V_{2^k}(f) &\leq c \left(\frac{\beta^*}{2^k} \int_{-\beta^* 2^k}^{-\frac{2^k}{\beta^*}} f(t) dt + \frac{\beta^*}{2^k} \int_{\frac{2^k}{\beta^*}}^{\beta^* 2^k} f(t) dt \right) \\ &\leq 2c(\beta^{*2} - 1) \sup_{|e| \geq 2^k} \frac{1}{|e|} \int_e f(x) dx, \end{aligned}$$

i.e. f satisfies (7).

(ii) Let

$$f(x) = \begin{cases} 1, & -1 \leq x \leq 1, \\ \frac{\sin x}{x}, & 1 < x, \\ \frac{1}{x^2}, & x < -1. \end{cases}$$

We note that

$$V_{2^k}(f) \asymp \begin{cases} 2^{-k}, & k \in \mathbb{Z}^+ \\ 0, & -k \in \mathbb{N}, \end{cases}$$

$$\sup_{|e| \geq 2^k} \frac{1}{|e|} \int_e f(x) dx \asymp \begin{cases} 2^{-k}, & k \in \mathbb{Z}^+ \\ 1, & -k \in \mathbb{N}, \end{cases}$$

so f satisfies (7). On the other hand when $k \in \mathbb{N}$, we have

$$V_{2^k}^{2^{k+1}}(f) \asymp 2^{-k},$$

$$\int_{\frac{2^k}{\beta}}^{\beta 2^k} t^{-1} |f(t)| dt \asymp 2^{-2k},$$

which means that f does not satisfy (4). The proof is complete. \square

Acknowledgement. The authors thank the careful referee for some generous suggestions, which have improved the final version of this paper.

The authors thank Luleå University of Technology for financial support for the research visit of the first author in December 2015. The research of the first author was also supported by the MESRK grants 5540/GF4 and the research of the second author was supported by the MESRK grants 4080/GF4.

The publication was supported by the Ministry of Education and Science of the Russian Federation (the Agreement number 02.a03.21.0008).

REFERENCES

- [1] M. DYACHENKO, E. LIFLYAND AND S. TIKHONOV, *Uniform convergence and integrability of Fourier integrals*, J. Math. Anal. Appl., **372** (2010), 328–338.
- [2] M. DYACHENKO AND S. TIKHONOV, *A Hardy-Littlewood theorem for multiple series*, J. Math. Anal. Appl., **339**, 1 (2008), 503–510.
- [3] M. I. D'YACHENKO AND P. L. ULYANOV, *Measure and integral*, Faktorial, Moscow, 1998, 160 pp. (in Russian).
- [4] D. GORBACHEV, E. LIFLYAND AND S. TIKHONOV, *Weighted Fourier inequalities: Boas' conjecture in \mathbb{R}^n* , J. Anal. Math., **114** (2011), 99–120.
- [5] A. KOPEZHANOVA AND L.-E. PERSSON, *On summability of the Fourier coefficients in bounded orthonormal systems for functions from some Lorentz type spaces*, Eurasian Math. J., **1**, 2 (2010), 76–85.
- [6] E. LIFLYAND AND S. TIKHONOV, *A concept of general monotonicity and applications*, Math. Nachr., **284**, 8–9 (2011), 1083–1098.
- [7] E. LIFLYAND AND S. TIKHONOV, *Extended solution of Boas' conjecture on Fourier transforms*, C. R. Acad. Sci. Paris. Ser. I., **346** (2008), 1137–1142.
- [8] E. LIFLYAND AND S. TIKHONOV, *The Fourier transforms of general monotone functions*, Analysis and Mathematical Physics, Trends in Mathematics, Birkhäuser, (2009), 377–395.
- [9] E. LIFLYAND AND S. TIKHONOV, *Two-sided weighted Fourier inequalities*, Ann. Sc. Norm. Super. Pisa Cl. Sci., **XI**, 5 (2012), 341–362.
- [10] E. D. NURSULTANOV, *On the coefficients of multiple Fourier series from L_p -spaces*, Izv. Russ. Akad. Nauk Ser. Mat., **64**, 1 (2000), 95–122. (in Russian). Translation in: Izv. Math., **64**, 1 (2000), 93–120.
- [11] E. D. NURSULTANOV, *Network space and Fourier transform*, Dokl. Russ. Akad. Nauk., **361**, 5 (1998), 597–599 (in Russian). Translation in: Acad. Sci. Dokl. Math., **58**, 1 (1998), 105–107.

- [12] E. D. NURSULTANOV, *Network spaces and inequalities of Hardy-Littlewood type*, Mat. Sb., **189**, 3 (1998), 83–102. (in Russian). Translation in: Sb. Math., **189**, 3 (1998), 399–419.
- [13] E. D. NURSULTANOV AND S. TIKHONOV, *Net Spaces and Boundedness of Integral Operators*, J. Geom. Anal., **21**, 4 (2011), 950–981.
- [14] Y. SAGHER, *Integrability conditions for the Fourier transform*, J. Math. Anal. Appl., **54** (1976), 151–156.
- [15] E. M. STEIN, *Interpolation of linear operators*, Trans. Amer. Math. Soc., **83** (1956), 482–492.
- [16] E. M. STEIN AND G. WEISS, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, 1972.
- [17] S. TIKHONOV, *Trigonometric series with general monotone coefficients*, J. Math. Anal. Appl., **326** (2007), 721–735.

(Received June 22, 2016)

Aigerim Kopezhanova
Department of Engineering Sciences and Mathematics Luleå
University of Technology
SE 97187, Luleå, Sweden
and
Faculty of Mechanics and Mathematics L. N. Gumilyov
Eurasian National University
Satpayev st., 2, 010008 Astana, Kazakhstan
e-mail: Kopezhanova@mail.ru

Erlan Nursultanov
Kazakhstan Branch of Lomonosov Moscow State University
Astana, Kazhymukan st., 11, 010010 Astana, Kazakhstan
e-mail: er-nurs@yandex.ru

Lars-Erik Persson
Department of Engineering Sciences and Mathematics Luleå
University of Technology
SE 97187, Luleå, Sweden
and
UiT, The Arctic University of Norway
P. O. Box 385, N 8505, Narvik, Norway
and
RUDN University
6 Miklukho-Maklay St
Moscow, 117198, Russia
e-mail: Lars-Erik.Persson@ltu.se